

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (TS-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.
655-776

DATE OF ISSUANCE
SEP 18 1985

NOTICE OF PESTICIDE

REGISTRATION
REREGISTRATION

(Under the Federal Insecticide, Fungicide
and Rodenticide Act, as amended)

TERM OF ISSUANCE

Until Reregistration

NAME OF PESTICIDE PRODUCT

Prentox Diazinon Turf Insecticide

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Prentiss Drug & Chemical Co., Inc.
C.R. 2000
Floral Park, NY 11001

190 / 149146
17 / 1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling change listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 655-776."
 - b. Change "Diazinon AG500 contains 4 lbs of Diazinon per gallon" to "Prentox Diazinon Turf Insecticide contains, etc."
 - c. Revise the first paragraph under Environmental Hazards to be in keeping with the use pattern on this (split) label, i.e., revise as follows:

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

9/18/85

- d. Change "agricultural crops" in GENERAL INFORMATION to "turf and ornamentals." Also delete the statements "Do not use in barns or dwellings. Do not use on livestock."
 - e. Change "Diazinon AG500" in the table of dilution to "this product."
 - f. This label does not appear to come under the requirements of PR Notice 83-2 unless you have determined that this product would involve any other crop activities that meet the definitions for hand labor tasks normally performed by farmworkers. Therefore you may delete the statements beginning "Do not apply this product in such a manner . . .," and "Do not enter treated areas"
 - g. The Confidential Statement of Formula is acceptable and has been made a part of the registration record for this product.
 - h. We suggest you revise your product name to "Prentox Diazinon Turf and Ornamental Insecticide."
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

Apples, Pears—Apply 1 qt per acre. Do not apply within 14 days of harvest.
Borers—Use a minimum of 1 gal of water for aerial applications and 5 gals of water for ground applications. Sorghum Midge Apply 1/2 pt per acre when 50% of heads have emerged from the boot. Repeat application every 7 days. **Beetles** Apply 1 pt per acre when infestation first appears and repeat applications as necessary. **Greenbug** Apply at the rates indicated below when infestation first appears and repeat as necessary. Texas and Oklahoma—Apply 1/2 pt per acre. All Other Areas—Apply 1 pt per acre. Crops may be fed to livestock immediately following treatment. Do not apply within 7 days of harvesting for grain. **Sugar Beets**—Leafhoppers, Dicotyledon Leaf Miners, Aphids Apply 1 pt per acre. NOTE: Tops may be fed to beef and dairy animals. **Sugarcane**—Aphids Apply 1/2 pt per acre. Allow 1 day between last application and harvest. **Bagasse and forage** may be fed to livestock. **Tobacco**—Aphids, Flea Beetles Apply 1/2 pt per acre. Do not apply within 3 days of harvest. **Grasshoppers** Apply 1 pt per acre. Do not apply within 14 days of harvest.

ORNAMENTAL INSECT CONTROL
 To control certain insects on Arborvitae, Azalea, Birch, Boxwood, Camellia, Carnation, Chrysanthemum, Douglas Fir, Elm, Gladioli, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Oak, Pine, Plum, Poplar, Rhododendron, Rose, Spruce, and Willow, apply the recommended rates indicated below.

Insects	Rate per 3 gals. water	Rate per 100 gal. water
Aphids, Bagworms, Carnation Bud Mites, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, Dicotyledon Leaf Miners, European Pine Shoot Moths, European Red Mites, Flea Beetles, Holly Bud Moths, Leafhoppers, Obscure Root Weevils, Omnivorous Leaf Thrips, Privet Mites, Scale Crawlers (Colony-Cushion, Lecanium, Pine Needle San Jose Soft Scale), Trips, Two-Spotted Mite Webworms, Whiteflies	1/2 pt	1 pt
Apple-and-Thorn Skeletonizers, Cottonweaver Webworms, Fall Webworms, Hemlock Cherries, Oak Looper, Oblique-banded Leaf Roller, Pear Slugs, Tent Caterpillars	1 1/2 fl oz	3 pts

LAWN INSECT CONTROL
 To control insects listed below, apply recommended rates. For control of Ants, Brown Dog Ticks, Clover Mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, and Colembola (Springtails), which are commonly found near the house and may enter the house, spray a 5-ft. band of soil around the house, as well as the house foundation wall to a height of 2-3 ft. Repeat application if necessary.

Insects	Rate/1,000 sq. ft. Diazinon Water AG500	Remarks
Lawn Chinch Bugs	2-3 fl oz or 3-6 fl oz	Use higher rate for longer residual control and in lawns of dense growth such as St. Augustine lawns.
Ants, Armyworms, Clover Mites, Colembola (Springtails), Crickets, Cutworms, Digger Wasps, Earwigs, Frit flies, Lawn Billbugs, Sod Webworms (Lawn Moths), Sowbugs	1/2 fl oz	Spot spray ant hills and wasp ground nest openings. For lawn billbugs, irrigate the lawn thoroughly immediately after treatment. For Frit flies, mow grass and water well before treatment but delay application until grass is dry. Do not water again until necessary.
Brown Dog Ticks, Bermuda Mites, Chiggers, Fleas, Leafhoppers	1 1/2 fl oz	Do not apply to animals. For brown dog ticks spray grass and under shrubbery particularly near house.
Millipedes	8 fl oz	
Rhodes-grass Scales	5 fl oz	Apply when crawlers first emerge.

* Alternately, make application in 3 gals of water per 1,000 sq. ft. and thoroughly water immediately following application. Boxelder Bugs Spray bugs outside the home—in 1 1/2-2 1/2 fl oz (2 1/2-3 tablespoons) in 1 gal water. At the same time thoroughly spray around foundation, windows, doors, tree trunks, and in to cracks and other places where the insects hide. **Billbugs and White Grubs**, Japanese beetle larvae, European chaffer and Southern chaffer—Apply 3 fl oz of product per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass is present, indicating

inches around each mound. Apply gently to avoid disturbing the ants. Repeat as necessary. Larger quantities of solution can be used for mound treatments. Use equipment capable of applying the product at a gentle rain at the rate of 1 gal per mound. The pressure of spray may disturb the ants and cause migration, reducing the effectiveness. NOTE: For best results, apply in cool weather, 65° to 85° F., in early morning or late evening hours. Treat new mounds as they appear.

FIELD AND FORAGE SOIL INSECTS
 Unless otherwise stated, soil applications of Diazinon AG500 should always be applied to field and forage crops as a spray in a minimum of 50 gal of water per acre for ground applications. **A Half and Over**—Cutworms (Surface and Subterranean) Broadcast 2-4 qts per acre just prior to planting. Work into the soil immediately 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. **Corn**—Corn Rootworm Larvae Broadcast Application at Planting Time Apply 1-2 qts per acre. Make application in a 4-inch band directly behind the planter shoe. For band treatment refer to the row spacing chart for amounts recommended per acre. **Postemergence Basal Treatment** Apply 1 1/2-2 1/2 oz per 1,000 linear feet of corn row in sufficient water to thoroughly wet base of corn plant and a narrow band of soil next to the plant. Refer to the row spacing chart for amounts recommended per acre. Cover treated band with 2-3 inches of soil using a disk hiller or cultivator immediately after spraying. Application should be made at the first sign of corn rootworm larvae, but not later than June 24th in the Middle Corn Belt, or July 1st in the Northern Corn Belt.

SUGGESTED RATES ACCORDING TO ROW SPACING FOR CORN ROOTWORM LARVAE CONTROL

Row Spacing	Linear feet of Corn row/acre	Rate per acre Planting time application	Postemergence basal treatment
48"	10,890	28-56 oz	14-28 oz
42"	12,455	31-62 oz	15.5-31 oz
40"	13,068	32-64 oz (1-2 qts)	16-32 oz (1/2-1 qt)
38"	13,939	35-70 oz	17.5-35 oz
36"	14,250	36-72 oz	18-36 oz
30"	17,424	44-88 oz	22-44 oz
24"	21,780	55-110 oz	27.5-55 oz
20"	26,136	66-132 oz (2-4 qts)	33-66 oz (1 qt)
15"	34,848	88-176 oz	44-88 oz

NOTE: For row spacings not listed, calculate the appropriate dosage based on the 40" row figure.

Seed Corn Maggots Broadcast 2-4 qts per acre just prior to planting. Work into soil 2 inches immediately. **Cutworms (Surface and Subterranean)** Broadcast 2-4 qts per acre. For surface cutworms apply just prior to planting or when larvae first appear and work into soil 1-2 inches immediately. Do not apply later than July 1st. For subterranean cutworms apply just prior to planting and work into soil 3-6 inches immediately. **Wireworms** Apply 3-4 qts per acre just prior to planting and work into the soil 4-8 inches immediately. Corn may be fed to dairy and beef cattle and sheep. **Cowpeas**—Cutworms (Surface and Subterranean) Broadcast 2-4 qts per acre just prior to planting. Work into the soil immediately 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. **Lespedeza**—Cutworms (Surface and Subterranean) Broadcast 2-4 qts per acre just prior to planting. Work into the soil immediately 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. **Peanuts**—Cutworms Apply 2 qts per acre in a 12-inch band over the furrow. Thorough coverage of the foliage is essential. Begin application when small larvae appear. Two or three additional applications at three-week intervals may be necessary. Do not apply within 21 days of peanut harvest. **Livestock** (1) may be fed peanut hulls (2) may be fed peanut forage 7 days following treatment and (3) may be fed peanut hay 21 days following treatment. **Sorghum**—Cutworms (Surface and Subterranean) Broadcast 2-4 qts per acre just prior to planting. Work into the soil immediately 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. **Soybeans**—Cutworms (Surface and Subterranean) Broadcast 2-4 qts per acre just prior to planting. Work into the soil immediately 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. **Sugarcane**—Wireworms Broadcast 3-4 qts per acre just prior to planting and work into the soil 4-8 inches immediately. **Tops** may be fed to beef and dairy animals. **Sugarcane**—Wireworms Apply 1 gal per acre in a 12-inch band centered over the furrow. Make application directly to seed pieces in open furrow at covering time. **Tobacco**—Cutworms Broadcast 2-3 qts per acre before transplanting and work into the soil 2-4 inches immediately. **Green June Beetle Larvae** Mix 1/2 pt in 50 gals of water and apply as a drench (100 gals per 100 sq. yds.) to uprooted areas of plant beds with sprinkling can. **Koala Crickets** Broadcast 1 qt per acre when insects are present. Make application to beds or fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches immediately. **Seed-Corn Maggots** Broadcast 3 pts per acre 3-4 weeks before transplanting. Work into soil 2 inches immediately. **Wireworms** Broadcast 2-3 qts per acre 3-4 weeks before transplanting. Work into soil 3-6 inches immediately.

RANGE, PASTURE, AND GRASSLAND INSECTS
 Unless otherwise stated, Diazinon AG500 should always be applied as a spray in a minimum of 1 gal of water per acre for aerial applications and a minimum of 5 gals of water per acre for ground applications. **Bermudagrass and Forage Grasses (other than Range and Pasture Grasses)**—Thrips, Whiteflies, Grasshoppers Apply 1 pt per acre. Dairy and beef cattle and sheep (1) may be grazed immediately following application. (2) may be fed green forage immediately following cutting and (3) forage grasses may be cut for hay 7 days following treatment. Do not

EPA Reg. No. 655-776
 Federal Insecticide, Fungicide, and Rodenticide Act
 registered under EPA Reg. No. 655-776

pre DIAZIN INSE

FOR THE CONTROL OF C VEGETABLES, FIELD CROPS
ACTIVE INGREDIENT:
 Diazinon (0,0-diethyl 4-pyridinyl) phosphorothioate
Xylene Range Aroma
INERT INGREDIENTS:

DIAZINON AG500 contains 4 lb PRENTOX®—Registered Trade Name
 Co., Inc.

PRECAUTION HAZARDS TO HUMANS

May be fatal if swallowed. May spray mist. Do not get in eyes or on hands. Avoid contamination. Do not permit children or pets to go down. Keep out of reach of domestic animals. Do not use on household pets or livestock.

ENVIRONMENTAL

This product is toxic to fish and aquatic life. Keep out of lakes, streams, ponds, rivers, and other bodies of water. Do not use in areas where impact on the water table may occur. Fish and Game Agencies should be notified of use. Do not use in areas where impact on the water table may occur. Fish and Game Agencies should be notified of use. Do not use in areas where impact on the water table may occur. Fish and Game Agencies should be notified of use.

PHYSICAL

Do not use, pour, spill or store

DIRECTIONS

For use on 100 Federal Insecticide Act. Do not apply to plants or animals. Do not use in areas where impact on the water table may occur. Fish and Game Agencies should be notified of use.

GENERAL INFORMATION

Diazinon is a phosphorothioate insecticide. It is effective against a wide range of insects. It is registered for use on a wide range of crops. It is a contact and stomach poison. It is effective against a wide range of insects. It is registered for use on a wide range of crops. It is a contact and stomach poison.

EPA EST NO 655-GA-1

NET CONTENTS

LOT NO.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
WARNING**

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get on eyes, on skin or on clothing. Wash thoroughly after handling. Avoid contamination of food. Do not use on humans or pets. Do not permit children or pets to go on treated areas until spray has completely dried. Keep out of reach of domestic animals. Do not contaminate feed or pasture for ruminant part of livestock.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, marshes, and estuaries except for the use of bait for treated water in fish ponds. Do not apply to water that will be used for irrigation purposes and human and livestock consumption. Shrimp and crab may be killed at application rates not recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are in contact or exposure. This product must not be used in areas where important and/or protected endangered species is likely to contact your spray, fish and birds. Apply a future application of this product. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by spraying. Frequent use or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or to drifting spray during flowering time. Do not apply when bees are actively visiting the crop or crop area. Avoid spraying during the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the arrival of bees to the area.

PHYSICAL & CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

This is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to cause by or through drift exposure workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

~~Apply this product in accordance with the directions on the label for various crops. Do not use this product on citrus, cotton, alfalfa, and other crops unless specifically labeled for such use.~~

GENERAL INFORMATION

PRENTISS is an emulsifiable solution which when diluted with water according to the directions herein is intended for the control of insect pests of agricultural crops, fruit trees, ornamentals, lawns, and lawns. Do not use on livestock. In the directions for use given below, best control is obtained when application is made at first sign of insects. Best control should be repeated, as necessary, to maintain control. **NOTE:** This product is highly toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, marshes, and estuaries except for the use of bait for treated water in fish ponds. Do not apply to water that will be used for irrigation purposes and human and livestock consumption. Shrimp and crab may be killed at application rates not recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are in contact or exposure. This product must not be used in areas where important and/or protected endangered species is likely to contact your spray, fish and birds. Apply a future application of this product. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by spraying. Frequent use or disposal of wastes. Apply this product only as specified on this label.

~~Apply this product in accordance with the directions on the label for various crops. Do not use this product on citrus, cotton, alfalfa, and other crops unless specifically labeled for such use.~~

EPA EST NO 655-GA-1

EPA REG. NO. 655-~~655~~

NET CONTENTS

LOT NO.

Manufactured by



Plant: Keolin Road Sandersville, GA. 31082
Office: C.B. 2000 Floral Park, N.Y. 11001

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged. Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material such as clay diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Do not store use pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

DIRECTIONS FOR USE (CONT'D)

~~**FRUIT AND NUT CROPS - NOTE:** Unless otherwise stated, the rates listed below should be applied as adult full cover spray. **Almonds** - Mites, Twig Borers. Apply 1 1/2 pts per 100 gals of water. **Apricots** - Mites, Scale. San Jose Scale. Mix 1 1/2 pts Diazinon plus 2-3 gals dormant oil per 100 gals of water and apply as a dormant spray. Almond hulls may be used for feed for livestock. **Apples** - Aphids, Brown Mites (Clover Mites), Two Spotted Spider Mite, Olive Scale Crawlers, San Jose Scale Crawler. Mix 1 pint AG500 with 10 gallons of light medium horticultural oil per 100 gallons of water. Apply when scale crawlers are present. **Apricot Mealybugs** - Mix 1 pt per 100 gals of water. Apply as a cover spray from petal fall to June. Do not apply within 10 days of harvest. **Blueberries** - Cranberry Fruitworms, Cherry Fruitworm, Blueberry Maggots, Aphids, Thrips, Two Spotted Mites. Mix 1 pt per 100 gals of water. Apply up to 200 gal of spray per acre. Do not apply within 7 days of harvest. **Caneberries** - Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries - Leafhoppers, Thrips, Aphids, Two Spotted Mites, Raspberry Sawflies. Apply 2 pts per acre in a minimum of 100 gals of water. Raspberry fruitworms. Apply at the above rates when blossoms buds separate and repeat application immediately before blossoms open. **Dryberry Mites** - Apply 1-2 pts per acre in a minimum of 100 gals of water. Apply as a dormant spray in early spring when canes are still on the ground and buds are closed. Make a second application when the buds are well developed but still closed. Do not apply within 7 days of harvest. **Raspberry Crown Borers (Raspberry Root Borers)** - Apply 2 qts per acre in a minimum of 100 gals of water as a drench to the crown area and to the lower cane. Apply either in the spring before berries begin to form or in the fall after harvest. **Cherries** - Eye Spotted Bud Moths, Fruit Tree Leaf Rollers. Mix 1 pt per 100 gals of water. Apply in cover sprays as necessary beginning with petal fall. **Leafhoppers** - Apply 1 pt per 100 gals of water in 3-5 applications beginning at petal fall and continuing through the growing season or apply as infestations develop. **Cherry Fruit Flies** - Mix 1/2 pt per 100 gals of water. Make 3 or 4 applications beginning when adult cherry fruit flies begin to emerge and repeat at 10-day intervals. **Black Cherry Aphids** - Mix 1 pt per 100 gals of water. Apply as a cover spray when aphids appear and repeat after 10 days. **San Jose Scale Crawlers** - Mix 1 pt per 100 gals of water. Apply during the pre-bark stage or when crawlers first appear. Do not apply within 10 days of harvest. **Cherry Rust Mites** - Mix 1/2 pt per 100 gals of water. Apply as a supplemental spray after harvest. **Citrus Fruits** - Grapefruit, Lemons, Limes, Oranges, Tangelos, Tangerines, Citrus, Citron, Kumquats, and hybrids of these - Citrus Aphids, Fruit Tree Leaf Rollers, Cottony Cushion Scale, Crawlers, and Soft Scale Crawlers. Apply 1/2 pt per 100 gals of water (maximum 1000 gals per acre) as a thorough cover spray when aphids and crawler stages of scale insects first appear. **Fruit Tree Leaf Rollers** - Use 1/2 pt per 100 gals of spray (maximum 1000 gals per acre) when an average of 50 percent of eggs in egg masses have hatched in the spring. **Citrus Thrips** - Apply 1 pt in 100 gals of water (maximum 1000 gals per acre) when thrips first appear. **Citrus Snow Scales** - Apply 1-2 pts per 100 gals of water (maximum 1000 gals per acre) as a post bloom spray and again as a summer spray. For heavy snow scale infestations use the higher dosage rate. Thorough coverage of tree trunks, limbs, and twigs is essential. Do not apply during bloom. Do not apply to citrus within 21 days of harvest. Shrimp and crab may be killed at application rates recommended on the label. Do not apply where there are important~~

gals of water and apply only). Apply 1 pt per 100 fall and continuing the develop. White Peach S 1 1/2 pts per 100 gals immature scale. Active applications may be ne to neclatines within 10 Case bearers Walnut C in a minimum of 100 **Pineapples** - Scales of medium horticultural on the plants following acre in a minimum of 1 Pineapple forage and application. **Plums and Red Mites** - Apply 1 pt Plum Aphids. Thistle apply within 10 days. **Spider Mites** - Apply 1 Apply 1 qt per 100 gal Plants should be suffi forage. Make additi Crickets. Broadcast 1 days before transpla immediately. **Strawbe Apply 200 gals of spr within 5 days of harve forage spray in a mi Crawlers. Walnut Case minimum of 100 gals. **VEGETABLE FOLIAR** Unless otherwise sta always be applied to water per acre for acre for ground app Beans - Black Bean 1 1/2 qts per acre. Cucu acre. Do not apply w may be fed to dairy treatment of forage is Leafhoppers, Mites harvest. Bean forage sheep. Wait 4 days for Cabbage, and Caulif Cabbageworms. Appl will usually be requi days or to cabbage a Diamondback Moth. Generally applicati Brussels sprouts with Carrots - Aphids, A harvest. Celery - Aph acre. Do not apply w trimmed (tops remove used for feed of food Aphids, Imported C Cabbage Bug, Corn collards, kale or turn harvest. Cucumbers - per acre. Do not app Aphids, Diverous Le; 10 days of harvest. **Cranberries, Honeyde of these, Watermelon Aphids, Leafhoppers, Miners, Thrips. Apply squash within 3 day Mushroom Moths - in 50 gals of water (inside walls and venti been pasteurized by spoaming. To treat w pasteurizing mix 1/2 pt compost. Onions - C 10 days of harvest. Pe may be picked immo and beef, cattle and tr out for hay. Peppers - Do not apply within****

emergence
resistant
1-1/2 qt
1-2 qts
state dosage
Work
prior to
1-2 inches
Cutworms
Immediately
the soil 4-
following last
and sheep
2-4 qts per
3 inches for
per acre just
for surface
Cutworms
coverage of
per Two or
necessary. Do
of leaf beetle
and (3) may
Cutworms
prior to
Cutworms
one (Surface
Work
6 inches for
2-4 qts per
immediately
Apply
application
Tea tree oil
work into
Apply as a
of buds with
effects are
of seed
Seed-Corn
Work
per acre 3-4
Apply
Apply as a
seasons and a
applications
and pasture
per acre Dairy
following
planting and
work. Do not
back. Banta,
Stamps -
right of fol
age of cattle
and may be fed
for hay
solution or
directly with
water.

**DIAZINON TM
INSECTICIDE**