

655-794

01/24/2003

118

Prentox Malathion 57% E.C.
EPA File Symbol 655-794

ACCEPTED
JAN 24 2003
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 655-794

Front Panel



MALATHION 57% E.C.

Not for use on turf being grown for sale or other commercial use as sod, or on plants being grown for sale, or on trees being grown for sale or other commercial use, or for the production of timber or wood products, or on crops, timber, or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds. For use on plants in non-crop and non-timber areas only.

ACTIVE INGREDIENTS:

Malathion: O,O-dimethyl phosphorothioate of diethyl mercaptosuccinate 57%

INERT OTHER INGREDIENTS*: 43%

TOTAL: 100%

*Contains Xylene Range Aromatic Solvent

PRENTOX® - Registered Trademark of Prentiss Incorporated

KEEP OUT OF REACH OF CHILDREN WARNING	
FIRST AID	
Have product container or label with you when calling a poison control center or physician, or going for treatment.	
If swallowed	<ul style="list-style-type: none"> • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 immediately for treatment advice. • Do not give person any liquid. • Do not induce vomiting unless told to do so by the Poison Control Center or physician. • Do not give anything by mouth to an unconscious or convulsing person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 for treatment advice.
NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, but should be administered only in conjunction with atropine sulfate. Contains xylene range aromatic solvent - vomiting may cause aspiration pneumonia.	
For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.	
See inside of booklet for additional precautionary statements and directions for use.	

EPA Reg. No. 655-794

598

EPA Est. No. 655-GA-1

Manufactured by:

PRENTISS INCORPORATED

Plant: Kaolin Road, Sandersville, GA 31082

Office: C.B. 2000, Floral Park, NY 11002-2000

PRENTOX[®] MALATHION 57% E.C.

Not for use on turf being grown for sale or other commercial use as sod, or on plants being grown for sale, or on trees being grown for sale or other commercial use, or for the production of timber or wood products, or on crops, timber, or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds. For use on plants in non-crop and non-timber areas only.

ACTIVE INGREDIENTS:

Malathion: O,O-dimethyl phosphorothioate of diethyl mercaptosuccinate 57%

INERT OTHER INGREDIENTS*: 43%

TOTAL: 100%

*Contains Xylene Range Aromatic Solvent

PRENTOX[®] - Registered Trademark of Prentiss Incorporated

KEEP OUT OF REACH OF CHILDREN WARNING	
FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 immediately for treatment advice. • Do not give person any liquid. • Do not induce vomiting unless told to do so by the Poison Control Center or physician. • Do not give anything by mouth to an unconscious or convulsing person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 for treatment advice.
<p>NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, but should be administered only in conjunction with atropine sulfate. Contains xylene range aromatic solvent – vomiting may cause aspiration pneumonia.</p>	
<p>For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.</p>	
<p>See inside of booklet for additional precautionary statements and directions for use.</p>	

EPA Reg. No. 655-794

598

EPA Est. No. 655-GA-1

Manufactured by:

PRENTISS INCORPORATED

Plant: Kaolin Road, Sandersville, GA 31082
Office: C.B. 2000, Floral Park, NY 11002-2000

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not apply this product through any type of irrigation system.

For use on vegetables: 10 gallons of water will cover approximately 400 square feet (an area 10' X 40').
For use in home vegetable gardens only.

Vegetable	Pests Controlled	Amount to Add to 10 Gal. Water	Interval (Days) Between Last Application and Harvest
Beans (Green beans, Lima beans, Navy beans, Red Kidney beans, Snap beans, Wax beans, Cowpeas and Blackeyed Peas)	Aphids, Bean Leaf Beetle, Blister Beetle, Cucumber Beetle, Potato Leafhoppers, Lygus Bugs	3 fl. oz. (6 Tbsp.)	Do not apply within 1 day of harvest.
	Japanese Beetle, Mexican Bean Beetle	2 ½ - 3 fl. oz. (5 - 6 Tbsp.)	
	Red Spider Mites	2 ½ fl. oz. (5 Tbsp.)	
Beets (Garden & Table)	Aphids, Leafhoppers	2 ½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply within 7 days of harvest if tops are to be used as food or feed.
Broccoli	Aphids, Cabbage Looper, Imported Cabbageworm	1 ½ - 3 fl. oz. (3 - 6 Tbsp.)	Do not apply within 3 days of harvest.
Brussels Sprouts	Aphids	1 ½ - 3 fl. oz. (3 - 6 Tbsp.)	Do not apply within 7 days of harvest.
Cabbage	Aphids, Cabbage Looper, Imported Cabbageworm	1 ½ - 3 fl. oz. (3 - 6 Tbsp.)	Do not apply within 7 days of harvest. For control of caterpillars on summer and fall plantings in the South, begin when true leaves appear. On the other plantings, and for control of other insects, begin when insects appear.
	Diamondback Moth, Webworm	3 fl. oz. (6 Tbsp.)	
Cauliflower	Diamondback Moth	3 - 6 fl. oz. (6 - 12 Tbsp.)	Do not apply within 7 days of harvest.
Celery, Anise	Aphids, Spider Mites	2 ½ fl. oz. (5 Tbsp.)	To be applied to fresh leaves and stalks only. Do not apply within 7 days of harvest.
Collards	Aphids	2 ½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply within 7 days of harvest. For control of caterpillars, on summer and fall plantings in the South, begin when true leaves appear. On other plantings, and for control of other insects, begin when insects appear.
	Harlequin Cabbage Bug	1 ½ fl. oz. (3 Tbsp.)	
Corn (Sweet)	Japanese Beetle	3 fl. oz. (6 Tbsp.)	Do not apply within 5 days of harvest. Injury may occur in the whorl or to the silks.
Cucumbers	Aphids, Pickleworms, Spider Mites	2 ½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply to cucumbers unless plants are dry. Do not apply within 1 day of harvest.
	Squash Vine Borer	5 fl. oz. (10 Tbsp.)	
	Cucumber Beetle, Leaf Miner	3 fl. oz. (6 Tbsp.)	
Dandelion, Parsley, Parsnip, Swiss Chard, Watercress	Aphids	1 ½ - 3 fl. oz. (3 - 6 Tbsp.)	Do not apply within 21 days of harvesting parsley or within 7 days of harvesting dandelion, parsnip, Swiss chard or watercress.
Eggplant	Aphids, Spider Mites	1 ½ fl. oz. (3 Tbsp.)	Do not apply within 3 days of harvest.
	Lacebugs	5 fl. oz. (10 Tbsp.)	
Endive	Aphids, Spider Mites	2 ½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply within 7 days of harvest.
Garlic, Shallots	Aphids, Thrips	2 ½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply within 3 days of harvest.

Vegetable	Pests Controlled	Amount to Add to 10 Gal. Water	Interval (Days) Between Last Application and Harvest
Kale	Aphids, Cabbage Looper, Imported Cabbageworm	1 ½ - 3 fl. oz. (3 - 6 Tbsp.)	Do not apply within 7 days of harvest. For control of caterpillars on summer plantings in the South, begin when true leaves appear. On other plantings, and for control of other insects, begin when insects first appear.
	Webworm, Diamondback Moth	3 fl. oz. (6 Tbsp.)	
Kohlrabi	Aphids	1 ½ - 3 fl. oz. (3 - 6 Tbsp.)	Do not apply within 7 days of harvest.
	Cabbage Looper, Diamondback Moth, Imported Cabbageworm, Webworm	3 fl. oz. (6 Tbsp.)	
Leek	Aphids	2 ½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply within 3 days of harvest.
Lentils	Cowpea Aphid, Pea Aphid	2 ½ fl. oz. (5 Tbsp.)	Do not apply within 3 days of harvest.
Lettuce	Aphids, Leafhoppers, Spider Mites	3 fl. oz. (6 Tbsp.)	Do not apply within 7 days of harvesting head lettuce or within 14 days of harvesting leaf lettuce.
Melons (Cantaloupe, Casaba, Crenshaw, Honeydew Melons, Honey Balls, Muskmelons, Persian Melons and hybrids of these, Watermelons and their hybrids)	Aphids, Spider Mites	2 ½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply to melons unless plants are dry. Do not apply within 1 day of harvest.
	Cucumber Beetles	3 - 5 fl. oz. (6 - 10 Tbsp.)	
	Leaf Miners	3 fl. oz. (6 Tbsp.)	
	Leafhoppers	2 ½ - 3 fl. oz. (5 - 6 Tbsp.)	
	Pickleworms	3 fl. oz. (6 Tbsp.)	
	Squash Vine Borer	5 fl. oz. (10 Tbsp.)	
Mustard Greens	Aphids, Cabbage Looper, Imported Cabbageworm	3 fl. oz. (6 Tbsp.)	For control of caterpillars on summer and fall plantings in the South, begin when true leaves appear. On other plantings, and for control of other insects, begin when insects appear. Do not apply within 7 days of harvest.
Okra	Aphids	2 ½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply within 1 day of harvest. Make no application after pods start to form.
	Japanese Beetle	3 fl. oz. (6 Tbsp.)	
Onions (Bulb & Green)	Onion Maggots	2½ - 4 fl. oz. (5 - 8 Tbsp.)	Do not apply within 3 days of harvest.
	Onion Thrips	2½ - 3 fl. oz. (5 - 6 Tbsp.)	
Peas	Aphids	2½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply within 7 days of harvest if vines are to be fed, or within 3 days of harvest if vines are not to be fed.
	Grasshoppers	3 fl. oz. (6 Tbsp.)	
	Leafhoppers	2 - 6 fl. oz. (4 - 12 Tbsp.)	
Peppermint, Spearmint	Aphids, Flea Beetles, Leafhoppers, Spider Mites	2½ fl. oz. (5 Tbsp.)	Do not apply within 7 days of harvest.
Peppers (Bell)	Aphids	1½ - 4 fl. oz. (3 - 8 Tbsp.)	Do not apply within 3 days of harvest.
	Pepper Maggots	4 fl. oz. (8 Tbsp.)	
Potatoes (White)	Aphids, Grasshoppers, Leafhoppers	3 fl. oz. (6 Tbsp.)	May be applied on the day of harvest.
	False Chinch Bugs	2½ fl. oz. (5 Tbsp.)	
	Mealybugs	3 - 4 fl. oz. (6 - 8 Tbsp.)	
Pumpkins	Aphids, Pickleworms, Spider Mites	3 fl. oz. (6 Tbsp.)	Do not apply to pumpkins unless plants are dry. Do not apply within 3 days of harvest.
	Leafhoppers	2½ - 3 fl. oz. (5 - 6 Tbsp.)	
	Cucumber Beetles, Squash Vine Borer	5 fl. oz. (10 Tbsp.)	

Vegetable	Pests Controlled	Amount to Add to 10 Gal. Water	Interval (Days) Between Last Application and Harvest
Radish, Horseradish	Aphids	2½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply within 7 days of harvest.
Rutabaga	Aphids	2½ fl. oz. (5 Tbsp.)	Do not apply within 7 days of harvest.
Spinach	Aphids, Leafhoppers	3 fl. oz. (6 Tbsp.)	Do not apply within 7 days of harvest.
Squash	Aphids, Leaf Miners, Spider Mites, Pickleworm	3 fl. oz. (6 Tbsp.)	Do not apply to squash unless plants are dry. Do not apply within 1 day of harvest.
	Cucumber Beetles, Squash Vine Borer	5 fl. oz. (10 Tbsp.)	
Strawberries	Aphids, Spider Mites, Field Crickets, Lygus Bugs, Spittlebugs, Thrips	2½ fl. oz. (5 Tbsp.)	Do not apply within 3 days of harvest.
	Potato Leafhopper, Strawberry Leafroller, Strawberry Root Weevil, Whiteflies	2½ - 4 fl. oz. (5 - 8 Tbsp.)	
Sweet Potatoes	Leafhoppers	2½ - 3 fl. oz. (5 - 6 Tbsp.)	Do not apply within 3 days of harvest.
	Morninglory leaf miner	4 - 5 fl. oz. (8 - 10 Tbsp.)	
Tomatoes	Aphids	2½ fl. oz. (5 Tbsp.)	Do not apply within 1 day of harvest.
	Drosophila	4 fl. oz. (8 Tbsp.)	
	Spider Mites	2½ fl. oz. (5 Tbsp.)	
	Armyworms, Fruitworms (California only)	8¼ fl. oz. (17½ Tbsp.)	Do not apply within 3 days of harvest.
Turnip	Aphids, Cabbage Looper, Imported Cabbageworm	1½ - 3 fl. oz. (3 - 6 Tbsp.)	For control of caterpillars on summer and fall plantings in the South, begin when true leaves appear. Do not apply within 7 days of harvest if tops are to be used for food or feed.

VEGETABLES GROWN IN GREENHOUSES: Apply to give complete coverage.
For use on vegetables grown in home greenhouses only.

Vegetable	Pests Controlled	Amount to Add to 10 Gal. of Water	Interval (Days) Between Last Application and Harvest
Cucumbers	Aphids, Armyworms, Cabbage Loopers, Earwigs, Garden Fleahoppers, Mealybugs, Spider Mites, Thrips, Whiteflies	2½ - 3 fl. oz. (5 - 6 Tbsp.)	Spray plants thoroughly. Do not apply to cucumber unless plants are dry. Do not apply within 1 day of harvest.
Endive	Aphids, Armyworms, Cabbage Loopers, Serpentine Leaf Miners, Thrips, Whiteflies	2½ - 3 fl. oz. (5 - 6 Tbsp.)	Spray plants thoroughly. Do not apply within 7 days of harvest.
Lettuce	Aphids, Armyworms, Cabbage Loopers, Serpentine Leaf Miners, Thrips, Whiteflies	2½ - 3 fl. oz. (5 - 6 Tbsp.)	Spray plants thoroughly. Do not apply within 14 days of harvest.
Tomatoes	Aphids, Armyworms, Cabbage Loopers, Cutworms, Drosophila, Garden Fleahoppers, Mealybugs, Serpentine Leaf Miners, Spider Mites, Thrips, Tomato Russet Mites, Whiteflies	2½ - 3 fl. oz. (5 - 6 Tbsp.)	Spray plants thoroughly. Do not apply within 1 day of harvest.

FRUITS: For use on fruits in home gardens only.

Fruit	Pests Controlled	Amount to Add to 10 Gal. of Water	Interval (Days) Between Last Application and Harvest
Apples	Budmoth, Forbes Scale, Woolly Apple Aphid	1½ fl. oz. (3 Tbsp.)	Do not apply within 3 days of harvest.
	Eastern Tent Caterpillar	1½ - 2½ fl. oz. (3 - 5 Tbsp.)	
	Green Apple Aphid, Rosy Apple Aphid	2½ fl. oz. (5 Tbsp.)	
	Codling Moth, Plum Curculio, Red-banded Leafroller	3 fl. oz. (6 Tbsp.)	
	Mealybug, Mites (such as Clover Mite, European Red Mite, Twospotted Spider, Mite, Williamette, Mite, Red Apple Aphid)	1½ - 3 fl. oz. (3 - 6 Tbsp.)	
Prentox Malathion 57% E.C. may cause fruit injury to McIntosh and Cortland varieties if spray is applied within 4 weeks of harvest.			
Dormant or Delayed Dormant Spray applications	Aphids, Mites, Redbanded Leafrollers	1.5 fl. oz. + 13 oz superior oil	Make full coverage dormant or delayed dormant only.
Apricots	Aphids, Codling Moths, European Fruit Lecanium, Orange Tortrix, Soft Brown Scale, Terrapin Scale	1½ - 2½ fl. oz. (3 - 5 Tbsp.)	Do not apply within 7 days of harvest.
Avocado	Greenhouse Thrips, Latania Scale, Omnivorous Looper, Orange Tortrix, Soft Brown Scale	2½ fl. oz. (5 Tbsp.)	Do not apply within 7 days of harvest.
Blackberry, Boysenberry, Dew berry, Loganberry, Raspberry	Aphids, Rose Scale Chafers, Japanese Beetle, Leafhoppers, Mites, Thrips	2½ fl. oz. (5 Tbsp.)	Do not apply within 1 day of harvest.
Cherry (Sweet & Tart)	Black Cherry Aphid, Fruit-Tree Leafroller	2½ fl. oz. (5 Tbsp.)	Do not apply within 3 days of harvest.
	Cherry Fruit Fly, Japanese Beetle	1½ fl. oz. (3 Tbsp.)	
Injury may occur on certain varieties of sweet cherries, particularly in the Northwest.			
Citrus (Grapefruit, Kumquat, Lemon, Lime, Orange, Tangerine, Tangelo)	Black Scale (single and off-brooded), California Red Scale, Citricola Scale, Purple Scale, Soft Brown Scale, Tangelo Yellow Scale	1½ - 2½ fl. oz. (3 - 5 Tbsp.)	Do not apply when trees are in bloom. Do not apply within 7 days of harvest.
	Florida Purple Scale, Florida Red Scale	3 fl. oz. (6 Tbsp.)	
	Citrus Red Mites, Thrips	1¼ fl. oz. (7 Teaspoons)	Do not apply when trees are in bloom. Do not apply within 7 days of harvest.
Currants, Gooseberries	Japanese Beetles	2½ fl. oz. (5 Tbsp.)	Do not apply within 3 days of harvest.
	Rose Chafers	1½ fl. oz. (3 Tbsp.)	
Grapes	Leafhoppers, Spider Mites	2½ fl. oz. (5 Tbsp.)	Do not apply within 3 days of harvest.
	European Fruit Lecanium	2½ fl. oz. (5 Tbsp.)	Make full coverage applications when newly hatched nymphs are migrating over vines, usually shortly after bloom. Do not apply within 3 days of harvest.
	Mealybugs	2½ fl. oz. (5 Tbsp.)	Do not apply within 3 days of harvest.
Injury may occur on grapes of Almeira, Cardinal, Italia and Ribier varieties when sprays containing malathion are applied after clusters appear.			
Nectarines	Spider Mites	1½ - 3 fl. oz. (3 - 6 Tbsp.)	Do not apply within 7 days of harvest.
	Plum Curculio	3 fl. oz. (6 Tbsp.)	
Malathion may cause fruit spotting on nectarines.			

Fruit	Pests Controlled	Amount to Add to 10 Gal. of Water	Interval (Days) Between Last Application and Harvest
Papaya	Aphids	2½ - 3 fl. oz. (5 - 6 Tbsp.)	May be applied on the day of harvest.
Peaches	Black Cherry Aphid, Black Peach Aphid, European Red Mite, Green Peach Aphid, Rusty Plum Aphid, Japanese Beetle, Spider Mites	1½ fl. oz. (3 Tbsp.)	Do not apply within 7 days of harvest.
	Oriental Fruit Moth, Plum Curculio	3 fl. oz. (6 Tbsp.)	Do not apply within 7 days of harvest.
	Cottony Peach Scale, European Fruit Lecanium, Tarrapin Scale	3 fl. oz. (6 Tbsp.)	Make full coverage applications at completion of hatching, about the last of June. Do not apply within 7 days of harvest.

OUTDOOR ORNAMENTALS: Including ornamental flowering plants, ornamental lawns and turf and ornamental woody plants.

Flowers, Shade Trees, Shrubs:

Injury may occur on Ferns, Hickory, Viburnum, Lantana, Crassula and Canareti Juniper following the use of Prentox Malathion 57% E.C. Slight injury has also been reported on Boston, Pteris, and Maidenhair Ferns, Petunias, Small-Leaf Spirea, White Pine and Maples. Under extreme heat, drought and disease conditions, emulsifiable concentrates may cause slight damage to elms.

For use on outdoor ornamentals grown around homes only.

Pest	Rate	Remarks
Aphids, Azalea Scale, Bagworms, Birch Leaf Miner, Black Scale Crawlers, Boxwood Leaf Miner, Euonymus Scale, European Pine Shoot Moth, Four-lined Leaf Bug, Japanese Beetle Adults, Lace Bugs, Magnolia Scale, Mealybugs, Monterey Pine Scale, Pine Leaf Scale, Potato Leafhopper, Rose Leafhopper, Scurfy Scale, Soft Brown Scale, Spider Mites, Tarnished Plant Bug, Tent Caterpillar, Thrips, Whiteflies	2 Teaspoons/Gallon	—
Fletcher Scale, Florida Red Scale, Juniper Scale, Oak Kermes, Oyster Shell Scale	2 Teaspoons/Gallon	Apply when scale crawlers have settled on foliage.
Pine Needle Scale	4 Teaspoons/Gallon	—
Wax Scale	4 Teaspoons/Gallon	Apply in Spring when crawlers are active. Repeat 1 or 2 full coverage applications at ten day intervals.

OUTDOORS, AROUND THE HOME AND OTHER STRUCTURES AND INSIDE HOME GREENHOUSES:

Site	Pest	Rate	Remarks
In and around home greenhouses and home gardens and outside homes and other structures.	Millipedes, Sawbugs Springtails	1 tbsp./gal.	Apply to 150 square feet of soil or where insects congregate. Repeat at 7 to 10 day intervals as needed.
Lawns, Turf and Golf Courses	Ant Mounds	2½ oz./gal.	Spray ant hills thoroughly so that they are well soaked. For other small ants in flower beds, lawns, around trees, spray lightly in the infested areas. Repeat in 10 to 15 days if ants return.
	Ground pearls	10-13 oz./gal.	Make full coverage to soil surface when ground pearl nymphs are in the pink, "crawler" or active stage and immediately wash into soil with additional water.

DO NOT ALLOW CHILDREN OR PETS IN TREATED AREAS UNTIL SPRAYS HAVE DRIED.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: WARNING. Causes substantial but temporary eye injury. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing vapors or spray mist. Do not get in eyes or on clothing. Wear goggles, face shield or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before re-use.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, aquatic invertebrates and aquatic life stages of amphibians. For terrestrial uses, do not apply directly to water, or to where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment wash waters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Use only with adequate ventilation when applying in greenhouses. Do not allow contact with treated surfaces until sprays have dried.

PHYSICAL OR CHEMICAL HAZARDS: Do not use, pour or spill near heat or open flame.

STORAGE AND DISPOSAL

STORAGE: Store in cool, dry area. If container is damaged, stop any leaks by repositioning container, or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent onto a shovel and put the sweepings into a salvage drum. Apply liquid household bleach to contaminated area. Scrub thoroughly using a long handled brush. Let stand for 15 minutes. Wash area thoroughly with water. Place leaking container into a similar drum or glass container. Mark the container with the name of the product, ingredient statement, precautionary statements, and signal word. Contact us for replacement label. Do not store, use pour or spill near heat or open flame. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** ~~Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse container. Rinse thoroughly before discarding in trash.~~ **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. **Do not reuse this container. Never place unused product down any indoor or outdoor drain.**

Adhesive Panel



MALATHION 57% E.C.

Not for use on turf being grown for sale or other commercial use as sod, or on plants being grown for sale, or on trees being grown for sale or other commercial use, or for the production of timber or wood products, or on crops, timber, or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds. For use on plants in non-crop and non-timber areas only.

ACTIVE INGREDIENTS:

Malathion: O,O-dimethyl phosphorothioate of diethyl mercaptosuccinate 57%

INERT OTHER INGREDIENTS*: 43%

TOTAL: 100%

*Contains Xylene Range Aromatic Solvent

PRENTOX® - Registered Trademark of Prentiss Incorporated

**KEEP OUT OF REACH OF CHILDREN
WARNING**

FIRST AID

Have product container or label with you when calling a poison control center or physician, or going for treatment.

If swallowed	<ul style="list-style-type: none"> • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 immediately for treatment advice. • Do not give person any liquid. • Do not induce vomiting unless told to do so by the Poison Control Center or physician. • Do not give anything by mouth to an unconscious or convulsing person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a Poison Control Center, physician, or the National Pesticide Information Center at 1-800-858-7378 for treatment advice.

NOTE TO PHYSICIAN: Malathion may cause cholinesterase inhibition. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, but should be administered only in conjunction with atropine sulfate. Contains xylene range aromatic solvent - vomiting may cause aspiration pneumonia.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

See inside of booklet for additional precautionary statements and directions for use.

EPA Reg. No. 655-794

598

EPA Est. No. 655-A-1

Manufactured by:

PRENTISS INCORPORATED

Plant: Kaolin Road, Sandersville, GA 31082
Office: C.B. 2000, Floral Park, NY 11002-2000