

Prentox Prentish Grass Carp Management Bait EPA Registration No. 655-795

RESTRICTED USE PESTICIDE DUE TO AQUATIC TOXICITY

For retail sale to, and use only by, Certified Applicators or Persons under their direct supervision and only for those uses covered by the Certified Applicators' certification.

PRENTOX

# PRENFISH GRASS CARP MANAGEMENT BAIT

ACCEPTED

ACTIVE INGREDIENTS:

Rotenone
Related Compounds
Piperonyl Butoxide Technical\*
INERT INGREDIENTS:

JUN 18 2001

Under the Federal Lecticide,
Fangistie: a posticide
rog: TOTAL

EPA R PANY 655 - 795

2.64% w/w 3.36% 0.500% 93.50% 100.00%

\*Equivalent to 0.4% (Butylcarbityl)(6-propypiperon ) \*Fether at PRENTOX\* - Registered Trademark of Prentiss Incorporated

PRENFISHTM - Trademark of Prentiss Incorporated

EPA Registration No. 655-795

EPA Est. No. 655-GA-1

U.S. Patent Reg. No. 5,674,518, SNO8/426,898 Filed 10/7/97

	KEEP OUT OF REACH OF CHILDREN
	CAUTION
	FIRST AID  Have product container or label with you when obtaining treatment advice.
If swallowed	<ul> <li>Call a physician, Poison Control Center, or the National Pesticide Telecommunications Network at 1-800- 858-7378 immediately for treatment advice.</li> </ul>
	Do not induce vomiting thless directed by a physician or Poison Control Center.     Do not give anything by mouth to an unconscious or convulsing person.
If on skin	<ul> <li>Wash with plenty of soap and water for 15-20 minutes.</li> <li>Call a physician, Poison Control Center, or the National Pesticide Telecommunications Network at 1-800-858-7378 immediately for treatment advice.</li> </ul>
If in eyes	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center, doctor, or the National Pesticide Telecommunications Network at 1-800-858-7378 immediately for treatment advice.</li> </ul>

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

# ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Fish kills are expected at recommended rates. Consult your State Fish and Game Agency before applying this product to public waters to determine if a permit is needed for such an application. Do not contaminate untreated water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

National Pesticide Telecommunications Network at 1-800-858-7378.

### USE RESTRICTIONS

This product may be used to reduce Prentish Grass Carp Management Bait is to be used to remove undesirable grass carp populations that have been trained to eat fish food pellets and that reside in freshwater lakes, ponds and reservoirs. Use This product is to be used only at locations, rates, and times authorized and approved by the appropriate State and Federal fish and wildlife agencies. This product is to be used only after the of water body to be treated has been probabiled prepared and grass carp within it have been trained to come to designated locations to feed upon floating fish feed. Apply this product only within the range of rates specified in this labeling. Do not apply Prentox Prenfish Grass Carp Management Bait at times when winds and/or waves are high. Collect and properly dispose of all dead fish and unconsumed or leftover bait. Do not use dead fish that were exposed to this product for feed or as food.


GENERAL USE CONSIDERATIONS

Page: 1 of 2

#### ESTABLISHING TEST SITES

Areas designated for baiting should be established away from shore, at or near the edge of littoral zone vegetation and away from areas where boat traffic is likely Where possible, bait stations should be placed so that prevailing winds will blow the GCMB-T away from shore. This will give the carp more time to find and consume the bait. The judgment of experienced Local. State and Federal fisheries personnel should be employed to determine optimal locations for feeding areas in each body of water to be treated. Fish can be trained by application of 1 - 11 lbs. (0.5-5kg) per day of Grass Carp Management Bait-Trainer (GCMB-T) by an automatic fish feeder or by hand. One to three weeks of training should attract any fish in the area. Test site should have a fish feed retention device placed to limit the movement of floating fish feed.

Circumscribe each area designated for baiting by use of a feed retention ring designed to restrain the movement of floating bait Use a feed retention ring to limit the movement of floating bait (see "SAMPLE STATION PLACEMENT" diagram). Each retention ring should encompass surround an area no less than of at least 0.1-9.2 acres (5.350-10,700 sq. ft.). Attach Center the feed retention ring to bait dispensing station on the mechanical feeder, if one is used (as indicated in the diagram). Secure bait station in place either by use of anchors or by mounting baiter on a post sunk into the bottom of the lake, pend, or reservoir. it on a fixed structure in the body of water. If no bait dispenses mechanical feeder is used, restrain the movements of the feed retention ring using with at least two anchors. Allowing enough slack in rope lines to accommodate the range of expected fluctuations in the water level.



#### PREBAITING DIRECTIONS

The goal of a prebaiting period is to have as many grass carp feeding on the trainer bait as possible. Prior to use of this product, train grass carp to eat pelleted feed by applying 1.1-3.3 lbs. (0.5 1.5 kg.) 2-50 lbs. (1-20 kg) of Prentox Fish Prenfish Grass Carp Management Bait-Trainer (GCMB-T). Apply probait trainer bait twice a day, at set times, preferably at dusk and dawn - the times when grass carp are likely to be feeding, most teadily. The amount of trainer bail applied should be increased as the number of fish feeding on the batt increases. If all or most of the trainer bait applied is consumed by grass carp in 20 minutes or less, the next application of bait should be increased by 2 lbs. (1 kg). Applicators should continue to increase the amount of trainer bait applied as long as all or most of the bait is consumed in 20 minutes. One to two three weeks of training trainer bait applications should attract any grass carp in the vicinity to the probated baiting area. The applicator should closely observe feeding by the fish Juring the prebatting period. The floating trainer batt will allow the applicator to determine the number and species of fish feeding on the bait.

#### BAITING DIRECTIONS

After prebait trainer bait is accepted consistently by grass carp, apply Prentox Prenfish Grass Carp Management Bait in the exact same manner and amount as in which the prebait had been was applied. Conditions at the time of application of this product should be the same conditions as existed during the application of the trainer bait. Therefore, care must be taken to prevent or avoid additional personnel, boats, vehicles or activity in the bait station area that may disturb the fish at the time this product is applied. One pellet of this product contains approximately enough rotenone to kill a 2.2 lb. (1 kg.) -fish, grass carp, although some variation can occur. The amount of GCMB applied should be equal to the amount of GCMB-T.

#### EXPECTED RESULTS

It is possible to kill all of the grass carp feeding at the bait station at one time. The number of fish killed is limited to the number of fish feeding on the bait station at one time. fooder and the number of Prentox Prentish Grass Carp Management Bait pellets that are used applied. Fish will take from 1-4 hours to die. The length of time will depend on the amount of bait consumed, temperature of the water and condition of the fish. Once the grass carp have been removed from the reduced in a given area, the bait station should be moved 200-300 meters to attract naive fish.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store only in the original container, in a cool, dry place inaccessible to children and pets. After opening, only store indoors at room temperature. For prolonged storage, refrigerate at less than 50° F (10° C). Do not expose to light for prolonged periods of time. If spilled, sweep up and dispose of as below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by

State and local authorities, by burning. If burned stay out of smoke.			<u>, , , , , , , , , , , , , , , , , , , </u>
NET CONTENTS	LOT NO		1)
Manufactured by:		1 , 3 3	,
PRENTISS INCORPORATED	Plant: Kaolin Ruad Sandersville GA 31082		

Office: C.B. 2000, Floral Park, NY 11002-2000

4/01

Page 2 of 2