

PM-13

Reg # 655-799

503986

1419



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

655-799

APR 22 1996

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product:

Prentox Diazinon Lawn
& Garden Insecticide

Name and Address of Registrant (include ZIP Code):

Prentiss, Inc.
C.B. 2000
Floral Park, NY 11002-2000

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the labeling change listed below before you release the product for shipment:
 - a. Add the phrase "EPA Reg. No. 655-799".
3. Submit three (3) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

Signature of Approving Official:

Date:

APR 22 1996

2919

page 2
EPA Reg. No. 655-799

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosures

Container Label

PRENTOX[®] DIAZINON LAWN & GARDEN INSECTICIDE

Water-Based Concentrate

This product must not be used on golf courses and sod farms. For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

Protects Lawns, Vegetables, Fruits and Nuts, Roses, Flowers, Trees, Shrubs

Controls Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants¹, Flea Beetles, Grubs, Mites, Sod Webworms and many others.

¹ Aids in control

ACTIVE INGREDIENT:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 22.4%

INERT INGREDIENTS: 77.6%

Total: 100.0%

PRENTOX[®] — Registered Trademark of Prentiss Incorporated

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

See full label in booklet under cap for complete directions for mixing and applying this product and a complete list of pests controlled.

EPA REG. NO. 655-*700*
Manufactured for:

EPA EST NO. _____
NET CONTENTS _____

PRENTISS INCORPORATED

Plant: Kaolin Road, Sandersville, GA 31082
Office: C.B. 2000, Floral Park, NY 11002-2000

- Concentrated
- Broad Spectrum Pest Control
- Less Smell
- Low Odor
- Reduced Odor Formulation
- Easy-to-Mix
- Easy-to-Apply
- Handles Easily
- Dilute and Spray
- Excellent Performance
- Controls soil, foliar, and sucking pests
- Convenient water-based formula
- Concentrated = Better Value
- Protects Home Lawns and Home Vegetable Gardens
- Controls pests in the home lawn and home garden
- Controls a wide spectrum of pests
- Soil and Foliar Insecticide with Broad Spectrum Control
- Spray on foliage or soil to control numerous pests
- Reliable control
- Covers up to _____ sq. ft. of lawn
- Fast acting
- Dependable Control
- Time Tested Insecticide
- Economical
- New Water-Based Formula*
- Use on home lawns, ornamentals, vegetables and fruits

ACCEPTED
with COMMENTS
in EPA Letter Dated

APR 22 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
655-799

*The term "new" may be used for no longer than 6 months following product introduction into the marketplace.

4 y 19

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

Prentox Diazinon Lawn & Garden Insecticide is a water-based concentrate product which must be diluted with water before application. To avoid having spray mixture left over, mix only the amount needed. This product kills many outdoor pests which damage lawns, roses, flowers, trees, shrubs, vegetables and fruits, and insects which annoy pets and people.

Notes: (1) Do not use inside the home. (2) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia, or orchids, since plant injury may occur. (3) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (4) Do not apply to food crops grown in greenhouses.

After spraying, wait the following number of days before harvesting fruits, nuts, or vegetables:

| | |
|----------------|---|
| 1 day | Tomatoes |
| 3 days | Melons, Winter Squash |
| 5 days | Peppers, Strawberries |
| 7 days | Beans, Blackberries, Blueberries, Boysenberries, Broccoli, Cauliflower, Cucumbers, Dewberries, Loganberries, Peas, Raspberries, Summer Squash, Sweet Corn |
| 10 days | Chinese Broccoli, Chinese Cabbage, Collards, Kale |
| 14 days | Carrots, Endive (Escarole), Lettuce, Onions (Bulb and Green), Parsnips, Radishes, Rutabagas, Spinach, Turnips |
| 21 days | Apricots, Cabbage, Cherries, Nectarines, Peaches, Plums, Prunes |
| 28 days | Grapes |
| 35 days | Potatoes (Irish) |
| 45 days | Walnuts |
| * | Almonds (one dormant spray only) |

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

*This table is only a partial listing of the pests controlled by this product. See booklet under cap for complete directions for use and pests controlled.

| Problem Area | Pests* |
|---|---|
| Nuisance Pests in Outside Areas | Ants, Brown Dog Ticks, Chiggers, Cockroaches, Crickets, Fleas, Fire Ants, Spiders, and others* |
| Lawns | Chinch Bugs, Ants, Sod Webworms, Brown Dog Ticks, Fleas, White Grubs, Lawn Billbugs, and others* |
| Vegetables: Foliar Application | |
| Beans, Peas (except dried varieties) | Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites |
| Broccoli, Chinese Broccoli, Cabbage, Chinese Cabbage, Cauliflower, Collards, Kale | Aphids, Flea Beetles, Dipterous Leafminers, Diamondback Moths, Imported Cabbageworms |
| Endive (Escarole), Lettuce, Spinach | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites |
| Peppers, Tomatoes | Aphids, Colorado Potato Beetles, Fall Armyworms (Tomatoes), Flea Beetles, Leafhoppers, Southern Armyworms (Tomatoes), Vinegar Flies - <i>Drosophila</i> spp. (Tomatoes) |
| Onions (Bulb and Green) | Onion Thrips |
| Carrots, Parsnips, Potatoes, Radishes, Rutabagas, Turnips | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Colorado Potato Beetles (Potatoes), Southern Armyworms (Potatoes) |
| Sweet Corn | Aphids, Flea Beetles, Grasshoppers, Spider Mites |
| Melons, Squash (Summer, Winter) | Aphids, Thrips, Spider Mites |
| Vegetables: Preplant Soil Application | |
| Beans, Peas (except dried varieties) | Cutworms, Mole Crickets, Wireworms (Snap Beans) |
| Beets, Red (Table) | Cutworms, Mole Crickets |
| Cabbage | Cutworms, Root Maggots, Wireworms |
| Carrots, Sweet Corn | Cutworms, Mole Crickets, Wireworms |
| Lettuce, Tomatoes | Cutworms, Wireworms |
| Radishes, Rutabagas, Turnips | Cutworms |

| | |
|--|---|
| Fruits & Nuts | |
| Almonds | Scale Crawlers |
| Walnuts (CA only - both crops) | Aphids, Scale Crawlers, Mites, and others* |
| Apricots, Peaches, Nectarines | Aphids, Apricot Mealybugs (Apricots), Spider Mites, Scale Crawlers |
| Blueberries | Aphids, Blueberry Maggots, Cherry Fruitworms, Cranberry Fruitworms, Two-Spotted Spider Mites, Thrips |
| Cherries | Black Cherry Aphids, Cherry Fruitflies, Cherry Rust Mites, Scale Crawlers, and others* |
| Plums, Prunes | Aphids, Spider Mites |
| Caneberries, Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries (CA, OR, WA only - all crops) | Aphids, Leafhoppers, Two-Spotted Spider Mites, Thrips, Raspberry Fruitworms |
| Strawberries | Aphids, Two-Spotted Spider Mites, Strawberry Leafrollers |
| Grapes | Aphids, Grape Berry Moths, Grape Leafrollers, Leafhoppers |
| Roses & Flowers | |
| Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses | Aphids, Flea Beetles, Dipterous Leafminers, Leafhoppers, Mealybugs, Two-spotted Spider Mites, Thrips, Whiteflies, and others* |
| Ornamental Trees & Shrubs | |
| Arborvitae, Azaleas, Birch, Boxwood, Camellias, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honeysuckle, Honey Locust, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Scotch Pine, Spirea, Sycamore, Wax Plant (Hoya), Willow, Yellowwood, Yew | Aphids, Mites, Scale Crawlers, Whiteflies, Tent Caterpillars, and others* |

STORAGE and DISPOSAL

Storage

Store in original container in an area inaccessible to children or pets. If product freezes, place container in a warm room (68° F or above) and allow to thaw. Shake well before using.

Disposal

Do not pour the concentrate, the spray mixture, or the rinse water down the drain or sewer. Do not reuse empty container. Triple rinse the empty container, pour the rinse water into the sprayer, the spray area to be treated. Securely wrap empty container in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

6919

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where irrigation (watering) is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least ¼ inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

7 9 19

Front Panel - Booklet Label

PRENTOX® DIAZINON LAWN & GARDEN INSECTICIDE

Water-Based Concentrate

This product must not be used on golf courses and sod farms. For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

Protects Lawns, Vegetables, Fruits and Nuts, Roses, Flowers, Trees, Shrubs

Controls Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants¹, Flea Beetles, Grubs, Mites, Sod Webworms and many others.

¹ Aids in control

ACTIVE INGREDIENT:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 22.4%

INERT INGREDIENTS: 77.6%

Total: 100.0%

PRENTOX® — Registered Trademark of Prentiss Incorporated

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 655-700
Manufactured for:

EPA EST NO.
NET CONTENTS _____

PRENTISS INCORPORATED

Plant: Kaolin Road, Sandersville, GA 31082
Office: C.B. 2000, Floral Park, NY 11002-2000

ACCEPTED
with COMMENTS
in EPA Letter Dated

APR 22 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
655-799

8 9 19

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Prentiss Incorporated or the Seller. All such risks shall be assumed by the Buyer.

Prentiss Incorporated warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. Prentiss Incorporated makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Prentiss Incorporated or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Prentiss Incorporated and the Seller offer this product and, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Prentiss Incorporated.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

Prentox Diazinon Lawn & Garden Insecticide is a water-based concentrate product which must be diluted with water before application. To avoid having spray mixture left over, mix only the amount needed. Refer to the section How to Mix and Apply. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruits and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. Prentox Diazinon Lawn & Garden Insecticide works fast and applications should be repeated only as directed to maintain control.

Notes: (1) Do not use inside the home. (2) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia, or orchids, since injury may occur. (3) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (4) Do not apply to food crops grown in greenhouses.

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

How to Mix and Apply

Step #1: Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

Step #2: Use clean sprayer. Carefully measure and mix the amount of Prentox Diazinon Lawn & Garden Insecticide and water as indicated in the rate chart.

Step #3: Spray as directed in rate chart. Thorough coverage is important. See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

Step #4: Flush sprayer with clean water after each use.

Measurement Conversions

1 tablespoon = 3 teaspoons
1 fl. oz. = 6 teaspoons or 2 tablespoons
8 fl. oz. = $\frac{1}{2}$ cup
 $1\frac{1}{2}$ tablespoons in 3 gals. of water = $1\frac{1}{2}$ teaspoons in 1 gal. of water
2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

Lawn Pest Control

Precautions

1. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.
3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl.
4. Avoid overlapping of sprays.
5. Where irrigation (watering) is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least $\frac{1}{4}$ inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Marion, and St. Augustinegrass.

10919

| Nuisance Pests Outside the Home | |
|--|---|
| Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (<i>Collembola</i>) | <p>Mix 2 fl. oz. in 1 gal. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.</p> <p>To prevent entry into houses, spray a 5 ft. band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 ft. Repeat application as necessary.</p> <p>Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate.</p> |
| Fire Ants | <p>To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.</p> <p>Note: For best results, apply in cool weather 65-80° F, or in early morning or late evening hours. Treat new mounds as they appear.</p> |

Insect Control on Ornamental Trees, Shrubs, and Flowers
Spray when listed pests first appear. Repeat as necessary.

| Trees and Shrubs | Insects | Rate per 1 Gal. of Water | Timing/Special Directions |
|--|--|--------------------------|--|
| Arborvitae, Azaleas, Birch, Boxwood, Camellias, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Scotch Pine, Spirea, Spruce, Sycamore, Wax Plant (Hoya), Willow, Yellowwood, Yew | Aphids, Apple-and-Thorn Skeletonizers, Bagworms, European Pine Sawfly, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak) Hemlock Chermes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Oak Loopers, Obliquebanded Leaf-rollers, Obscure Root Weevils, Omnivorous Leaf-tiers, Pear Slugs, Scale Crawlers (Cottony-cushion, Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft), Tent Caterpillars, Webworms (Cotoneaster, Fall, Juniper, Mimosa), Whiteflies | 2 teaspoons | <p>Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage with spray.</p> <p>Mimosa Webworms: apply when adults first appear. Make 2 additional applications at 4 - 5 week intervals.</p> |

| Roses and Flowers | Insects | Rate per 1 Gal. of Water | Timing/Special Directions |
|--|---|--------------------------|---|
| Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses | Aphids, Dipterous Leafminers, Euonymus Scale Crawlers, Flea Beetles, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies | 2 teaspoons | Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage. |

Fruits and Nuts

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted.

Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

| Crop | PHI* |
|---------------|------|
| Almonds | ** |
| Apricots | 21 |
| Blackberries | 7 |
| Blueberries | 7 |
| Boysenberries | 7 |
| Cherries | 21 |
| Dewberries | 7 |
| Grapes | 28 |
| Loganberries | 7 |
| Nectarines | 21 |
| Peaches | 21 |
| Plums | 21 |
| Prunes | 21 |
| Raspberries | 7 |
| Strawberries | 5 |
| Walnuts | 45 |

*PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

**Dormant spray only.

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

| Crop | Pest | Rate | Timing/Special Directions |
|---|--|--------------------------------|---|
| Almonds (CA only) | San Jose Scale, Parlatoria Scale | 2 teaspoons in 1 gal. of water | Apply only as a dormant spray. Do not apply more than 6 gals. per 500 sq. ft. of home orchard. This product may be used on almonds only in CA. Do not use this product on almonds grown in other states. |
| Apricots | Aphids, Apricot Mealybugs, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose) | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. |
| Blueberries | Aphids, Blueberry Maggots, Cherry Fruitworms, Cranberry Fruitworms, Thrips, Two-spotted Spider Mites | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Allow a minimum of 14 days between applications. Do not apply more than 4 gals. per 500 sq. ft. of blueberry patch. Do not make more than 2 applications per season. Allow a minimum of 7 days between the last application and harvest. |
| Caneberries, (CA, OR, WA only): Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries | Aphids, Leafhoppers, Raspberry Fruitworms, Two-spotted Spider Mites, Thrips | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 5 gals. per 500 sq. ft. of caneberry patch. Do not make more than 5 applications per season a minimum of 14 days apart. Do not apply within 7 days of harvest. This product may be used on caneberries only in CA, OR, and WA. Do not use this product on caneberries grown in other states. |
| Cherries | Black Cherry Aphids, Cherry Fruit-flies, Cherry Fruitworms (Maggots), Fruitree Leafrollers, Mites (Cherry Rust), Scale Crawlers (San Jose) | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest. |

12 9 19

| Crop | Pest | Rate | Timing/Special Directions |
|-------------------|---|---------------------------------|--|
| Grapes | Aphids, Grape Berry Moths, Grape Leafhoppers, Leafhoppers | 1½ teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 3 gals. per 500 sq. ft. of home vineyard. Do not make more than 5 applications per season a minimum of 7 days apart. Do not apply within 28 days of harvest. |
| Nectarines | Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose) | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. |
| Peaches | Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose) | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest. |
| Plums and Prunes | Aphids (Leafcurl Plum, Mealy Plum, Thistle), Mites (Brown, European Red) | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest. |
| Strawberries | Aphids, Two-spotted Spider Mites, Strawberry Leafrollers | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 3 gals. per 500 sq. ft. of strawberry patch. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 5 days of harvest. |
| Walnuts (CA only) | Aphids, Scale Crawlers (Walnut, Italian Pear, Frosted), Spider Mites, Twig Borers | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 6 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest. This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states. |

13 9 19

Vegetables

For foliar or soil applications, Pretox Diazinon Lawn & Garden Insecticide should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil.

Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Intervals (PHI*) for Foliar Applications

| Crop | PHI* (Days) |
|--|-------------|
| Beans ¹ (succulent varieties only: Lima, Pole and Snap) | 7 |
| Broccoli | 7 |
| Cabbage | 21 |
| Cantaloupes | 3 |
| Carrots | 14 |
| Casabas | 3 |
| Cauliflower | 7 |
| Chinese Broccoli, Chinese Cabbage, Collards | 10 |
| Crenshaws | 3 |
| Cucumbers | 7 |
| Endive (Escarole) | 14 |
| Honeydew Melons | 3 |
| Kale | 10 |
| Lettuce | 14 |
| Muskmelons | 3 |
| Onions (Bulb and Green), Parsnips | 14 |
| Peas ¹ | 7 |
| Peppers | 5 |
| Persian Melons | 3 |
| Potatoes (Irish) | 35 |
| Radishes, Rutabagas, Spinach | 14 |
| Squash, Summer | 7 |
| Squash, Winter | 3 |
| Sweet Corn | 7 |
| Tomatoes | 1 |
| Turnips (Roots and Tops) | 14 |
| Watermelons | 3 |

*PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

¹ Do not use this product on dried pea or bean varieties, such as pinto beans, dried limas, split peas, or dried black-eyed peas.

14 8 19

Vegetables

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

| Crop | Pest | Rate | Timing/Special Directions |
|---|--|---|---|
| Beans, Lima, Pole and Snap (succulent varieties only) | Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches. |
| | Wireworms (snap beans only) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |
| | Note: (1) Do not apply this product to dried bean varieties, such as pinto beans or dried limas. (2) To protect bees, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom. | | |
| Beets, Red (Table) | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches. |
| Broccoli | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Cabbage | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 21 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Root Maggots | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |

15919

| Crop | Pest | Rate | Timing/Special Directions |
|-------------------------------------|--|---|---|
| Carrots | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |
| Cauliflower | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Chinese Broccoli | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticide. |
| Chinese Cabbage (Bok Choy and Napa) | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Collards | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Corn (see sweet corn) | | | |
| Cucumbers | Aphids, Thrips, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. |
| Endive (Escarole) | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |

16 9 19

| Crop | Pest | Rate | Timing/Special Directions |
|---|---|---|--|
| Kale | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Lettuce | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |
| Melons (Cantaloupes, Casabas, Crenshaw's, Honeydews, Muskmelons, Persians and Hybrids of these) and Watermelons | Aphids, Thrips, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest. |
| Onions (Bulb and Green) | Onion Thrips | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. |
| Parsnips | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat applications every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Peas (succulent only) | Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |
| | Notes: (1) Do not apply this product to dried pea varieties, such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom. | | |

17919

| Crop | Pest | Rate | Timing/Special Directions |
|----------------|--|---|---|
| Peppers | Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest. |
| Potatoes | Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden. | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Radishes | Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| Rutabagas | Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| Spinach | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Squash, Summer | Aphids, Thrips, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. |
| Squash, Winter | Aphids, Thrips, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest. |

18 9 19

| Crop | Pest | Rate | Timing/Special Directions |
|--|---|---|---|
| Sweet Corn | Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last application. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |
| Note: To protect bees, do not apply to corn foliage during the pollen shed period. | | | |
| Tomatoes | Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies (<i>Drosophila</i> spp.) | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |
| Turnips and Turnip Tops | Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |

STORAGE AND DISPOSAL

Storage: Store in original container in an area inaccessible to children and pets. If product freezes, place container in a warm room (68° F or above) and allow to thaw. Shake well before using.

Disposal: Do not pour the concentrate, the spray mixture, or the rinse water down the drain or sewer. Do not reuse empty container. Triple rinse the empty container, pour the rinse water into the sprayer, then spray the area to be treated. Securely wrap empty container in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

19 9 19
7

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.
If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.
If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
If on skin: Wash with plenty of soap and water. Get medical attention.
Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where irrigation (watering) is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Adhesive Panel

PRENTOX® DIAZINON LAWN & GARDEN INSECTICIDE

Water-Based Concentrate

This product must not be used on golf courses and sod farms. For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

Protects Lawns, Vegetables, Fruits and Nuts, Roses, Flowers, Trees, Shrubs

Controls Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants¹, Flea Beetles, Grubs, Mites, Sod Webworms and many others.

¹ Aids in control

ACTIVE INGREDIENT:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 22.4%

INERT INGREDIENTS:

..... 77.6%

Total: 100.0%

PRENTOX® — Registered Trademark of Prentiss Incorporated

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.
If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.
If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
If on skin: Wash with plenty of soap and water. Get medical attention.
Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.

SEE END OF BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 655-

Manufactured for

PRENTISS INCORPORATED

EPA EST NO.

NET CONTENTS

Plant: Kaolin Road, Sandusville, GA 31082

Office: C.B. 2000, Floral Park, NY 11002-2000

9919

To control pests listed below, apply recommended rates.

| Pests | Rate Diazinon Lawn & Garden Insecticide | Water | Remarks |
|--|---|-------------------------|--|
| Lawn Chinch Bugs | 1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Thoroughly wet down grass a few hours before applying. In the North, if Chinch Bugs are the only problem, a lower rate of application can be used: 3 teaspoons in 3 gals. of water covers 125 sq. ft. or 4 fl. oz. in 24 gals. of water per 1,000 sq. ft. |
| Ants, Armyworms, Bermudagrass Mites, Clover Mites, Springtails (<i>Collembola</i>), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs) | 1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Thoroughly wet down grass a few hours before applying. For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active. Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect. For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry. |
| Brown Dog Ticks, Chiggers, Fleas | 1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Note: Do not apply this product to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house. |
| Hyperodes Weevil | 1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application. |
| Millipedes | 1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Thoroughly wet grass a few hours before applying. |
| Rhodesgrass Scates | 1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Apply when crawlers first emerge. |
| White Grubs of Japanese Beetle, European Chafer, Southern Chafer | 1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Apply any time between late July and early October. Water grass thoroughly after application. |