

PM 03

655-799

2-18-98

10416

537825

FEB 18 1998

Mr. Richard Miller
Prentiss Incorporated
C.B. 2000
Floral Park, NY 11002-2000

Dear Mr. Miller:

Subject: Amendment - Add Apples and Pears,
Update label
Prentox Diazinon Lawn & Garden Insecticide
EPA Registration Number 655-799
Your submission dated August 15, 1996, and
associated authorization dated October 20, 1997

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a) At the top of the front panel of the booklet label, revise the claim "and many others" to "and many others as listed in directions".
 - b) In the second Note under the heading General Information, revise "since injury" to "since plant injury".
 - c) In the directions for Trees and Shrubs, add the missing vertical line separating the Rate column from the Timing/Special Directions column.
 - d) In the directions for Roses and Flowers, add the missing vertical line separating the Rate column from the Timing/Special Directions column.

- e) For **Fruits and Nuts**, in the definition of PHI appearing under the Pre-Harvest Interval table, add "to wait" between "days" and "between".
- f) For **Pears**, in the **Pests** listing add a comma between "San Jose" and "Forbes".
- g) For **Vegetables**, in the definition of PHI appearing under the Pre-Harvest Interval table, add "to wait" between "days" and "between".
- h) For **Lettuce**, add the missing horizontal lines in the **Pests**, **Rate** and **Timing/Special Directions** columns between the three pest entries.
- i) For **Peas (succulent only)**, add the missing horizontal lines in the **Pests**, **Rate** and **Timing/Special Directions** columns between the four pest entries.
- j) For **Sweet Corn**, add the missing horizontal lines in the **Pests**, **Rate** and **Timing/Special Directions** columns between the four pest entries.
- k) For **Tomatoes**, add the missing horizontal lines in the **Pests**, **Rate** and **Timing/Special Directions** columns between the three pest entries.
- l) For **Turnips and Turnip Tops**, add the missing horizontal lines in the **Pests**, **Rate** and **Timing/Special Directions** columns between the two pest entries.
- m) In the **Disposal** text, replace the first sentence with:

To avoid harming aquatic organisms in rivers and other surface waters, do not pour this concentrate, spray mixture, or rinse water into sanitary drains (for example, toilets and sinks) or storm sewers (for example, street drains).
- n) Also in the **Disposal** text, replace the last sentence with:

If disposal of a partially full container is necessary, take the container to your local household waste disposal facility, or make sure the cap is tightly closed, securely wrap the container in several layers of newspaper, and discard in trash.
- o) At the top of the "Adhesive Panel" label, revise the claim "and many others" to "and many others as listed in directions".

p) On the "Adhesive Panel" correct the EPA Reg. No. to read "655-799", rather than "655-".

3. Submit three (3) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager 13
Insecticide/Rodenticide Branch
Registration Division (7505C)

40116

Prentox Diazinon Lawn & Garden Insecticide,
EPA Reg. No. 655-799

Front Panel - Booklet Label

PRENTOX® DIAZINON LAWN & GARDEN INSECTICIDE

Water-Based Concentrate

This product must not be used on golf courses and sod farms. For outdoor use on the home lawn and the home garden only.
Spray concentrate. Contains diazinon insecticide.

Protects Lawns, Vegetables, Fruits and Nuts, Roses, Flowers, Trees, Shrubs

Controls Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants¹, Flea Beetles, Grubs, Mites, Sod Webworms and many others.
¹ Aids in control

ACTIVE INGREDIENT:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 22.4%

INERT INGREDIENTS: 77.6%

Total: 100.0%

PRENTOX® — Registered Trademark of Prentiss Incorporated

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 655-799
Manufactured for:

EPA EST NO.
NET CONTENTS _____

PRENTISS INCORPORATED

Plant: Kaolin Road, Sandersville, GA 31082
Office: C.B. 2000, Floral Park, NY 11002-2000

ACCEPTED
with COMMENTS
in EPA Letter Dated

FEB 18 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
655-799

96 AUG 21 AIO:19

RECD EPA/OPP/DPD1

5016

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Prentiss Incorporated or the Seller. All such risks shall be assumed by the Buyer.

Prentiss Incorporated warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. Prentiss Incorporated makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Prentiss Incorporated or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Prentiss Incorporated and the Seller offer this product and, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Prentiss Incorporated.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

Pretox Diazinon Lawn & Garden Insecticide is a water-based concentrate product which must be diluted with water before application. To avoid having spray mixture left over, mix only the amount needed. Refer to the section How to Mix and Apply. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruits and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. Pretox Diazinon Lawn & Garden Insecticide works fast and applications should be repeated only as directed to maintain control.

Notes: (1) Do not use inside the home. (2) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia, or orchids, since injury may occur. (3) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (4) Do not apply to food crops grown in greenhouses.

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

How to Mix and Apply

Step #1: Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

Step #2: Use clean sprayer. Carefully measure and mix the amount of Pretox Diazinon Lawn & Garden Insecticide and water as indicated in the rate chart.

Step #3: Spray as directed in rate chart. Thorough coverage is important. See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

Step #4: Flush sprayer with clean water after each use.

Measurement Conversions

1 tablespoon = 3 teaspoons
1 fl. oz. = 6 teaspoons or 2 tablespoons
8 fl. oz. = $\frac{1}{2}$ cup
1 $\frac{1}{2}$ tablespoons in 3 gals. of water = 1 $\frac{1}{2}$ teaspoons in 1 gal. of water
2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

Lawn Pest Control

Precautions

1. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.
3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl.
4. Avoid overlapping of sprays.
5. Where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least $\frac{1}{4}$ inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Marion, and St. Augustinegrass.

60916

To control pests listed below, apply recommended rates.

Pests	Rate Diazinon Lawn & Garden Insecticide	Water	Remarks
Lawn Chinch Bugs	1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Thoroughly wet down grass a few hours before applying. In the North, if Chinch Bugs are the only problem, a lower rate of application can be used: 3 tea-spoons in 3 gals. of water covers 125 sq. ft. or 4 fl. oz. in 24 gals. of water per 1,000 sq. ft.
Ants, Armyworms, Bermudagrass Mites, Clover Mites, Springtails (<i>Collembola</i>), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs)	1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Thoroughly wet down grass a few hours before applying. For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active. Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect. For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.
Brown Dog Ticks, Chiggers, Fleas	1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Note: Do not apply this product to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Hyperodes Weevil	1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application.
Millipedes	1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Thoroughly wet grass a few hours before applying.
Rhodesgrass Scales	1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Apply when crawlers first emerge.
White Grubs of Japanese Beetle, European Chafer, Southern Chafer	1½ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Apply any time between late July and early October. Water grass thoroughly after application.

70116

Prentox Diazinon Lawn & Garden Insecticide,
EPA Reg. No. 635-799

Nuisance Pests Outside the Home	
Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (<i>Collembola</i>)	<p>Mix 2 fl. oz. in 1 gal. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.</p> <p>To prevent entry into houses, spray a 5 ft. band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 ft. Repeat application as necessary.</p> <p>Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate.</p>
Fire Ants	<p>To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.</p> <p>Note: For best results, apply in cool weather 65-80° F, or in early morning or late evening hours. Treat new mounds as they appear.</p>

Insect Control on Ornamental Trees, Shrubs, and Flowers
Spray when listed pests first appear. Repeat as necessary.

Trees and Shrubs	Insects	Rate per 1 Gal. of Water	Timing/Special Directions
Arborvitae, Azaleas, Birch, Boxwood, Camellias, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Scotch Pine, Spirea, Spruce, Sycamore, Wax Plant (<i>Hoya</i>), Willow, Yellowwood, Yew	Aphids, Apple-and-Thorn Skeletonizers, Bagworms, European Pine Sawfly, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak) Hemlock Chermes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Oak Loopers, Obliquebanded Leaf-rollers, Obscure Root Weevils, Omnivorous Leaf-tiers, Pear Slugs, Scale Crawlers (Cottony-cushion, Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft), Tent Caterpillars, Webworms (Cotoneaster, Fall, Juniper, Mimosa), Whiteflies	2 teaspoons	<p>Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage with spray.</p> <p>Mimosa Webworms: apply when adults first appear. Make 2 additional applications at 4 - 5 week intervals.</p>

Roses and Flowers	Insects	Rate per 1 Gal. of Water	Timing/Special Directions
Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses	Aphids, Dipterous Leafminers, Euonymus Scale Crawlers, Flea Beetles, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies	2 teaspoons	Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.

80116

Pretox Diazinon Lawn & Garden Insecticide,
EPA Reg. No. 655-799

Fruits and Nuts

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted.

Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

Crop	PHI*
Almonds	**
Apples	21
Apricots	21
Blackberries	7
Blueberries	7
Boysenberries	7
Cherries	21
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Pears	21
Plums	21
Prunes	21
Raspberries	7
Strawberries	-5
Walnuts	45

*PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

**Dormant spray only.

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

Crop	Pests	Rate	Timing/Special Directions
Almonds (CA only)	San Jose Scale, Parlatoria Scale	2 teaspoons in 1 gal. of water	Apply only as a dormant spray. Do not apply more than 6 gals. per 500 sq. ft. of home orchard. This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.
Apples	Aphids, Apple maggots, Codling moths, Eyespotted budmoths, Fruit Tree leafrollers, Leafhoppers, Mealybugs, Scale crawlers (San Jose, Forbes) Spider mites	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest. Note: Resistant apple maggots may not be controlled by diazinon or similar type insecticides.
Apricots	Aphids, Apricot Mealybugs, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest.
Blueberries	Aphids, Blueberry Maggots, Cherry Fruitworms, Cranberry Fruitworms, Thrips, Two-spotted Spider Mites	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of blueberry patch. Do not make more than 5 applications per season, a minimum of 14 days apart. Do not apply within 7 days of harvest.

10/16

Crop	Pests	Rate	Timing/Special Directions
Caneberries, (CA, OR, WA only): Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries	Aphids, Leafhoppers, Raspberry Fruitworms, Two-spotted Spider Mites, Thrips	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 5 gals. per 500 sq. ft. of caneberry patch. Do not make more than 5 applications per season, a minimum of 14 days apart. Do not apply within 7 days of harvest. This product may be used on caneberries only in CA, OR, and WA. Do not use this product on caneberries grown in other states.
Cherries	Black Cherry Aphids, Cherry Fruit- flies, Cherry Fruitworms (Maggots), Fruitree Leafrollers, Mites (Cherry Rust), Scale Crawlers (San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season, a minimum of 7 days apart. Do not apply within 21 days of harvest.
Grapes	Aphids, Grape Berry Moths, Grape Leafrollers, Leafhoppers	1½ teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 3 gals. per 500 sq. ft. of home vineyard. Do not make more than 5 applica- tions per season, a minimum of 7 days apart. Do not apply within 28 days of harvest.
Nectarines	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest.
Peaches	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season, a minimum of 14 days apart. Do not apply within 21 days of harvest.
Pears	Aphids, Apple maggots, Codling moths Mealybugs, Scale crawlers (San Jose Forbes), Spider Mites, Tentiform leafminers	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season, a minimum of 14 days apart. Do not apply within 21 days of harvest. Note: Resistant apple maggots will not be controlled by diazinon or similar type insecticides.
	Pear Psylla	1 - 2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Plums and Prunes	Aphids (Leafcurl Plum, Mealy Plum, Thistle), Mites (Brown, European Red)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season, a minimum of 7 days apart. Do not apply within 21 days of harvest.
Strawberries	Aphids, Two-spotted Spider Mites, Strawberry Leafrollers	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 3 gals. per 500 sq. ft. of strawberry patch. Do not make more than 3 applications per season, a minimum of 7 days apart. Do not apply within 5 days of harvest.
Walnuts (CA only)	Aphids, Scale Crawlers (Walnut, Italian Pear, Frosted), Spider Mites, Twig Borers	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 6 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest. This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.

100916

Vegetables

For foliar or soil applications, Pretox Diazinon Lawn & Garden Insecticide should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil.

Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Intervals (PHI*) for Foliar Applications

Crop	PHI* (Days)
Beans ¹ (succulent varieties only: Lima, Pole and Snap)	7
Broccoli	7
Cabbage	21
Cantaloupes	3
Carrots	14
Casabas	3
Cauliflower	7
Chinese Broccoli, Chinese Cabbage, Collards	10
Crenshaws	3
Cucumbers	7
Endive (Escarole)	14
Honeydew Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Onions (Bulb and Green), Parsnips	14
Peas ¹	7
Peppers	5
Persian Melons	3
Potatoes (Irish)	35
Radishes, Rutabagas, Spinach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Tomatoes	1
Turnips (Roots and Tops)	14
Watermelons	3

*PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

¹ Do not use this product on dried pea or bean varieties, such as pinto beans, dried limas, split peas, or dried black-eyed peas.

11/2/16

Vegetables

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

Crop	Pests	Rate	Timing/Special Directions
Beans, Lima, Pole and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days, up to 3 applications per season. Do not apply within 7 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms (snap beans only)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
	Note: (1) Do not apply this product to dried bean varieties, such as pinto beans or dried limas. (2) To protect bees, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom.		
Beets, Red (Table)	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 21 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Root Maggots	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.

20/16

Prentox Diazinon Lawn & Garden Insecticide,
EPA Reg. No. 655-799

Crop	Pests	Rate	Timing/Special Directions
Carrots	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Cauliflower	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Chinese Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticide.
Chinese Cabbage (Bok Choy and Napa)	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Collards	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Corn (see sweet corn)			
Cucumbers	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 7 days of harvest.
Endive (Escarole)	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

130416

Prentox Diazinon Lawn & Garden Insecticide,
EPA Reg. No. 655-799

Crop	Pests	Rate	Timing/Special Directions
Kale	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Lettuce	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians and Hybrids of these) and Watermelons	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 3 days of harvest.
Onions (Bulb and Green)	Onion Thrips	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest.
Parsnips	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat applications every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Peas (succulent only)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days, up to 3 applications per season. Allow at least 7 days between the last application and harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Notes: (1) Do not apply this product to dried pea varieties, such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom.			

14/01/16

Crop	Pests	Rate	Timing/Special Directions
Peppers	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 5 days of harvest.
Potatoes	Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 35 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Radishes	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 3 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
Rutabagas	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Squash, Summer	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 7 days of harvest.
Squash, Winter	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 3 days of harvest.

150416

Crop	Pests	Rate	Timing/Special Directions
Sweet Corn	Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Corn may be picked 7 days following last application.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Note: To protect bees, do not apply to corn foliage during the pollen shed period.			
Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies (<i>Drosophila</i> spp.)	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 1 day of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Turnips and Turnip Tops	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

STORAGE AND DISPOSAL

Storage: Store in original container in an area inaccessible to children and pets. If product freezes, place container in a warm area (68° F or above) inaccessible to children or pets and allow to thaw. Shake well before using.

Disposal: Do not pour the concentrate, the spray mixture, or the rinse water down the drain or sewer. Do not reuse empty container. Triple rinse the empty container, pour the rinse water into the sprayer, then spray the area to be treated. Securely wrap empty container in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

160916

Prentox Diazinon Lawn & Garden Insecticide,
EPA Reg. No. 655-799

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/2 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Adhesive Panel

PRENTOX® DIAZINON LAWN & GARDEN INSECTICIDE

Water-Based Concentrate

This product must not be used on golf courses and sod farms. For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

Protects Lawns, Vegetables, Fruits and Nuts, Roses, Flowers, Trees, Shrubs

Controls Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants¹, Flea Beetles, Grubs, Mites, Sod Webworms and many others.

¹ Aids in control

ACTIVE INGREDIENT:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 22.4%

INERT INGREDIENTS: 77.6%

Total: 100.0%

PRENTOX® — Registered Trademark of Prentiss Incorporated

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.

SEE END OF BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 655-

Manufactured for:

EPA EST NO.

NET CONTENTS _____

PRENTISS INCORPORATED

Plant: Kaolin Road, Sandersville, GA 31082

Office: C.B. 2000, Floral Park, NY 11002-2000