

Wyandotte™

STERI-CHLOR

ALL-SOLUBLE CHLORINE SOLUTION

CAUTION

**KEEP OUT OF REACH OF CHILDREN
MAY BE HARMFUL IF SWALLOWED**

Undiluted product may cause skin irritation. Avoid contact with eyes. In case of contact, flush with plenty of water. If irritation persists, get medical attention. Avoid breathing dust. Avoid contamination of food.

REPLACE COVER. STORE IN COOL, DRY PLACE.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

DIRECTIONS

GLASSES, DISHES, UTENSILS

1. Scrape and pre-flush—articles to be washed.
2. Wash—in clean water using recommended Wyandotte product.

3. Rinse—in clean running water.
4. Sanitize—in Wyandotte Steri-Chlor solution using 2 ounces in each 6 gallons water (100 p.p.m. Av. Cl.). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code. Test periodically with chlorine tester. When solution depreciates to 50 p.p.m. Av. Cl., discard.
5. Air-dry on clean drainboard.


DAIRY, BEVERAGE, FOOD PROCESSING EQUIPMENT— FARM TANKS, PAILS, STRAINERS, MILKING MACHINE PARTS

- A. Clean equipment immediately after use.
 1. Rinse—with clean water.
 2. Wash—with recommended Wyandotte product.
 3. Rinse—with clean water.
 4. Air-dry—protect from dust and dirt.

FOR INDUSTRY USE ONLY

350 LBS. NET WEIGHT

BASF



BASF Wyandotte Corporation
Chemical Specialties Division
Wyandotte, Michigan 48192

Wyandotte™

STERI-CHLOR

4-13-72
662-29

SOLUBLE CHLORINE SANITIZER

EPA REG. NO. 662-29

IF CHILDREN SWALLOWED

irritation. Avoid contact, flush with water, get medical attention, avoid contamination.

DRY PLACE.

to drum reconditioning or crushing and

washed. Recommended Wyandotte

3. Rinse—in clean running water.
4. Sanitize—in Wyandotte Steri-Chlor solution using 2 ounces in each 6 gallons water (100 p.p.m. Av. Cl.). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code. Test periodically with chlorine tester. When solution depreciates to 50 p.p.m. Av. Cl., discard.
5. Air-dry on clean drainboard.

DAIRY, BEVERAGE, FOOD PROCESSING EQUIPMENT—FARM TANKS, PAILS, STRAINERS, MILKING MACHINE PARTS

- A. Clean equipment immediately after use.
 1. Rinse—with clean water.
 2. Wash—with recommended Wyandotte product.
 3. Rinse—with clean water.
 4. Air-dry—protect from dust and dirt.

- B. Sanitize immediately before re-use with Wyandotte Steri-Chlor solution using 2 ounces in each 6 gallons water (100 p.p.m. Av. Cl.). Test periodically with chlorine tester. When solution depreciates to 50 p.p.m. Av. Cl. discard.

Wyandotte Steri-Chlor fulfills the criteria of Appendix F, the Grade "A" Pasteurized Milk Ordinance 1965 recommendations of the United States Public Health Service when tested by the Chambers Method.

Do not rinse after sanitizing with Wyandotte Steri-Chlor solution unless required by local Health Authorities, then rinse with clean, safe water.

ACTIVE INGREDIENT

[Mono (trichloro) tetra (monopotassium dichloro)]-penta-s-triazinetriene	6.0%
INERT INGREDIENTS	94.0%
	<hr/> 100.0%

FOR INDUSTRY USE ONLY

350 LBS. NET WEIGHT




BASF Wyandotte Corporation
 Chemical Specialties Division
 Wyandotte, Michigan 48192



Wyandotte

STERI-CHLOR[®]

ALL-SOLUBLE CHLORINE SANITIZER



EPA REG. NO. 662-29

CAUTION — KEEP OUT OF REACH OF CHILDREN. MAY BE HARMFUL IF SWALLOWED.

Undiluted product may cause skin irritation. Avoid contact with eyes. In case of contact, flush with plenty of water. If irritation persists, get medical attention. Avoid contamination of food.

Replace Cover. Store in Cool, Dry Place.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

DIRECTIONS

GLASSES, DISHES, UTENSILS

1. Scrape and pre-flush—articles to be washed.
2. Wash—in clean water using recommended Wyandotte product.

3. Rinse—in clean running water.
4. Sanitize—in Wyandotte Steri-Chlor solution using 2 ounces in each 6 gallons water (100 p.p.m. Av. Cl.). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code. Test periodically with chlorine tester. When solution depreciates to 50 p.p.m. Av. Cl. discard.

5. Air-dry on clean drain board.

DAIRY, BEVERAGE, FOOD PROCESSING EQUIPMENT—FARM TANKS PAILS, STRAINERS, MILKING MACHINE PARTS

- A. Clean equipment immediately after use.
 1. Rinse—with clean water.
 2. Wash—with recommended Wyandotte product.
 3. Rinse—with clean water.

4. Air-dry—protect from dust and dirt.
- B. Sanitize—immediately before re-use with Wyandotte Steri-Chlor solution using 2 ounces in each 6 gallons water (100 p.p.m. Av. Cl.). Test periodically with chlorine tester. When solution depreciates to 50 p.p.m. Av. Cl. discard.

Wyandotte Steri-Chlor fulfills the criteria of Appendix F, the Grade "A" Pasteurized Milk Ordinance 1965 recommendations of the United States Public Health Service when used by the Chambers Method.

ACTIVE INGREDIENT

[Mono (trichloro) tetra (monopotassium dichloro)]-penta-s-triazinetriene . . . 6.0%

INERT INGREDIENTS 94.0%

100.0%

FOR INDUSTRY USE ONLY

50 LBS. NET WEIGHT

BASF



BASF Wyandotte Corporation
Chemical Specialties Division
Wyandotte, Michigan 48192