



Wyandotte

ANTIBAC[®] B

UNDER THE FEDERAL
FUNGICIDE AND
FOR ECONOMIC POISON
ED UNDER NO. 662

ALL-SOLUBLE CHLORINE SANITIZER-GERMICIDE DISINFECTANT DEODORANT

CAUTION

**KEEP OUT OF REACH OF
CHILDREN**

**MAY BE HARMFUL
IF SWALLOWED**

Undiluted product may cause skin irritation. Avoid contact with eyes. In case of contact, flush with plenty of water. If irritation persists, get medical attention.

Avoid contamination of food.

ACTIVE INGREDIENT

Sodiumdichloro-(s)-triazinetrione . . . 25.8%

INERT INGREDIENTS*

74.2%

*Includes buffering agents.

100.0%

U.S.D.A. Registration No. 662-41

DIRECTIONS FOR USE

Glasses, Dishes, Utensils

1. Scrape and pre-flush—articles to be washed.
2. Wash—in clean water using recommended Wyandotte product.
3. Rinse—in clean running water.
4. Sanitize—in ANTIBAC B solution using 1/4 oz. to each 3 gallons water (100 ppm available chlorine). Test periodically with chlorine tester. When the available chlorine decreases to 50 ppm, discard.

CONTACT TIME: The laws of some states require that utensils and equipment remain in contact with germicidal solution for at least 2 minutes. Check with your Health Department.

5. Air-dry—on clean drain board.

Dairy, Beverage, Food Processing Equipment, Farm Tanks, Pails, Strainers, Milking Machine Parts.

- A. Clean equipment immediately after use.

1. Rinse—with clean water.
2. Wash—with recommended Wyandotte product.
3. Rinse—with clean water.
4. Air-dry—protect from dust and dirt.

- B. Sanitize—immediately before re-use with ANTIBAC B solution using 1 oz. of ANTIBAC B in each 12 gallons water (100 ppm). Test periodically with chlorine tester. When the available chlorine decreases to 50 ppm discard.

Sanitizing Eggs

1. Wash eggs with approved Wyandotte product.
2. Sanitize eggs in spray rinse containing 100 ppm available chlorine (1 oz. ANTIBAC B in 12 gallons of water).

For Deodorizing and Disinfecting

Refrigerators, Walls, Floors, Cold Storage Rooms, Refuse Containers, Dog Kennels, Poultry Houses, etc. Use 1 ounce to each 6 gallons of water (200 ppm available chlorine).

Replace cover. Store in cool, dry place.

ANTIBAC B fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance, 1965 recommendations of the U.S. Public Health Service, when tested by the method outlined by Chambers.

50 LBS. NET WEIGHT

CREATIVE RESEARCH • QUALITY PRODUCTS • PROFESSIONAL SERVICE

Wyandotte
CHEMICALS CORPORATION

J. B. FORD DIVISION
WYANDOTTE, MICHIGAN 48192

Wyandotte

ANTIBAC B

ALL-SOLUBLE CHLORINE SANITIZER-GERMICIDE
DISINFECTANT DEODORANT

662

CAUTION

KEEP OUT OF REACH OF
CHILDREN

MAY BE HARMFUL
IF SWALLOWED

Undiluted product may cause skin
irritation. Avoid contact with eyes. In
case of contact, flush with plenty of
water. If irritation persists, get medical
attention.

Avoid contamination of food.

ACTIVE INGREDIENT

Sodium dichloro (s-triazine) trihydrate 25.8%

INERT INGREDIENTS*

Includes buffering agents 74.2%

U.S.D.A. Registration No. 662-41

DIRECTIONS FOR USE

Glasses, Dishes, Utensils

1. Scrape and pre-flush—articles to be washed.
2. Wash—in clean water using recommended Wyandotte product.
3. Rinse—in clean running water.
4. Sanitize—in ANTIBAC B solution using 4 oz. to each 3 gallons water (100 ppm available chlorine). Test periodically with chlorine tester. When the available chlorine decreases to 50 ppm, discard.

CONTACT TIME The laws of some states require that utensils and equipment remain in contact with germicidal solution for at least 2 minutes. Check with your Health Department.

5. Air-dry—on clean drain board.

Dairy, Beverage, Food Processing Equipment Farm Tanks, Pails, Strainers, Milking Machine Parts

- A. Clean equipment immediately after use.
 1. Rinse—with clean water.
 2. Wash—with recommended Wyandotte product.
 3. Rinse—with clean water.
 4. Air-dry—protect from dust and dirt.

- B. Sanitize—immediately before re-use with ANTIBAC B solution using 1 oz. of ANTIBAC B in each 12 gallons water (100 ppm). Test periodically with chlorine tester. When the available chlorine decreases to 50 ppm, discard.

Sanitizing Eggs

1. Wash eggs with approved Wyandotte product.
2. Sanitize eggs in spray rinse containing 100 ppm available chlorine (1 oz. ANTIBAC B in 12 gallons of water).

For Deodorizing and Disinfecting

Refrigerators, Walls, Floors, Cold Storage Rooms, Refuse Containers, Dog Kennels, Poultry Houses, etc. Use 1 ounce to each 6 gallons of water (200 ppm available chlorine).

Replace cover. Store in cool, dry place.

ANTIBAC B fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance, 1965 recommendations of the U.S. Public Health Service, when tested by the method outlined by Chambers.

50 LBS. NET WEIGHT

CREATIVE RESEARCH • QUALITY PRODUCTS • PROFESSIONAL SERVICE

Wyandotte
CHEMICALS CORPORATION

J. B. FORD DIVISION
WYANDOTTE, MICHIGAN 48192