

662-43

Wyandotte PER-VAD fulfills the criteria of Appendix F of the Grade A Pasteurized Milk Ordinance, 1965, recommendations of the U. S. Public Health Service when tested by the Chambers Modification of the Weber-Black method in water up to 1000 ppm hardness (58 grains per gallon as CaCO₃).

IMPORTANT—Store at room temperature. If product becomes chilled, bring back to 60° F., before use.

ACCEPTED

May 26, 1971

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 662-43

U.S.D.A. Reg. No. 662-43

Wyandotte

PER-VAD

TRADEMARK

LOW FOAM LIQUID ACID SANITIZER

ACTIVE INGREDIENTS:

Orthophosphoric Acid	15.0%
Sulfonated Oleic Acid, Sodium Salt	2.6%
Inert Ingredients	82.4%
	<hr/> 100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN.

HARMFUL IF SWALLOWED. Causes skin or eye irritation. Avoid contact with skin and eyes. In case of contact, flush skin or eyes immediately with plenty of water. For eyes, get medical attention. Avoid contamination of food.

CONTENTS ONE GALLON

Wyandotte

CHEMICALS CORPORATION

J. B. FORD DIVISION

WYANDOTTE, MICHIGAN 48192

USE DIRECTIONS

Dairy, Beverage, Brewery, Food Processing Equipment.

Clean Equipment Immediately After Use.

1. Rinse with clean water.
2. Wash with recommended Wyandotte product.
3. Rinse with clean water.
4. Air dry—protect from dust and dirt.
5. Sanitize—before reuse with one fluid ounce of Wyandotte PER-VAD per gallon of water. Test periodically with Wyandotte PER-VAD Test Kit. When solution depreciates to 100 ppm, discard.
6. Drain for 15 minutes prior to use in contact with food.

Contact Time: The laws of some states require that utensils and equipment remain in contact with sanitizing solution for at least 2 minutes. Check with your Health Department.

05/26/1971

11