

**DIRECTIONS FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT**

Clean equipment in the normal manner. Just before using rinse all surfaces thoroughly with 200 ppm of Sodium Hypochlorite. Maintain contact with disinfectant for a minimum of 2 minutes. Do not rinse with water after Chlorine treatment. Do not soak overnight. Use Sodium Hypochlorite according to the Dilution Chart below.

**USE DILUTIONS**

Gallons Water	Ozs. Sodium Hypochlorite		
	50 ppm av. Cl.	100 ppm av. Cl.	200 ppm av. Cl.
2	1/4	1/2	1

To feed Sodium Hypochlorite with a Topper Cub or Topper Cougar Pump, dilute 1 gal. Sodium Hypochlorite with 4 gals. water (1:4). This will permit feeding 100 ppm chlorine into a rinse flow of 1/2 gpm. At stronger concentrations and higher flows use less water.

Wyandotte™

# SODIUM HYPOCHLORITE

**Cleans • Disinfects • Deodorizes and Bleaches**

**ACTIVE INGREDIENT**

Sodium Hypochlorite..... 5.25% BY WT.

**INERT INGREDIENTS**..... 94.75% BY WT.

**WARNING**

**KEEP OUT OF REACH OF CHILDREN**

**SEE WARNING TEXT ON RIGHT SIDE PANEL**

**RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD IT**

**CONTENTS ONE GALLON**

Sold by

**BASF**



**BASF Wyandotte Corporation** ::  
**Chemical Specialties Division** ::  
 Wyandotte, Michigan 48192

## WARNING

KEEP OUT OF REACH OF CHILDREN.  
HARMFUL IF SWALLOWED.

Skin and Eye Contact May Cause Severe Irritation.  
Avoid Contact with Skin and Eyes.

## ANTIDOTE

**INTERNAL**—Drink mucilage, raw egg white, milk or rice gruel. Follow with emetic (tablespoon of mustard in tumbler of warm water).

**EXTERNAL**—Wash with water.

**ACCEPTED**

MAY 22 1972

DATE PACKAGED:

EPA Reg. No. 662-50

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 662-50



for us



**POWER**  
Kills a broad spectrum of  
equipment and  
supplies

**EASY T**  
Wyandotte  
fed to m  
dissolv

**ECONO**  
Only sp  
One out  
tion. Or  
200 ppm

