

Wyandotte™

# STERI-CHLOR® D

## Highly-Soluble Chlorine Sanitizer

**ACCEPTED**  
 662-60  
 JUL 8 1970  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTRATION  
 UNDER NO. \_\_\_\_\_ SUBJECT  
 TO ATTACHED COMMENTS.

**WARNING—KEEP OUT OF REACH OF CHILDREN. MAY BE HARMFUL IF SWALLOWED.**

Undiluted product may cause skin irritation. Avoid contact with skin or eyes. Avoid contamination of food.

**FIRST AID**—In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician immediately.

If swallowed, drink milk, egg white, starch paste, milk of magnesia, aluminum hydroxide gel, or magnesium trisilicate gel. Call a physician. Avoid sodium bicarbonate because of the release of carbon dioxide. Do not use acid first aid.

Store in a cool dry place. Avoid steam pipes and damp areas. Replace cover after each use. Avoid exposure to heat or moisture. Do not keep in hot areas or on hot surfaces.

The empty container should be thoroughly rinsed with water and discarded.

**USE DIRECTIONS**

Terminal Sanitizing Rinse for Glasses, Dishes and Utensils

1. Scrape and pre-flush articles to be washed.
2. Wash with recommended Wyandotte brand product.
3. Rinse in potable water.
4. Sanitize in Wyandotte Steri-Chlor D solution using 1 ounce in 3 gallons of water (providing 100 ppm available chlorine). Immerse all articles for at least two minutes, or for contact time specified by

governing sanitary regulations. Test solution periodically with chlorine tester. When solution depreciates to 50 ppm available chlorine, it should be discarded.

**5. Air-dry on clean drain board.**

Sanitizing Dairy, Beverage and Food Processing Equipment

1. Rinse equipment with water immediately after use.
2. Wash with recommended Wyandotte brand product.
3. Rinse with potable water.
4. Air-dry, protect from dust and dirt.
5. Sanitize immediately before re-use, using a solution of 1 ounce of Steri-Chlor D in 3 gallons of water (this provides 100 ppm available chlorine). Test this solution periodically with chlorine test kits and when it depreciates to 50 ppm available chlorine, it should be discarded.

Do not rinse after sanitizing with Wyandotte Steri-Chlor D solution unless required by local health regulations; then rinse with potable water.

Sanitizing Counters, Tables and Food Preparation Surfaces

1. Scrape or pre-flush surfaces.
2. Clean with recommended Wyandotte brand product.
3. Rinse with potable water.
4. Sanitize immediately before use with Wyandotte Steri-Chlor D solution using 1 ounce in 3 gallons of water. When solution depreciates to 50 ppm available chlorine, it should be discarded.

**Special Applications**

1. Deodorizing—Use 2 ounces of Steri-Chlor D in one gallon of water to provide a solution containing 600 ppm available chlorine. Rinse or spray odor problem areas. After deodorizing, rinse treated food contact surfaces with potable water before reuse.
2. Garbage Disposal Units—Prepare solution containing 2 ounces of Steri-Chlor D in each gallon of water and freeze the solution in ice cube trays. Pour the ice cubes into garbage disposal unit and run unit, without water, until cubes are gone. This will clean, deodorize and sharpen blades.
3. Coolers and Refrigerators—Soak charcoal overnight in solution containing 2 ounces of Steri-Chlor D in each gallon of water. Place charcoal in corners of units to deodorize and help eliminate odor transfer.

**ACTIVE INGREDIENT**

Potassium dichloro-s-triazinetrione	6.0%
<b>INERT INGREDIENTS</b>	<b>93.1%</b>
Available Chlorine, min.	4%

Wyandotte Steri-Chlor D fulfills the criteria of Appendix F, the Grade "A" Pasteurized Milk Ordinance 1965 Recommendations of the USPHS when tested by the AOAC Germicidal and Detergent Sanitizer Test.

EPA Reg. No. 662-60 AA  
EPA Est. 662-MI-1

**FOR INDUSTRY USE ONLY**

### 50 LBS. NET WEIGHT

# BASF



BASF Wyandotte Corporation  
Chemical Specialties Division  
Wyandotte, Michigan 48192

Freight Classification: Disinfectant, O/T Medicinal

# Wyandotte™ **STERI-CHLOR D** Highly-Soluble Chlorine

## **WARNING—KEEP OUT OF REACH OF CHILDREN. MAY BE HARMFUL IF SWALLOWED.**

Undiluted product may cause skin irritation. Avoid contact with skin or eyes. Avoid contamination of food.

**FIRST AID**—In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician immediately.

If swallowed, drink milk, egg white, starch paste, milk of magnesia, aluminum hydroxide gel, or magnesium trisilicate gel. Call a physician. Avoid sodium bicarbonate because of the release of carbon dioxide. Do not use acid first aid.

Store in a cool dry place. Avoid steam pipes and damp areas. Replace cover after each use. Avoid exposure to heat or moisture. Do not keep in hot areas or on hot surfaces.

The empty container should be thoroughly rinsed with water and discarded.

### **USE DIRECTIONS**

Terminal Sanitizing Rinse for Glasses, Dishes and Utensils

1. Scrape and pre-flush articles to be washed.
2. Wash with recommended Wyandotte brand product.
3. Rinse in potable water.
4. Sanitize in Wyandotte Steri-Chlor D solution using 1 ounce in 3 gallons of water (providing 100 ppm available chlorine). Immerse all articles for at least two minutes, or for contact time specified by

governing sanitary regulation periodically with available chlorine, it should be discarded.

5. Air-dry on clean drain board.  
Sanitizing Dairy, Beverage Processing Equipment

1. Rinse equipment with water after use.
2. Wash with recommended Wyandotte brand product.
3. Rinse with potable water.
4. Air-dry, protect from dust.
5. Sanitize immediately by using a solution of 1 ounce of product in 3 gallons of water (this provides 100 ppm available chlorine). Test periodically with chlorine test kit. When it depreciates to 50 ppm available chlorine, it should be discarded.

Do not rinse after sanitizing.  
Wyandotte Steri-Chlor D solution is approved by local health regulations for use as potable water.


Sanitizing Counters, Table Preparation Surfaces

1. Scrape or pre-flush surface.
2. Clean with recommended Wyandotte brand product.
3. Rinse with potable water.
4. Sanitize immediately by using 1 ounce of Wyandotte Steri-Chlor D in 3 gallons of water (providing 100 ppm available chlorine). When available chlorine depreciates to 50 ppm, it should be discarded.

**FOR INDUSTRY USE**

## **50 LBS. NET WT.**

# **BASF**



**BASF Wyandotte**  
Chemical Division  
Wyandotte, Michigan

Wyandotte<sup>TM</sup>  
STERI-CHLOR<sup>®</sup> D

Highly-Soluble Chlorine Sanitizer

FEATURES

- Easy to use
- Safe on metals
- Easy on hands
- Economical
- Dependable odor control

BENEFITS

Instantly soluble, makes clear solutions,  
leaves no films or residues


Safe on metals when used as directed

Buffered for controlled pH

Only 1 ounce in 3 gallons of water  
provides effective sanitizing

Steri-Chlor D eliminates odors, it does  
not just mask them

EPA Reg. No. 662-60 AA



Wyandotte STERI-CHLOR D

JUL 6 1974  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. SUBJECT  
TO ATTACHED COMMENTS.

Terminal Sanitizing Rinse for Glasses, Dishes and Utensils

1. Scrape and pre-flush articles to be washed.
2. Wash with recommended Wyandotte brand product.
3. Rinse in potable water.
4. Sanitize in Wyandotte Steri-ChlorD solution using 1 ounce in 3 gallons of water (providing 100 ppm available chlorine). Immerse all articles for at least two minutes, or for contact time specified by governing sanitary regulations. Test solution periodically with chlorine tester. When solution depreciates to 50 ppm available chlorine, discard.
5. Air-dry on clean drain board.

Sanitizing Dairy, Beverage and Food Processing Equipment

1. Rinse equipment with water immediately after use.
2. Wash with recommended Wyandotte brand product.
3. Rinse with potable water.
4. Air-dry, protect from dust and dirt.
5. Sanitize immediately before re-use, using a solution of 1 ounce of Steri-Chlor D in 3 gallons of water (this provides 100 ppm available chlorine). Test this solution periodically with chlorine test kits and when it depreciates to 50 ppm available chlorine, it should be discarded.

Do not rinse after sanitizing with Wyandotte Steri-Chlor D solution unless required by local health regulations; then rinse with potable water.

Sanitizing Counters, Tables and Food Preparation Surfaces

1. Scrape or pre-flush surfaces.
2. Clean with recommended Wyandotte brand product.

3. Rinse with potable water.
4. Sanitize immediately before use with Wyandotte Steri-Chlor D solution using 1 ounce in 3 gallons of water. When solution depreciates to 50 ppm available chlorine, it should be discarded.

#### Special Applications

1. Deodorizing – Use 2 ounces of Steri-Chlor D in one gallon of water to provide a solution containing 600 ppm available chlorine. Rinse or spray odor problem areas. After deodorizing, rinse food contact surfaces with potable water before reuse.
2. Garbage Disposal Units – Prepare solution containing 2 ounces of Steri-Chlor in each gallon of water and freeze the solution in ice cube trays. Pour the ice cubes into garbage disposal unit and run unit, without water, until cubes are gone. This will clean, deodorize and sharpen blades.
3. Coolers and Refrigerators – Soak charcoal overnight in solution containing 2 ounces of Steri-Chlor D in each gallon of water. Place charcoal in corners of units to deodorize and help eliminate odor transfer.

For technical assistance and recommendations concerning specific site problems, consult your Wyandotte Products representative.

#### Storage

Store Steri-Chlor D in a cool dry place. Avoid steam pipes and damp areas.

Replace cover after each use. Avoid exposure to heat or moisture.

Do not keep in hot areas or on hot surfaces.

The empty container should be thoroughly rinsed with water and discarded.