

Front

**BULK
LYSOL®
BRAND DISINFECTANT**

GERMICIDAL* Staphylocidal, Pseudomonacidal**, Tuberculocidal
FUNGICIDAL* (Against pathogenic fungi)
VIROCIDAL* (Against Influenza Virus Type A₁ [Japan], Herpes Simplex Type 1, Vaccinia, Adenovirus Type 2 and Human Immunodeficiency Virus Type 1 (HIV-1) (AIDS Virus))

*On Hard, Nonporous Surfaces **At 2% Solution

Prevents Mold and Mildew

ACTIVE INGREDIENTS: o-Phenylphenol, 2.80%, o-Benzyl-p-chlorophenol, 2.70%.

INERT INGREDIENTS: 94.50%.

EPA REGISTRATION NO. 675-19 EPA EST. NO. 777-OH-1

KEEP OUT OF REACH OF CHILDREN


WARNING: Harmful if swallowed. Causes eye and skin irritation. See Back Panel for Statement of Practical Treatment and Additional Precautionary Statements.

NET CONTENTS ONE GALLON

Alternate Sizes

**NET CONTENTS: 64 FL. OZ. (1.892 liters)
5 GALLONS
55 GALLONS**

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.



29 5 6

HOW TO USE BULK LYSOL® BRAND DISINFECTANT
(See Brochure for detailed instructions)

Bulk LYSOL Brand Disinfectant is recommended for use in hospitals, nursing homes, dental offices and other institutional facilities. Bulk LYSOL Brand Disinfectant may be used on floors, walls, bathroom fixtures and surgical instruments. (See brochure for additional information).

BULK LYSOL IS IN CONCENTRATED FORM. DILUTE ACCORDING TO DIRECTIONS.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

To prepare proper dilutions in required quantities, add full strength Bulk LYSOL to water as below.

FOR CLEANING OF SURGICAL AND DENTAL INSTRUMENTS:

Prior to disinfection instruments must be cleared to remove all adhering blood and serous exudates, body fluids and human wastes.

FOR DISINFECTION OF SURGICAL AND DENTAL INSTRUMENTS:

Immerse instruments in Bulk LYSOL Brand Disinfectant solution for 15 minutes to wet all surfaces thoroughly.

Flush and fill the lumen of hollow instruments/equipment, avoiding entrapment of air bubbles. If instruments are to remain in solution

for substantially longer periods of time, add 1 gram of sodium bicarbonate to each quart of solution; stir to dissolve completely. Also add 2 oz. of Isopropyl alcohol per quart to solution.

Fresh solution should be prepared at least daily or more often if the solution becomes diluted or soiled.

Correct
Solution
Strength


3%
(3:100)

BULK
LYSOL

WATER

30 cc.
4 oz.
12 oz.
20 oz.

1 quart
1 gallon
3 gallons
5 gallons




**FOR SANITARY DISPOSAL OF
URINE, FECES:**

To urine, add equal amount of Bulk LYSOL Brand Disinfectant solution. For feces, place solution in pan in amount slightly larger than expected discharge. After receiving feces, mix thoroughly with solution then cover and let stand for one hour before disposal.

**FOR SANITIZING LAUNDRY, LINENS,
BLANKETS AND DIAPERS:**

Add 1 cup of Bulk LYSOL Brand Disinfectant to each 17 gallons of water.

(See brochure for detailed instructions)



**BULK
LYSOL®
BRAND DISINFECTANT**

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Harmful if swallowed. Causes eye and skin irritation. Do not get in eyes. Avoid contact with eyes and skin. Avoid contamination of food.

STATEMENT OF PRACTICAL TREATMENT:

In case of eye contact, IMMEDIATELY flush eyes thoroughly with water, remove any contact lenses, and continue to flush eyes with plenty of water for at least 15 minutes. Get medical attention.

In case of skin contact, wash with soap and water. If irritation persists, consult a physician.

If swallowed, drink promptly a large quantity of water to dilute the product. Get immediate medical attention.

STORAGE AND DISPOSAL

64 fl. oz.
& 1 gallon

STORAGE: Store in original container in areas inaccessible to small children. Keep securely closed.

DISPOSAL: Do not reuse empty container. Rinse thoroughly with water and detergent. Wrap and discard in trash.

5 & 55
gallons


Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Store in original container in areas inaccessible to small children. Keep securely closed.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**NATIONAL LABORATORIES
Lehn & Fink Products Group
Montvale, NJ 07645**



HOW TO USE BULK LYSOL® BRAND DISINFECTANT

(See Brochure for detailed instructions)

BULK LYSOL IS IN CONCENTRATED FORM. DILUTE ACCORDING TO DIRECTIONS.

	Correct Solution Strength	To prepare proper dilutions in required quantities, add full strength Bulk LYSOL to water as below.	
		BULK LYSOL	WATER
FOR PREVENTING RUST IN HEAT DISINFECTION: Bulk Lysol Brand Disinfectant neutralizes the rusting action of boiling water.	1/2% (1:200)	20 cc. 2 oz.	1 gallon 3 gallons

FOR GENERAL DISINFECTION:

(including areas occupied by tubercular patients)


To disinfect hard non-porous surfaces, first clean surface thoroughly prior to application of Bulk LYSOL Brand Disinfectant. Apply solution to surface using a sponge or mop to wet all surfaces thoroughly. Let

stand 10 minutes before wiping or allow to air dry. Fresh solution should be prepared daily or more often if solution becomes diluted or soiled. For tuberculocidal activity 1% Bulk LYSOL Brand Disinfectant is effective in 10 minutes at 20°C (68°F).

FOR MOLD AND MILDEW ON HARD NON-POROUS SURFACES:
Wash surfaces with Bulk LYSOL solution; allow to air dry. Repeat application as necessary

FOR PSEUDOMONACIDAL ACTIVITY:
Clean surface thoroughly. Apply solution using a sponge or mop. Wet surface thoroughly. Let stand 10 minutes before wiping or let it air dry.

10 cc.	1 quart
40 cc.	1 gallon
4 oz.	3 gallons



Panel 4 (continued)

PURPOSE

FOR HIV-1 (AIDS VIROS) ACTIVITY:
KILLS HUMAN IMMUNODEFICIENCY
VIRUS TYPE 1 (HIV-1) (AIDS
VIRUS) ON PRECLEANED
ENVIRONMENTAL SURFACES/OBJECTS
PREVIOUSLY SOILED WITH BLOOD/
BODY FLUIDS.

SPECIAL INSTRUCTIONS FOR CLEANING
AND DECONTAMINATION AGAINST HIV-1
OF SURFACES/OBJECTS SOILED WITH
BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When
handling items soiled with blood
or body fluids, use disposable
latex gloves, gowns, masks or
eye coverings.

CLEANING PROCEDURES: Blood and
other body fluids must be
thoroughly cleaned from surfaces
and objects before application of
Bulk LYSOL Brand Disinfectant.

CONTACT TIME: Allow surface
to remain wet for 10 minutes.

DISPOSAL OF INFECTIOUS MATERIALS:
Blood and other body fluids should
be autoclaved and disposed of
according to local regulations for
infectious waste disposal.

Correct
Solution
Strength
For HIV-1
Activity

2 1/2
(2:100)

To prepare proper dilutions
in required quantities,
add full strength
Bulk LYSOL to water
as below.

BULK
LYSOL

80 cc.
8 oz.

WATER

1 gallon
3 gallons

ONE TABLESPOONFUL EQUALS ABOUT 1/2 FLUID OUNCE OR 15 CC.

