

09.093
07.101

GUARANTEE:

Modern manufacturing facilities and careful laboratory control assure materials of uniform quality at all times. National Laboratories' products are guaranteed to give complete satisfaction, when used as directed, or they may be returned for credit.



**Lehn & Fink Industrial Products Division
of Sterling Drug Inc.**
225 Summit Avenue, Montvale, N. J. 07645

Amphyl[®]

**DISINFECTANT
DEODORANT • DETERGENT**

PHENOL COEFFICIENT 10

GERMICIDAL, * staphylocidal, pseudomonacidal, tuberculocidal.

FUNGICIDAL* (Against pathogenic fungi)

VIRUCIDAL* (against most Lipophilic viruses such as Influenza A₂, Herpes simplex and Vaccinia)

***On Environmental Surfaces**

ACTIVE INGREDIENTS:

Soap, Anhydrous 44.0%, o-Phenylphenol 15.0%, p-tert-Amylphenol 6.3%, Alcohol 4.7%.

INERT INGREDIENTS: 30.0%

EPA REGISTRATION NO. 675-21-ZA

DANGER:

KEEP OUT OF REACH OF CHILDREN

See first aid and precautions on side panel.

NATIONAL LABORATORIES

**LEHN & FINK INDUSTRIAL PRODUCTS DIVISION
OF STERLING DRUG INC.**

Montvale, New Jersey 07645 Plant: Toledo, Ohio 43612

HOW TO USE HOSPITAL TYPE Amphyl[®]

ALWAYS DILUTE AMPHYL ACCORDING TO DIRECTIONS

To prepare proper dilutions in required quantities, add full strength Amphyl to water as below:

CORRECT SOLUTION STRENGTH	AMPHYL	WATER
---------------------------	--------	-------

FOR GENERAL DISINFECTION — (including areas occupied by tubercular patients) — Floors, walls, and furniture should be mopped, wiped, or scrubbed with the recommended solution to assure thorough wetting.

1/2%
(1:200)

5 cc.	1 quart
20 cc.	1 gallon
40 cc.	2 gallons

FOR PREVENTING RUST in heat sterilization—Amphyl prevents the rusting action of boiling water.

FOR DISINFECTION OF SURGICAL AND DENTAL INSTRUMENTS—Rinse off blood and exudates. Immerse instruments in Amphyl solution for 15 minutes. If Amphyl solution is to be stored for any period of time add 2 oz. of isopropyl alcohol to each quart of Amphyl solution.
FOR DISINFECTION OF RUBBER GOODS (e.g. tubing, gloves) —Rinse off blood and exudates. Immerse for 15 minutes. Remove, then immerse in 70% alcohol for one minute.

2%
(2:100)

20 cc.	1 quart
2 1/2 oz.	1 gallon
5 oz.	2 gallons

ONE TABLESPOONFUL EQUALS ABOUT 1/2 FLUID OUNCE OR 15 CC.

May be fatal or harmful if swallowed. Concentrate is corrosive to skin. Do not get concentrate or dilutions on skin or in eyes. Use rubber gloves for manual cleaning operations. Properly shield the eyes when handling the concentrate, in case of splashing. Avoid contamination of food. Do not get on clothing. Rinse empty container thoroughly with water and discard it.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, promptly drink a large quantity of milk, egg whites or gelatin solution; if these are not available drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

AB-557

ACCEPTED

MAY 28 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 675-21