

one operation
 Ntrol-eCON-omy
 lomonacidal, tuber-
 ungi)
 c viruses including
 ovirus Type 2 and
 enol 11.75%, o-Ben-
 m Ethylene Diamine
 ylbenzenesulfonate
 ntial Oils 0.30%.
 o phosphorous compounds.
 rosive. Causes eye
 get in eyes, on skin
 kin when handling.
 nation of food. Rinse
 er and discard it.
 diately flush eyes or
 5 minutes. For eyes,
 ntaminated clothing
 quantity of milk, egg
 not available, drink
 cian immediately.
 cosal damage may
 e. Measures against
 sion and convulsion

Con -O- Syl*

DISINFECTANT- DETERGENT

EPA REGISTRATION No. 675-27-AA

NET CONTENTS
 55 U.S. GALLONS


HOW TO USE Con-O-Syl HOSPITAL TYPE DISINFECTANT-DETERGENT

Con-O-Syl is economical, effective in 400 ppm hard water, noncombustible, and contains no phosphorous compounds. Con-O-Syl is in concentrated form. Dilute according to directions.

Purpose	Correct Solution Strength	To prepare proper dilutions in required quantities, add full strength Con-O-Syl to water as below:	
		Con-O-Syl	Water
FOR GENERAL CLEANING AND DISINFECTING	0.39% (1:256)	4 cc.	1 quart
		½ oz.	1 gallon
		1½ oz.	3 gallons
		2½ oz.	5 gallons

DIRECTIONS: No need to add soap or other detergents. Con-O-Syl does a complete job—cleans, disinfects, deodorizes. Avoid waste, dilute as directed.

High Dilution SAVES Time and Labor. See Brochure for detailed instructions. One tablespoon equals about ½ fluid ounce or 15 cc.



NATIONAL LABORATORIES

Lehn & Fink Industrial Products—Division of Sterling Drug Inc.
 Montvale, New Jersey 07645—Plant: Toledo, Ohio 43612

*TRADEMARK

3760-055