

CON-O-SYL**Disinfectant-Detergent**

Cleans, disinfects and deodorizes in one operation **CON**centrated
For **CON**venience-**CON**trol-**CON**omy

GERMICIDAL* staphylocidal, pseudomonacidal, tuberculocidal

FUNGICIDAL* (Against pathogenic fungi)

VIRUCIDAL* (Against most lipophilic viruses including Influenza
A₂, Herpes simplex, Adenovirus Type 2 and Vaccinia)

* On Hard, Non-porous Surfaces.

ACTIVE INGREDIENTS:	O-Phenylphenol	11.75%
	O-Benzyl-p-chlorophenol	7.82%
INERT INGREDIENTS:		<u>80.43%</u>
	Total	100.00%

EPA REGISTRATION No. 675.27

DANGER: Keep Out of Reach of Children.

See left panel for additional precautions.


NET CONTENTS ONE HALF GALLON

NATIONAL LABORATORIES

Lehn & Fink Products Group

Montvale, NJ 07645 . Plant: Toledo, Ohio 43612

Net Contents and Registration claims to be in accordance with
EPA's Pesticide Product Guidance for Phase 2
Pesticides, 40 CFR 155.10, Active to Inert Change in Status.



**FOR USE BY PROFESSIONAL MAINTENANCE PERSONNEL
ONLY IN INSTITUTIONAL AND INDUSTRIAL FACILITIES.**

**HOW TO USE Con-O-Syl HOSPITAL TYPE
DISINFECTANT-DETERGENT**


Con-O-Syl is economical, effective in 400 ppm hard water, noncombustible, and contains no phosphorous compounds. Con-O-Syl is in concentrated form. Dilute according to directions.

Purpose	Correct Solution Strength	To prepare proper dilutions in required quantities, add full strength Con-O-Syl to water as below:	
		Con-O-Syl	Water
FOR GENERAL CLEANING AND DISINFECTING	0.39% (1:256)	4 cc. 1/2 oz. 1 1/2 oz. 2 1/2 oz.	1 quart 1 gallon 3 gallons 5 gallons

DIRECTIONS:

No need to add soap or other detergents. Con-O-Syl does a complete job--cleans, disinfects, deodorizes. Avoid waste, dilute as directed.

High Dilution SAVES Time and Labor
See Brochure for detailed instructions.
One tablespoon equals about 1/2 fluid ounce or 15 cc.



DANGER:

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STORAGE/DISPOSAL:

Storage: Store in original container in areas inaccessible to small children, Keep away from food or feed. Disposal: Do not re-use empty container. Rinse thoroughly with water and detergent. Discard in trash.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

09 4
5 5
5 5
5 5