

P11 32 675-43

1075

**FRONT PANEL**

**PART OF YOUR INFECTION CONTROL PROGRAM**

For Use in Institutional and Industrial Facilities

**AMPHYL®  
DISINFECTANT • DEODORANT • DETERGENT**

<b>GERMICIDAL*</b>	Staphylocidal (including methicillin and gentamicin resistant staphylococcus), Pseudomonacidal, Tuberculocidal
<b>FUNGICIDAL*</b>	Against pathogenic fungi
<b>VIRUCIDAL*</b>	Against the following viruses — Herpes Simplex Type 1 Influenza A <sub>2</sub> (Japan), Vaccinia, and HIV-1 (AIDS Virus)

\*On environmental surfaces.

<b>ACTIVE INGREDIENTS:</b>	o-Phenylphenol.....	10.5%
	o-Benzyl-p-chlorophenol.....	5.0%
<b>INERT INGREDIENTS*:</b>	.....	84.5%

\*Contains no phosphates.

**KEEP OUT OF REACH OF CHILDREN**

**DANGER:**


Corrosive. Causes eye damage and skin irritation. See side/back panel for statement of practical treatment and additional precautionary statements.

EPA Reg. No. 675-43  
EPA Est. No. 777-OH-1

**1:200**

NATIONAL LABORATORIES (logo)  
L&F Products•Montvale, NJ 07645

Net Contents:  
6 Fl. Oz. (177 mL)  
32 Fl Oz. (1 Qt.) 946 mL  
1 U.S. Gallon (3.79 Liters)  
5 U.S. Gallons (18.95 Liters)



2075

**HOW TO USE AMPHYL® DISINFECTANT•DEODORANT•DETERGENT:**

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AMPHYL is recommended for use in hospitals, nursing homes, dental and doctors' offices, and other institutional facilities. It disinfects hard, nonporous surfaces such as wheelchairs, patient bed frames, bedside tables, medical lamps, stands, and other similar equipment.

**ONE TABLESPOONFUL EQUALS ABOUT ½ FLUID OUNCE OR 15 CC.**

AMPHYL is a concentrate. Always dilute in accordance with label instructions. Thoroughly preclean all soiled surfaces with AMPHYL prior to disinfecting, as described below.


) **NOTE:** Prepare fresh AMPHYL solution for each use. If a freshly prepared ½% AMPHYL solution is not to be utilized immediately, add 2 cc.(oz.) of Isopropyl alcohol to each quart of AMPHYL solution to maintain clarity.

To prepare proper dilutions in required quantities, add full strength AMPHYL to water as below:

<b>CORRECT SOLUTION STRENGTH</b>	<b>AMPHYL</b>	<b>WATER</b>
½% (1:200)	5cc (1 tsp.)	1 quart
	20cc (4 tsp.)	1 gallon
	40cc (1-1/3 oz.)	2 gallons

) **FOR GENERAL CLEANING:** Use a ½% solution for lightly soiled areas and a 1% solution for more difficult soil conditions.

**FOR GENERAL DISINFECTION (Including tuberculocidal effectiveness):** When required, remove heavy soil and gross filth prior to disinfection as per above cleaning instructions. Floors, walls and environmental surfaces should be mopped, wiped, or scrubbed with the recommended solution, maintaining a thoroughly wet surface for 10 minutes and allowed to air dry. Discard used solution daily or more often if soiling is evident. AMPHYL is effective in 10 minutes at 20°C (68°F).



**FOR DISINFECTION OF RUBBER GOODS (e.g. tubing, gloves):** All instruments and devices must be thoroughly cleaned to remove excess debris, rinsed and then rough dried before immersion. Clean and rinse the lumens of hollow instruments before filling with solution. Immerse in solution of AMPHYL for 10 minutes. Remove, then immerse in 70% alcohol for one minute. Discard used solution daily or more often if soiling is evident.

**This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical devices prior to sterilization or high level disinfection.**

) **FOR DISINFECTION OF SURGICAL AND DENTAL INSTRUMENTS:** All instruments and devices must be thoroughly cleaned to remove excess debris, rinsed and rough dried before immersion. Clean and rinse the lumens of hollow instruments before filling with solution. Immerse instruments in AMPHYL solution for 10 minutes. Discard used solution daily or more often if soiling is evident.

**KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) (AIDS VIRUS) ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of (HIV-1) (associated with AIDS).

) **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks or eye coverings.

**CLEANING PROCEDURES:** Blood or other body fluids must be thoroughly cleaned from surfaces and objects before application of AMPHYL Disinfectant.

**CONTACT TIME:** Allow surface to remain wet for 10 minutes.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

4-25

**BACK LABEL (PANEL #2)**

DISINFECTS AND DEODORIZES IN ONE OPERATION.

**PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**KEEP OUT OF REACH OF CHILDREN**

**DANGER:** Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Avoid contamination of food.

**STATEMENT OF PRACTICAL TREATMENT:**

In case of eye contact, IMMEDIATELY flush eyes thoroughly with plenty of water. Remove any contact lenses and continue to flush eyes with plenty of water for at least 15 minutes. Get medical attention. If on skin, wash with plenty of soap and water. If irritation persists, get medical attention. If swallowed, do not induce vomiting. Drink a large amount of water. Call a physician.

**STORAGE AND DISPOSAL**

(For sizes 1 gallon or less)

**STORAGE:** Store in original container in areas inaccessible to small children. Keep securely closed.

**DISPOSAL:** Do not reuse empty container. Rinse thoroughly with water and detergent. Discard in trash.

(For sizes larger than 1 gallon)

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container in areas inaccessible to small children. Keep securely closed.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**BACK LABEL (PANEL #2)**

Please encourage your state and local authorities to establish a recycling program so that this bottle/pail/drum can be recycled.

AMPHYL is part of a system of infection control products provided by National Laboratories. People respect the LYSOL Brand name for providing high quality and trusted surface disinfection, and gain the added benefit of a clean, sanitary environment. We invite you to call us at 1-800-677-9218, Monday through Friday, 8:45 a.m. to 4:30 p.m. (Eastern time), if you have any questions or comments on our products.

(For 6 oz. and 32 oz. size)

This label is printed on 50% recycled paper (10% post-consumer).

