

GENERAL USE DIRECTIONS

1. To clean, disinfect and deodorize walls, floors, tables, drinking fountains, sinks, refrigerators, doors, restroom fixtures, telephone booths, animal and bird cages, kennels, and garbage cans. Remove gross filth, heavy soils and grease, then wash with solution containing 1 ounce of Calgo San per gallon of water.

2. To clean and disinfect such articles as combs, brushes, razors, scissors, instruments and rubber goods. Wipe article clean and soak for ten minutes in a solution containing 2 ounces of Calgo San per gallon water.

3. To disinfect fabrics such as sheets, linens, aprons and uniforms. Soak for ten minutes in solution containing 1 ounce of Calgo San per gallon of water.

4. To control mold and mildew on surfaces such as floors, walls, woodwork and metal surfaces, remove gross filth, heavy soil and grease then apply 1 oz. of Calgo San per gallon of water and repeat the application when new growth appears or weekly.

Kills Trichophyton interdigitale (athletes foot fungi) on environmental surfaces when applied at 1 oz. per gallon.



Calgon Chemical Laboratories
Division of Calgon Corp.
Subsidiary of March & Co., Inc.
St. Louis, MO 63133

CALGO-SAN®

CLEANS DISINFECTS DEODORIZES
BACTERICIDAL FUNGICIDAL TUBERCULOCIDAL
VIRUCIDAL* PSEUDAMONACIDAL

A synthetic detergent with broad spectrum kill of both gram positive and gram negative Micro organisms for outstanding cleaning and disinfecting in one operation. An effective deodorizer for surfaces contaminated by bacterial putrefaction. No unpleasant residual perfume odor. *Inactivates suspensions of Influenza Type A/575 and Adenovirus Type 2 viruses.

ACTIVE INGREDIENTS

Oxalis Benzyl Para Chlorophenol 12.0%

INERT INGREDIENTS

88.0%

Nonflammable and nonvolatile in use dilutions. Excellent solubility in hot, cold, soft, or hard water. Contains biodegradable detergent. Dilute as directed. Do not mix with other cleaners.

DANGER: Keep out of reach of children. See First Aid and other precautions on right side panel. Do not use, pour, spill, or store near heat or open flame.

CONTENTS: 1 U.S. GAL., 7 FL. OZ. (4 LITERS)

HOSPITAL USE DIRECTIONS

1. To clean, disinfect and deodorize surfaces such as floors, walls, woodwork and metal surfaces, use 1 ounce of Calgo San per gallon of water.

2. To fog an unoccupied room as an adjunct either preceding or following regular cleaning and disinfecting procedures, use a dilution of 1/42 or 3 ounces of Calgo San per gallon of water per average size room. Allow two hours after fogging has stopped before entering room.

DANGER: Keep out of reach of children. Corrosive. Causes eye damage and severe skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Avoid contact with eyes and skin with diluted product. Wash thoroughly after using dilutions. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probably mucosal damage may contraindicate the use of gastric lavage. Measure against circulatory shock, respiratory depression and convulsion may be needed.

EPA Est. No. 8011M01
EPA Reg. No. 843-1-101

PRODUCT MADE IN U.S.A.

443

00 00 1

Not Reviewed. Registration of this to be in accordance with
Registration Guidance for Phase 2
Response, Page 11 and 12. Not Change in Status.

1/30/90

2422
1043-100
Pg. 1561

GERMICIDAL* . Tuberculocidal . Pseudomonacidal . Staphylocidal
VIRUCIDAL* (Against most lipophilic viruses including Herpes
simplex, Vaccinia, Influenza A₂ (Asian), and
Adenovirus Type 2)
FUNGICIDAL* (Against pathogenic fungi)
FUNGISTATIC* Prevents the growth of mold and mildew

*On hard non-porous surfaces in the presence of organic soil (5%
blood serum) and 400 ppm hard water (as CaCO₃,)

ACTIVE INGREDIENTS

o-Benzyl-p-chlorophenol	7.24%
o-Phenylphenol	2.23%
INERT INGREDIENTS	<u>90.53%</u>
TOTAL	100.00%

*Includes biodegradable detergents and contains no phosphorous
compounds

DANGER: KEEP OUT OF REACH OF CHILDREN

See Additional Precautions and Statement of Practical Treatment on
Side Panel


EPA Registration No. 675-46

NET CONTENTS: ONE GALLON (3.76 liters)

NATIONAL LABORATORIES
LEHN & FINK PRODUCTS GROUP
Montvale, NJ 07645 Plant Toledo, Ohio 43612


5.4.90

Not Reviewed. Registrant claims to be in accordance with
Reregistration of Pesticide Product Guidance for Phase 2
Response. Page 2.12, Active to Inactive Change in Status.



2nd side panel

AMERICA'S FOREMOST MANUFACTURER OF
DISINFECTANTS AND DISINFECTANT-DETERGENTS
FOR ENVIRONMENTAL SANITATION.



3RD PANEL

**HOW TO USE HOSPITAL TYPE
O-SYL
DISINFECTANT-DETERGENT**

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

O-SYL has been tested and found effective in hard water and in the presence of organic matter. O-SYL solution is in concentrated form, dilute only according to directions, following precautions indicated on this label.

PURPOSE

FOR GERMICIDAL CLEANING and DEODORIZING and CONTROL of MOLD and MILDEW (including premises occupied by tubercular patients).

To prepare proper dilution in required quantities, add full strength O-SYL to water as below

Prepare a fresh solution and apply to environmental surfaces. Maintain a thoroughly wet surface for 10 minutes. When required, remove gross filth and heavy soil prior to disinfection.

Correct
Solution
Strength

O-SYL

WATER

Floors - Walls - Carts -
Garbage Cans - Fixtures

0.78%
(1:128,
1 oz. per
gallon)

8 cc
1 oz.
3 oz.
5 oz.

1 quart
1 gallon
3 gallons
5 gallons

FLOORS Use O-SYL applied via mopping or spraying. Using a wet vacuum pick-up, rinsing not necessary


WALLS Wipe walls at least 8 ft. up from floor with O-SYL solution. Do not rinse.

FOR DISINFECTION OF SURGICAL DENTAL INSTRUMENTS AND THERMOMETERS. Rinse off gross blood and exudates. Immerse instruments for 10 minutes in O-SYL solution to maintain aseptic conditions. Rinse with sterile water. Discard used solutions daily or more often if soiling is evident.

FOR CONTROLLING MOLD and MILDEW Clean surface first, then apply via mop, cloth or sprayer until surface is moistened thoroughly. Do not rinse. Repeat as necessary.

EFFECTIVE IN HARD, SOFT AND SOILED WATER
TWO TABLESPOONFULS EQUALS ABOUT 1 FLUID OUNCE OR 30 CC

O-SYL saves time and labor. Cleans, Disinfects and Deodorizes in one operation.



3RD PANEL

HOW TO USE HOSPITAL TYPE
O-SYL
DISINFECTANT-DETERGENT

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

O-SYL has been tested and found effective in hard water and in the presence of organic matter. O-SYL solution is in concentrated form, dilute only according to directions, following precautions indicated on this label.

PURPOSE

FOR GERMICIDAL CLEANING and DEODORIZING and CONTROL of MOLD and MILDEW (including premises occupied by tubercular patients).

To prepare proper dilution in required quantities, add full strength O-SYL to water as below

Prepare a fresh solution and apply to environmental surfaces. Maintain a thoroughly wet surface for 10 minutes. When required, remove gross filth and heavy soil prior to disinfection.

Correct
Solution
Strength

O-SYL

WATER

Floors - Walls - Carts -
Garbage Cans - Fixtures

0.78%
(1:128,
1 oz. per
gallon)

8 cc
1 oz.
3 oz.
5 oz.

1 quart
1 gallon
3 gallons
5 gallons

FLOORS Use O-SYL applied via mopping or spraying. Using a wet vacuum pick-up, rinsing not necessary

WALLS Wipe walls at least 8 ft. up from floor with O-SYL solution. Do not rinse.

FOR DISINFECTION OF SURGICAL DENTAL INSTRUMENTS AND THERMOMETERS. Rinse off gross blood and exudates. Immerse instruments for 10 minutes in O-SYL solution to maintain aseptic conditions. Rinse with sterile water. Discard used solutions daily or more often if soiling is evident.

FOR CONTROLLING MOLD and MILDEW Clean surface first, then apply via mop, cloth or sprayer until surface is moistened thoroughly. Do not rinse. Repeat as necessary.

EFFECTIVE IN HARD, SOFT AND SOILED WATER

TWO TABLESPOONFULS EQUALS ABOUT 1 FLUID OUNCE OR 30 CC

O-SYL saves time and labor. Cleans, Disinfects and Deodorizes in one operation.

