

Front Panel

4/1/71
1/4
CME

**PREMEASURED
QUATSYL® 256
Detergent-Disinfectant-Deodorizer**

EPA Reg. No. 675-47 FFA Est. No. 46275-II-1

**FOR USE BY PROFESSIONAL MAINTENANCE PERSONNEL
ONLY IN INSTITUTIONAL AND INDUSTRIAL FACILITIES**

Water soluble packets provide germicidal activity in the presence of organic soil.*

BACTERICIDAL* Salmonella Choleraesuis, Staphylococcus aureus, Pseudomonas aeruginosa

FUNGICIDAL* (Against pathogenic fungi)

VIRUCIDAL* (Against viruses such as Herpes simplex Type I, Influenza A₂ (Japan), Vaccinia, Adenovirus Type 2 and HIV-1 (AIDS virus))

PREVENTS MOLD & MILDEW*

*On Environmental Surfaces in the presence of 5% blood serum.

ACTIVE INGREDIENTS: n-Alkyl (60% _{C₁₄} , 30% _{C₁₆} , 5% _{C₁₂} , 5% _{C₁₈})	
Dimethyl Benzyl Ammonium Chlorides	7.45%
n-Alkyl (68% _{C₁₂} , 32% _{C₁₄}) Dimethyl Ethylbenzyl Ammonium Chlorides	7.45%
INERT INGREDIENTS	85.10%
TOTAL	100.00%


KEEP OUT OF REACH OF CHILDREN

DANGER: Corrosive. Harmful if swallowed. Causes eye and skin injury. Do not handle pouches with wet hands. See Side Panel for Precautionary Statements.

NET CONTENTS 75 Water-Soluble 1/2 oz. Packets

NATIONAL LABORATORIES

**Lehn & Fink Industrial Products
Division of Sterling Drug Inc.
Montvale, NJ 07645**



Not to be used. Registrant claims to be in accordance with
EPA's Phase 2 Guidance for Phase 2
Active to Inert Change in Status.

Panel 2

20/11

Premeasured QUATYSL® 256, in 1/2 oz. water soluble film packets, provides a controlled, convenient and economical method for effective CLEANING, DISINFECTION AND DEODORIZATION.

Formulated for use in Hospitals, (including Housekeeping Services, Nursing Services, Operating Rooms, Physical Therapy Departments, Autopsy Facilities, Dietary) Nursing Homes and all other institutions requiring environmental surface disinfection.

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

DANGER: Corrosive. Harmful if swallowed. Causes eye and skin injury. Do not get powder in eyes, on skin or in clothing. Do not open pouches directly. Avoid contamination of food. Do not remove pouches from package except for immediate use. Wash hands thoroughly after using. Remove and wash contaminated clothing.

STATEMENT OF PRACTICAL TREATMENT


In case of eye contact, IMMEDIATELY flush eyes thoroughly with water, remove any contact lenses, and continue to flush eyes with plenty of water for at least 15 minutes. Call a physician.
In case of skin contact, immediately flush with soap and water. If irritation persists, consult a physician.
If swallowed, drink promptly a large quantity of water to dilute the product. Get immediate medical attention.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in original container in areas inaccessible to all children. Keep securely closed.
DISPOSAL: Do not reuse empty container. Rinse thoroughly with water and detergent. Wrap and discard in trash.



Panel 3

**PREMEASURED QUATSYL® 256
Detergent-Disinfectant-Deodorizer**

FOR HOSPITAL AND INSTITUTIONAL USE 1/2 OZ. WATER SOLUBLE PACKET

DIRECTIONS FOR USE

Always use according to directions. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Prepare a fresh solution daily. Use warm water. Change solution frequently if grossly soiled or diluted. Always remove filth and heavy soil prior to use.

**FOR ONE STEP CLEANING AND DISINFECTION
ONE PACKET PER 1 GALLON WATER**

Apply to environmental surfaces, such as floors, walls, furniture and fixtures, and similar hard surfaces with a mop, cloth, sponge or mechanical sprayer. Maintain thoroughly wet surface for 10 minutes.


**FOR MOLD AND MILDEW CONTROL
ONE PACKET PER 1 GALLON WATER**

Clean surface first, then apply via mop or sprayer until surface is moistened thoroughly. Let air dry. Repeat procedure and reapply to surface on a weekly basis or as necessary.

**KILLS HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) (AIDS VIRUS) ON
PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY
FLUIDS.**

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OF
SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eye coverings.



Panel 3 (continued)

4, 4

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of Pressurized Quatsyl 256.

CONTACT TIME: Allow surface to remain wet for 10 minutes.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

