

Precautionary Statements
Hazards to Humans and Domestic Animals
Caution

Harmful if swallowed
May cause skin irritation
Avoid contact with skin, eyes or clothing
If contact with eyes occurs, flush with plenty of water

Environmental Hazards

Do not contaminate irrigation ditches or water used for domestic purposes. Flush sprayer out on suitable non-crop area after use. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Trace amounts of spray drift may cause severe injury to both growing and dormant plants. Course sprays are less likely to drift. Spray only on calm days using low pressure and lowered booms. Vapors from this product may injure 2,4-D sensitive plants in the vicinity.

Net Contents Gallons/ Liters



Diamond Shamrock

A 7
D

677-105

Crop Rider®

2,6,7D

Weed Killer

Active Ingredient: Butyl Ester of 2,4-Dichlorophenoxyacetic Acid* 36.8%
Inert Ingredients: 63.2%
Total 100.0%

*2,4-Dichlorophenoxyacetic Acid Equivalent: 29.35% (2.4 lbs/gal) Isomer specific by AOAC Method No. 6 D01-5

Keep Out Of Reach Of Children

CAUTION

See side panel for additional precautionary statements

Diamond Shamrock Corporation
Agricultural Chemicals Division
Cleveland, Ohio 44114

AG-05980

EPA Reg No 677-105-AA

001 21

01003

Crop Rider 2,67D

Read entire label before using this product

General Information

Crop Rider 2,67D Weed Killer is recommended for selective control of weeds such as bindweed, dandelion, dock, goldenrod, lambsquarters, mustard, peppergrass, pigweed, ragweed, shepherdspurse, thistles, wild carrot, wild morningglory, wild radish and many other broadleaved weeds on crops.

Treating Small Areas: One tablespoonful of Crop Rider 2,67D in 100 gallons of water is about equal to 1 quart in 100 gallons.

When stored at temperatures below freezing, it may be necessary to warm up to at least 45°F and shake thoroughly before using.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not use the same spray equipment for applying other materials to plants as injury will result. Use care to avoid spray drift to 2,4-D susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals.

Preparation of the Spray: Fill the spray tank with half the required amount of water, then add the recommended amount of Crop Rider 2,67D with agitation and continue filling the spray tank. The amount of water required for low-volume applications may vary from 5 to 25 gallons per acre. For high-volume applications, 100 gallons or more of water will be needed for good coverage. In any case, use the same amount of 2,4-D recommended per acre.

Time of Application: Best results are obtained when Crop Rider 2,67D is used on weeds that are young and in a rapidly-growing condition. Applications of lower rates to susceptible annual weeds usually will be satisfactory, but for perennial weeds and other conditions where kill is difficult, use higher rates. When used as a selective spray on crops, the stage of growth of the crop must be considered. Some woody plants and weeds are hard to kill and repeat applications may be necessary.

Small Grain Crops (Wheat, Barley, Rye): Apply Crop Rider 2,67D at the rate of 1 to 1½ pints per acre in sufficient water for uniform coverage on small grains when fully tillered or stooled (4 to 8 inches tall) but before head emerges from the boot. **Do not use** on grain undersown with legumes, such as alfalfa or clovers, except where some legume injury can be tolerated. **Do not apply** when grain is headed.

Do not forage or graze treated grain fields within two weeks after treatment with 2,4-D.

Corn: Preemergence - Use Crop Rider 2,67D at the rate of 1½ to 3 quarts in sufficient water for uniform coverage. Best results are obtained when applied 3 to 5 days after planting, but before corn emerges. **Do not apply** on light, sandy soils. **Postemergence -** Apply at the rate of ¼ to 1 pint per acre. When spraying corn above 10 inches in height, use nozzle extensions (corn drops) directing the spray down to the base of the corn plant. **Do not apply** from tasseling to dough stage. Avoid cultivation soon after treatment when corn may be brittle. Hybrids vary in tolerance to 2,4-D. Consult local agricultural experiment station or extension service weed specialists regarding the use of 2,4-D on your specific hybrid.

Sorghum: Use Crop Rider 2,67D at the rate of 1 to 1½ pints per acre on plants 6 to 12 inches high and when second roots are well established. Use drop nozzles when crop is over 10 inches tall. **Do not apply** from flowering to dough stage.

Pastures: To control many broadleaved weeds in pastures, meadows and rangelands, use 3 pints of Crop Rider 2,67D in sufficient water to provide for uniform application. Treat pastures when weeds are growing actively. **Do not apply** to recently-seeded pastures until grass is well established. Most legumes are usually injured or killed at the rates recommended. For control of wild garlic and onion in pastures, apply 3 to 4½ quarts per acre in very early spring and repeat annually until these weeds are eradicated. Injury may result if applied to St. Augustinegrass and bentgrass.

Do not graze treated areas by dairy animals within seven days after treatment.

Woody Plant Control: To control 2,4-D susceptible woody plants such as alder, buckbrush, elderberry, sumac and willow in non-cropped or other waste areas, use 5 to 8 pints in 100 gallons of water. Wet thoroughly all parts of the plants, including foliage and stems, to the point of run-off.

General Weed Control: Along fence rows, ditchbanks, around farm buildings and other waste areas, use 3 to 4 pints of Crop Rider 2,67D in 100 gallons of water per acre. Thoroughly wet all foliage to run-off.

Storage and Disposal

Storage - Do not contaminate water, food or feed by storage or disposal. **Do not store** near fertilizers, seeds, insecticides or fungicides.

Disposal - Do not reuse empty container. Pesticide spray mixture or rinse that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Triple rinse (or equivalent) container and offer for disposal in approved landfill or bury in a safe place. Open dumping is prohibited.

Consult federal, state or local disposal authorities for approved alternative procedures.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and **Buyer** assumes the risk of any use contrary to such directions. **Seller** makes no other express or implied warranty including any other express or implied warranty of **Fitness** or of **Merchantability**, and no agent of **Seller** is authorized to do so except in writing with a specific reference to this warranty. In no event shall **Seller's** liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

66