

NET CONTENTS

GALLONS

DIRECTIONS FOR USE

TIME OF APPLICATION: Best results are obtained when CROP RIDER LV-4D is used on weeds that are young and in a rapid growing condition. Applications of lower rates to susceptible annual weeds usually will be satisfactory, but for perennial weeds and other conditions where kill is difficult, use higher rates. When used as a selective spray in crops, the stage of growth of the crop must be considered. Some woody plants and weeds are hard to kill and repeat applications may be necessary.

SMALL GRAIN CROPS (Wheat, Oats, Barley, Rye): Apply CROP RIDER LV-4D at the rate of $\frac{1}{2}$ to 1 pint per acre, in sufficient water for uniform coverage, on small grains when fully tillered or stooled 4 to 8 inches tall but before head emerges from the "boot". Do not use on grains undersown with legumes such as alfalfa or clovers except where some legume injury can be tolerated. Do not apply when grain is headed.

DO NOT forage or graze treated grain fields within two weeks after treatment with 2,4-D.

DO NOT feed treated straw to livestock.

CORN: PRE-EMERGENCE — use CROP RIDER LV-4D at the rate of 1 to 2 quarts per acre in sufficient water for uniform coverage. Best results are obtained when applied 3 to 5 days after planting but before corn emerges. **DO NOT** apply to light sandy soils. **POST-EMERGENCE** — apply at the rate of $\frac{1}{2}$ pint per acre. When spraying corn above 12 inches in height, use nozzle extensions ("corn drops"), directing the spray down to the base of the corn plant.

SORGHUM: Apply CROP RIDER LV-4D at the rate of $\frac{1}{2}$ pint per acre on plants 6 to 12 inches high when secondary roots are well established.

PASTURES: To control many broadleaved weeds in pastures, meadows and rangelands, use 2 pints of CROP RIDER LV-4D in sufficient water to provide for uniform application. Treat pastures when weeds are growing actively. Do not apply to recently seeded pastures until grass is well established. Most legumes are usually injured or killed at the rates recommended. For control of wild garlic and onion in pastures, apply 2 to 3 quarts per acre in very early spring and repeat annually until these weeds are eradicated. Do not forage or graze treated areas by dairy animals within seven days after treatment.


LAWNS, GOLF COURSES AND SIMILAR TURF: Apply 2 pints per acre of CROP RIDER LV-4D in sufficient water to provide uniform coverage. Do not apply to newly seeded lawns until grasses become well established, or on creeping grasses such as bent. For wild garlic and wild onion, apply 2 to 3 quarts per acre in very early spring each year for at least 3 years.

FOR TREATING SMALL AREAS: One tablespoonful of CROP RIDER LV-4D in $1\frac{1}{2}$ gallons of water is about equal to 1 quart in 100 gallons of water.

WOODY PLANT CONTROL: To control 2,4-D-susceptible woody plants such as Willow, Alder, Buckbrush, Sumac, and Elderberry in non-cropped or other waste areas, use 2 to 3 quarts in 100 gallons of water. Wet thoroughly all parts of the plants, including foliage and stems, to point of run-off.

GENERAL WEED CONTROL: On vacant lots, roadsides, and along fence rows use 1 to 2 quarts of CROP RIDER LV-4D in 100 gallons of water per acre. Thoroughly wet all foliage to run-off.

U.S.A. REG. NO. 677-110



®

A DIAMOND CHEMICAL

Weed Killer

(2,4-D Low-Volatile Emulsifiable Concentrate)


CROP RIDER® LV-4D

Contains 4.0 Pounds of 2,4-D Acid Equivalent Per Gallon

ACTIVE INGREDIENT:

Isooctyl Ester of 2,4-Dichlorophenoxyacetic Acid*	69%
INERT INGREDIENTS:	31%
TOTAL	100%

*Equivalent to 2,4-Dichlorophenoxyacetic Acid 45.8%



GENERAL INFORMATION

CROP RIDER LV-4D WEED KILLER is recommended for controlling Arrowhead, Dandelion, Docks, Bindweed, Henbit, Lamb's-quarter, Mallow, Mustard, Peppergrass, Pigweed, Plantain, Ragweed, Shepherd's-purse, Thistles, Wild Carrot, Wild Morningglory, Wild Garlic, Wild Onion, and many other broadleaved weeds.

PREPARATION OF THE SPRAY: Fill the spray tank with half the required amount of water, then add the recommended amount of CROP RIDER LV-4D with agitation and continue filling the spray tank with water. Use enough water per acre to give uniform coverage. The amount of water required for low volume applications may vary from 5 to 25 gallons per acre. For high volume applications 100 gallons, or more, of water will be needed for good coverage. In any case, use the same amount of 2,4-D recommended per acre.

CAUTION: Avoid spray drift to susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees, and ornamentals. Spray drift may cause severe injury to both growing and dormant plants. Coarse sprays are less likely to drift. Although this ester is much less volatile than butyl or isopropyl esters, high temperatures may produce vapors after application which might injure 2,4-D-sensitive plants growing nearby.

Contamination of shallow fish bearing waters at application rates recommended on this label may kill fish. **DO NOT** contaminate irrigation ditches or water used for domestic purposes.

DO NOT store near fertilizers, seeds, insecticides, or fungicides.

DO NOT take internally. Avoid contact with skin, eyes, or clothing.

DO NOT use the same spray equipment for applying other materials to plants, as injury will result.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and **BUYER** assumes the risk of any use contrary to such directions. **SELLER** makes no other express or implied warranty, including any other express or implied warranty of **FITNESS** or of **MERCHANTABILITY**, and no agent of **SELLER** is authorized to do so except in writing with a specific reference to this warranty. In no event shall **SELLER's** liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

CAUTION KEEP OUT OF REACH OF CHILDREN.
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

DIAMOND SHAMROCK CORPORATION - CLEVELAND, OHIO 44115

A-31A-L-368