

677-130

AUG 1 1 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROUENWEEDS ACT
FOR ECONOMIC POISON REGISTRATION
EPA UNCLAS NO. 677-130-1

CONTENTS

GALLONS

READ ENTIRE LABEL
BEFORE USING THIS PRODUCT

GENERAL INFORMATION

CROP RIDER AMINE 6D WEED KILLER is recommended for controlling arrowweed, dandelion, docks, henbit, garden rad, lambsquarters, mustards, peppergrass, pigweed, plantains, ragweed, shepherdspurse, thistles, wild carrot, wild morning glory, wild radish and many other 2,4-D susceptible broad-leaved weeds in crops. This product also can be used on lawns, golf courses, parks, along highways, vacant lots and other similar areas without injury to most grasses.

TREATING SMALL AREAS: One tablespoonful of this product in 1 1/2 gallons of water is about equal to one quart in 100 gallons.

When stored at temperatures below freezing, it may be necessary to warm up to at least 45° F. and shake thoroughly before using.

DIRECTIONS FOR USE

PREPARATION OF THE SPRAY: Fill the spray tank with half the required amount of water, then add the recommended amount of CROP RIDER AMINE 6D WEED KILLER with agitation and continue filling the spray tank with water. Use enough water per acre to give uniform coverage. The amount of water required for low-volume applications may vary from 5 to 25 gallons per acre. For high-volume applications, 100 gallons or more of water will be needed for good coverage. In any case, use the same amount of 2,4-D recommended per acre.

TIME OF APPLICATION: Best results are obtained when CROP RIDER AMINE 6D is used on weeds that are young and in a rapidly growing condition. Applications at lower rates to susceptible annual weeds usually will be satisfactory, but for perennial weeds and other conditions where weed kill is difficult, use higher rates. When used as a selective spray in crops, the stage of growth of the crop must be considered.

SMALL GRAIN CROPS: Wheat, Barley, Rye. Apply CROP RIDER AMINE 6D at the rate of 1.3 to 1 pint per acre on small grains when fully tillered or headed, 4 to 8 inches tall, but before head emerges from the boot. Injury will be likely to occur if this product is applied earlier than tiller or later than boot. DO NOT use on grains undersown with legumes such as alfalfa or clovers, except where some legume injury can be tolerated. DO NOT apply when grain is headed.

DO NOT forage or graze treated grain fields within two weeks after treatment with 2,4-D.


AERIAL APPLICATION: Use CROP RIDER AMINE 6D at the rate of 1.3 to 2 pints in 1 to 10 gallons of water per acre.

FLAX: Apply CROP RIDER AMINE 6D at the rate of 1.3 pint per acre as weed growth warrants, including the flax plants have formed 4 to 5 leaves, 2 to 6 inches tall. DO NOT treat flax after the early bud stage.

RICE: In the South and Southwest, apply from 1 to 1.3 pints per acre in 5 to 10 gallons of water 7 to 10 weeks after planting or when the rice is fully tillered, but not yet in the boot. California: Apply 1 to 1.3 pints per acre 55 to 65 days after planting, when the rice is through the water and well established. For difficult weeds and where daily temperatures do not exceed 95° F., 2 pints per acre may be used.

CORN, PREEMERGENCE: Use CROP RIDER AMINE 6D at the rate of 2.3 to 1.3 pints per acre in sufficient water for uniform coverage. Best results are obtained when applied 3 to 5 days after planting, but before corn emerges. DO NOT apply to light, sandy soils. **POSTEMERGENCE:** Apply at the rate of 1 pint per acre. When spraying corn above 10 inches in height, use nozzle extensions, corn drops, directing the spray down to the base of the corn plant. DO NOT apply

EPA REG. NO. 677-130-AA



®
**Diamond Shamrock
Chemical Company**

Weed Killer

CROP RIDER® AMINE 6D

Contains 6.0 Pounds 2,4-D Acid Equivalent Per Gallon

ACTIVE INGREDIENT

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*

70%

INERT INGREDIENTS

30%

TOTAL

100%

*Equivalent to 2,4-Dichlorophenoxyacetic Acid 58.0%

**CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.**

AGRICULTURAL CHEMICALS DIVISION CLEVELAND, OHIO 44115

DIAMOND SHAMROCK CORPORATION

AQ-124 A

3644-A-19C-1-571

from tasseling to dough stage. Hybrids vary in tolerance to 2,4-D. Consult local agricultural experiment station or extension service weed specialists regarding the use of 2,4-D on your specific hybrid.

SORGHUM: Use spray at the rate of 1.3 pint per acre on plants 6 to 12 inches high when secondary roots are well established. Use drop nozzles when crop is over 10 inches tall. DO NOT apply from flowering to dough stage.

SUGAR CANE: Apply at the rate of 2.3 to 1.3 pints per acre in the spring after cane emerges and through layby.

PASTURES: To control many broadleaved weeds in pastures, meadows and rangelands, use 2.3 to 1 pint per acre of CROP RIDER AMINE 6D in sufficient water to provide for uniform application. Treat pastures when weeds are growing actively. DO NOT apply when grass is in boot to neck stage. DO NOT apply to recently-seeded pastures until grass is well established. Most legumes are usually injured or killed at the rates recommended. DO NOT graze dairy animals in treated areas within 7 days after application.

LAWNS AND GOLF FAIRWAYS: Apply 2.3 to 1 pint of CROP RIDER AMINE 6D per acre in sufficient water to obtain uniform coverage. DO NOT apply to bentgrasses or to freshly-seeded turf until grasses have become well established. Injury may result if applied to Dichondra, St. Augustinegrass, carpetgrass and clovers.

GENERAL WEED CONTROL: On vacant lots, roadsides and along fence rows, use 2.3 to 1 quart of CROP RIDER AMINE 6D in 100 gallons of water per acre. Thoroughly wet all foliage to run-off.

CAUTION

Harmful if swallowed or absorbed through the skin.

CAUSES IRRITATION OF EYES AND SKIN

DO NOT get in eyes. In case of contact, flush eyes with plenty of water for at least 15 minutes and get medical attention.

DO NOT allow contact with skin or clothing. Wash exposed skin gently with plenty of soap and water.

DO NOT store near fertilizers, seeds, insecticides or fungicides.

DO NOT contaminate irrigation ditches or water used for domestic purposes.

Use care to avoid spray drift to 2,4-D susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals. Trace amounts of spray drift may cause severe injury to both growing and dormant plants. Coarse sprays are less likely to drift. Spray only on calm days using low pressure and lowered boom.

Flush sprayer out on suitable non-crop area after use. DO NOT use the same spray equipment for applying other materials to plants as injury will result.

DO NOT reuse empty container. Return to drum reconditioner, or destroy by perforating, crushing and burying or discarding in a safe place.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and BUYER assumes the risk of any use contrary to such directions. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of SELLER is authorized to do so except in writing with a specific reference to this warranty. In no event shall SELLER's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.