



Diamond Shamrock

Daconil 2787 Fungicide

Turf Care (75% Wettable Powder)

Precautionary Statements

Hazards To Humans And Domestic Animals

Danger

- Corrosive. Causes Severe Eye Damage.
- Do not get in eyes. wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.
- Avoid contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using.
- Do not take internally.
- Avoid breathing dust or spray mist.
- Note: Daconil 2787 Fungicide may produce temporary allergic effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Affected persons respond to treatment with antihistamines or steroid creams and or systemic steroids.

Environmental Hazards

- This product is toxic to fish. Keep out of lakes, streams or ponds.
- Do not apply where runoff is likely to occur.
- Do not apply when weather conditions favor drift from areas treated.
- Do not graze treated areas. Do not feed clippings to livestock.
- Apply only to areas specified on label.

Active ingredient:	Chlorothalonil (tetrachloroisophthalonitrile)*	75.0%
Inert ingredients:		25.0%
Total		100.0%

*Covered under U.S. Pat. No. 3,290,353 and No. 3,331,735

DANGER: Keep Out Of Reach Of Children
See side panel for additional precautionary statements

(75% Wettable Powder)

A Broad-Spectrum Fungicide for Use on Golf Course Tees, Greens and Fairways; Ornamental Turfgrass and Ornamentals.
Read Entire Label and Use Only as Directed

General Information

Daconil 2787 Fungicide is highly effective and widely recommended for the control of a number of important golf course turf and ornamental lawn diseases as well as a number of commonly occurring diseases of ornamentals. Daconil 2787 Fungicide is specifically formulated for use on greens, tees and fairways, ornamental turfgrasses as well as on a number of ornamentals.


Do not combine Daconil 2787 Fungicide in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible and noninjurious under your conditions of use.

The required amount of Daconil 2787 Fungicide should be added slowly into the spray tank during filling with agitator running. Maintain agitation during spray operations.

Directions For Use

ONLY FOR SALE TO, USE, AND STORAGE BY LANNCARE AND LANDSCAPING PERSONNEL OR COMMERCIAL APPLICATORS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Turf

Do not mow or water after treatment until greens or turfgrasses are thoroughly dry. A 24 hour waiting period following treatment is preferred. Daconil 2787 Fungicide should always be used in conjunction with good turf management practices.

Golf Course Fairways: Apply 5 to 10 pounds of Daconil 2787 Fungicide per acre in 30 to 40 gallons of water. Under severe disease conditions, use higher indicated rate and lower interval between applications.

Helminthosporium leaf spot: Apply Daconil 2787 Fungicide on a 14 to 21 day interval, beginning after first or second cutting and continuing as long as conditions favor disease.

Dollar spot: Apply initial application when conditions favor disease development and repeat at 14 to 21 day intervals as needed.

Large brown patch: Make initial application when disease threatens and repeat at 7 to 14 day intervals as needed.

Golf Course Tees and Greens and Ornamental Turfgrass: Apply Daconil 2787 Fungicide in an adequate amount of water to provide complete coverage. This amount may vary from 5 to 15 gallons per 1,000 square feet. See below for suggested rates and timing. Under severe disease conditions, use the curative rates and spray on a 7 day schedule.

Disease	Interval Of Application	Rate Ounces per 1,000 sq. ft.	
		Preventive*	Curative**
Copper spot	7 - 10 day	4 - 6	6 - 8
Curvularia leaf spot	7 - 10	2 - 4	4 - 8
Dollar Spot	7 - 14	2 - 4	4 - 8
Gray leaf spot	7 - 10	2 - 4	4 - 8
Helminthosporium leaf spot and melting out	7 - 10	2 - 4	4 - 8
Large brown patch	7 - 10	2 - 4	4 - 8
Red thread	7 - 10	2 - 6	6 - 8
Stem rust of bluegrass	7 - 14	4 - 6	6 - 8
Dichondra Allertonia leaf spot (California only)	7 - 14	4 - 6	6 - 8

* Recommended rates for preventing disease establishment; use lower rates when disease conditions are light to moderate, higher indicated rates when conditions are severe.

** Rates for use on a 7 day schedule when disease is present. Higher indicated rate should be applied under severe conditions.

BEST DOCUMENT AVAILABLE.

180

Ornamentals

Daconil 2787 Fungicide is recommended for the control of a number of diseases of ornamentals. Apply at first sign of disease and repeat at 7 to 14 day intervals throughout the season. Make applications on a 7 day schedule during periods of wet weather or severe disease conditions. Rates provided are in pounds of **Daconil 2787 Fungicide** per 100 gallons of water. (One level spoonful per gallon approximates a rate of one pound of **Daconil 2787 Fungicide** per 100 gallons of water as a dilute spray.) Application should be thorough to obtain adequate coverage of the foliage.

Chrysanthemums: Ascochyta ray blight and gray mold—Use 1½ pounds per 100 gallons of water.

Impatiens: Botrytis blight—Use 1½ pounds per 100 gallons of water.

Botrytis blight and leaf spot—Use 1½ pounds per 100 gallons of water.

Black spot and blossom blight—Use 1 pound per 100 gallons of water.

Pink Petunias: Powdery mildew—Use 1½ pounds per 100 gallons of water.

Large plants (Florida only): Weekly applications should be made at rate of 1½ pounds **Daconil 2787 Fungicide** per 100 gallons of water beginning when conditions are favorable for disease on the following crops:

Herbaceous Fern—for control of Ascochyta blight, Cercospora spot, Cylindrocladium leaf spot and Rhizoctonia blight.

Podocarpus—for control of Phytophthora blight and Dactylaria spot.

Water Plant (Rhoeo discolor)—for control of tan leaf spot.

Do not spray plants during extremely hot and sunny conditions or when conditions will not promote drying. Do not spray wilted plants.

Storage And Disposal

Storage—Do not contaminate water, food or feed by storage or disposal.

Store in a dry place.

Disposal—Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of containers in an incinerator or landfill approved for pesticide containers or bury in a safe place. Open dumping is prohibited.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

EPA Est. No. 35982-TX-1



Diamond Shamrock

Diamond Shamrock Corporation
Agricultural Chemicals Division
Cleveland, Ohio 44114

EPA Reg. No. 677-229-AA

BEST DOCUMENT AVAILABLE

181