

NET CONTENTS


GALLONS

READ ENTIRE LABEL BEFORE USING THIS PRODUCT

GENERAL INFORMATION

TRAILWAY 2D/2T is a selective herbicide... It is effective against... weeds and brush... It is used on... roadsides, railroads, and other areas... It is applied as a spray...

EPA REG. NO. 677-238-AA



RECORDED
10-22-71
677-238

BEST DOCUMENT AVAILABLE

Diamond Shamrock Chemical Company

Weed and Brush Killer TRAILWAY 2D/2T

Contains 2 Pounds Each of 2,4-D and 2,4,5-T Acid Equivalent Per Gallon

ACTIVE INGREDIENTS	
N-Oleyl 1,3-propylene diamine salt of 2,4-Dichlorophenoxyacetic Acid*	29.0%
N-Oleyl 1,3-propylene diamine salt of 2,4,5-Trichlorophenoxyacetic Acid**	27.0%
INERT INGREDIENTS	44.0%
TOTAL	100.0%

*Equivalent to 2,4-Dichlorophenoxyacetic Acid 20.9%
**Equivalent to 2,4,5-Trichlorophenoxyacetic Acid 20.9%

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

AGRICULTURAL CHEMICALS DIVISION-CLEVELAND, OHIO 44115

DIAMOND SHAMROCK CORPORATION

AG-137A

3219 A 1 MC 1 871

USE IN ACCORDANCE WITH THE LABEL... TRAILWAY 2D/2T is a selective herbicide... It is effective against... weeds and brush... It is used on... roadsides, railroads, and other areas... It is applied as a spray...

GENERAL WEED CONTROL... TRAILWAY 2D/2T is a selective herbicide... It is effective against... weeds and brush... It is used on... roadsides, railroads, and other areas... It is applied as a spray...

Most regimens are usually cured... TRAILWAY 2D/2T is a selective herbicide... It is effective against... weeds and brush... It is used on... roadsides, railroads, and other areas... It is applied as a spray...

CAUTION

Harmful if swallowed or absorbed through the skin

CAUSES IRRITATION OF EYES AND SKIN

DO NOT get in eyes. In case of contact, flush eyes with plenty of water for at least 15 minutes and get medical attention.

DO NOT allow contact with skin or clothing. Wash exposed skin with plenty of soap and water.

DO NOT store near fertilizers, seeds, insecticides or fungicides.

Rinse out sprayer and container after use and properly dispose of wastes in non-crop areas away from other supplies.

DO NOT use the same spray equipment for applying other materials to plants as injury will result.

DO NOT contaminate water used for irrigation or domestic purposes.

Avoid spray drift to 2,4-D and 2,4,5-T susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals. Coarse sprays are less likely to drift. Spray only on still days with low pressure, coarse spray and lowered booms.

DO NOT graze dairy animals on treated areas within 6 weeks after application. DO NOT graze meat animals on treated areas within 2 weeks of slaughter.

DO NOT reuse empty container. Return to drum recycler, tiner or destroy by perforating, crushing and burying or discarding in a safe place.

This product is toxic to fish. Keep out of lakes, streams or ponds.

Learned under U.S. Patent No. 2,900,411

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to the chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and BUYER assumes the risk of any use contrary to such directions. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY and no agent of SELLER is authorized to do so except in writing with a specific reference to this warranty. In no event shall SELLER's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

52
57

**READ ENTIRE LABEL
BEFORE USING THIS PRODUCT**

GENERAL INFORMATION

TRAILWAY 2D/2T is a water-emulsifiable diamine formulation of 2,4-D and 2,4,5-T. It combines the advantages of both the ester and amine forms of 2,4-D and 2,4,5-T. TRAILWAY 2D/2T contains both the non-volatile features of amine and the effectiveness of esters. It is designed for control brush and woody plants along highways, railroad and utility rights-of-way, rangelands, fence or hedgerows and around buildings, tank farms, transformer stations and similar non-crop or waste areas where it is desired to kill or repress the growth of woody plants. TRAILWAY 2D/2T, besides being non-volatile, is especially effective against deep-rooted perennial weeds and woody plants because it is slow acting, thus allowing more translocation of the herbicide into the roots of the plant to afford a more complete kill. The following plants are included among those susceptible to TRAILWAY 2D/2T:

<u>Alder</u>	<u>Dogwood*</u>	<u>Persimmon</u>
<u>Am. crabapple</u>	<u>Elderberry</u>	<u>Pine*</u>
<u>Ash*</u>	<u>Elm</u>	<u>Poison ivy</u>
<u>Aspen</u>	<u>Greenbrier*</u>	<u>Poison oak</u>
<u>Barberry</u>	<u>Hackberry</u>	<u>Poplar</u>
<u>Bindweed</u>	<u>Hawthorn</u>	<u>Raspberry</u>
<u>Birch*</u>	<u>Hickory*</u>	<u>Rose</u>
<u>Blackberry</u>	<u>Honeylocust</u>	<u>Sagebrush</u>
<u>Blackoak</u>	<u>Honeylocust</u>	<u>Saltcedar*</u>
<u>Black locust</u>	<u>Honeylocust</u>	<u>Sassafras</u>
<u>Bowlder</u>	<u>Honeylocust</u>	<u>Servicberry</u>
<u>Brambles</u>	<u>Hornbeam</u>	<u>Silverberry</u>
<u>Buckbrush</u>	<u>Horse chestnut</u>	<u>Skunkbrush</u>
<u>Canada thistle</u>	<u>Juneberry</u>	<u>Shoebush</u>
<u>Catalpa</u>	<u>Knapweed</u>	<u>Spicebrush</u>
<u>Chestnut</u>	<u>Maple*</u>	<u>Sunox</u>
<u>Chinaberry</u>	<u>Mesquite</u>	<u>Sweetfern</u>
<u>Chokecherry</u>	<u>Mulberry</u>	<u>Sweetgum</u>
<u>Cocklebur</u>	<u>Mustard</u>	<u>Wild cherry</u>
<u>Cottonwood</u>	<u>Oak*</u>	<u>Willow</u>
<u>Currant</u>	<u>Orangeorange</u>	

For best results on these plants, use only the rates given in the directions under ideal control conditions. Lower rates may be found desirable on the species without an asterisk ().

DO NOT use around the home, recreation area or similar sites.

DIRECTIONS FOR USE

PREPARATION OF THE SPRAY: Fill the spray tank with half the required amount of water. Then add the recommended amount of TRAILWAY 2D/2T with agitation and continue filling the spray tank with balance of water. Keep agitator running when filling spray tank and during spray operations. Use enough water per acre to give uniform coverage.

TIME OF APPLICATION: Best results are obtained when TRAILWAY 2D/2T is used on weeds and brush during the early part of the growing season when the plants are in an actively-growing condition. DO NOT treat during periods of severe drought or in early fall when leaves have lost their healthy green color.

FOLIAGE TREATMENT: Foliage sprays are used during the growing season on dense brush or where woody vine growth, such as Smilax and greenbrier, is a problem. Apply TRAILWAY 2D/2T after foliage is well developed, using a drenching spray containing 1 gallon in 100 gallons of water (1/3 cupful in 2 gallons). For treating small areas, use 2 to 3 ounces of TRAILWAY 2D/2T in one gallon of water. Coverage should be complete, wetting all parts of the plants, including leaves and stems, to point of run-off. In controlling low-growing, undesirable woody plants in the hedgerow, a directed spray should be used to avoid damaging the taller, desirable hedgerow plants. High volumes are necessary where



**Weed and
TRAILWAY**

Contains 2 Pounds Each of 2,4-D and

ACTIVE INGREDIENTS:
N-Oleyl-1,3-propylenediamine salt
phenoxyacetic Acid*
N-Oleyl-1,3-propylenediamine salt
phenoxyacetic Acid**
INERT INGREDIENTS:

TOTAL
*Equivalent to 2,4-Dichloro
**Equivalent to 2,4,5-Trichloro

KEEP OUT OF REACH OF CHILDREN