


Agricultural Chemicals Division  
Diamond Shamrock Corporation  
1100 Superior Avenue  
Cleveland, Ohio 44114  
TE 691 5000

## Bravo W-75



**Diamond Shamrock**

### Supplemental Labeling



SUPPLEMENTAL DIRECTIONS FOR THE USE OF

BRAVO W-75  
(EPA Reg. No. 677-282)

FOR CONTROL OF DISEASES ON CONIFERS

#### Directions for Use on Tree and Orchard Crops

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply BRAVO W-75 in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Application through either ground equipment or aircraft is recommended except where otherwise indicated. When concentrate sprays are used the lower rate per acre of BRAVO W-75 listed may be used. The following spray volumes are recommended as gallons of spray per acre:

CROP	SPRAY VOLUME (Gallons per acre)	
	DILUTE	CONCENTRATE
CONIFERS:		
Forest stands	Not used	10 to 20 (aircraft)
Christmas trees	100	10 to 50 (aircraft or ground equipment)
Nursery beds	100	5 to 10 (ground equipment only)

Warranty and Limitation of Damages

Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of seller is authorized to do so except in writing with a specific reference to this warranty.

4/1/77

CROP	DISEASES	BRAVO W-75 RATE PER		APPLICATION DIRECTIONS
		ACRE	100 GAL *	
Conifers	Swiss needlecast	2 $\frac{3}{4}$ to 5 $\frac{1}{2}$ lbs	2 $\frac{3}{4}$ to 5 $\frac{1}{2}$ lbs	Single application technique: In Christmas tree plantations or forest stands make one application in the spring when new shoot growth is $\frac{1}{2}$ to 2 inches in length.
	Scleroderris canker (pines), Swiss needlecast and Rhabdocline needlecast (Douglas-fir)	1 $\frac{1}{2}$ to 2 $\frac{3}{4}$ lbs	1 $\frac{1}{2}$ to 2 $\frac{3}{4}$ lbs	Make the first application in spring when new shoot growth in $\frac{1}{2}$ to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.
	Sirococcus tip blight	2 to 3 $\frac{1}{2}$ lbs	2 to 3 $\frac{1}{2}$ lbs	
	Rhizosphaera needlecast (spruces), Scirrhia brown spot (pines)	5 $\frac{1}{2}$ lbs	5 $\frac{1}{2}$ lbs	
	Lophodermium needlecast	1 $\frac{1}{2}$ to 2 $\frac{3}{4}$ lbs	1 $\frac{1}{2}$ to 2 $\frac{3}{4}$ lbs	North Central and Northeastern States: Begin applications in mid-July to early August before infection occurs. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply 2 $\frac{3}{4}$ lbs per acre on a 3 week schedule.
	Botrytis seedling blight, Phoma twig blight	1 $\frac{1}{2}$ to 2 $\frac{3}{4}$ lbs	1 $\frac{1}{2}$ to 2 $\frac{3}{4}$ lbs	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.

\*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

NOTE: It is important to read and observe the precautionary statements and all other information appearing on the BRAVO W-75 label. Use only as directed.

NOTE: This labeling must be in the possession of the user at the time of pesticide application.