

Precautionary Statements
Hazards to Humans and Domestic Animals
Caution

Environmental Hazards

Butyl 4D

Butyl 4D

Butyl 4D

Butyl 4D

Warranty and Limitation of Damages

Butyl 4D

Butyl 4D

Butyl 4D

Butyl 4D

Butyl 4D

Butyl 4D

Butyl 4D

Net Contents 5 Gallons/18.9 Liters



Diamond Shamrock

Butyl 4D

Weed Killer

Active Ingredient

Inert Ingredients

Total

2,4-Dichlorophenoxyacetic Acid Equivalent 42.8% 3.8 lb/gal
(isomer specific by AOAC Method No. 6 D01-5)

Keep Out Of Reach Of Children

CAUTION

Diamond Shamrock Corporation
Agricultural Chemicals Division
Cleveland Ohio 44114
EPA Reg. No. 677-294-AA

76

78

Butyl 4D

Bulky 4D
Read entire label before using this product

General Information

Butyl 4D is a selective herbicide for use in corn, soybean, and sorghum. Many troublesome weeds, including the following, are controlled in farm buildings and storage areas. For full details, see a full list of weeds controlled by **Butyl 4D**:

[illegible]

Note: Local conditions, crop varieties and application of regulations vary and may differ significantly from those of local agricultural experiment stations. Consultations with local specialists and state regulatory agencies for the most current information are advised.

Treating Small Areas One tablespoon of Butyl 4D is added to 1 gallon of water. It is diluted to 1 quart in 100 gallons.

Note: when stored at temperatures below 45°F (7°C), it is necessary to warm contents to 45°F and equilibrate thoroughly before using.

Directions For Use

Directions For Use

[illegible][illegible][illegible]

Do not use the product on children under 12 years of age. Do not use the product on pregnant women or nursing mothers. Do not use the product on people with known hypersensitivity to any of the ingredients. Do not use the product on people with known hypersensitivity to any of the ingredients. Do not use the product on people with known hypersensitivity to any of the ingredients.

Corn Preemergence— Butyl 4D

Postemergence—Appl. Butyl 4D from emergence to 10 days after emergence. Spray 100 to 150 gal/acre that contains 1.5 to 2.5 lb of active ingredient.

Do not

1. 1990年12月15日，在北京市召开的“中国环境与发展”高层论坛上，江泽民总书记发表了重要讲话，指出：“中国是一个发展中国家，在现代化过程中，必须走出一条既发展经济，又保护环境的道路。”

$\text{C}_{10}\text{H}_8\text{N}_2\text{O}_2 + 2\text{H}_2\text{O} \rightarrow \text{C}_{10}\text{H}_{10}\text{N}_2\text{O} + \text{H}_2\text{O}_2$
 $\text{C}_{10}\text{H}_{10}\text{N}_2\text{O} + \text{H}_2\text{O}_2 \rightarrow \text{C}_{10}\text{H}_{12}\text{N}_2 + \text{H}_2\text{O}$
 $\text{C}_{10}\text{H}_{12}\text{N}_2 + \text{H}_2\text{O} \rightarrow \text{C}_{10}\text{H}_{14}\text{N}_2 + \text{H}_2$

[illegible]

Sorghum A-1, Butyl 4D

Amount of Butyl 4D per Acre

Crop (See Detailed Directions Above)	For Average Conditions	For Dry Conditions as in Western States*
Wheat, Barley, Rye	1 1/2 bushels	1 1/2 bushels
Alfalfa, Lucerne	1 1/2 bushels	1 1/2 bushels
Perennial Weeds Control	1 bushel	1 1/2 bushels
Preemergence	1 1/2 bushels	1 1/2 bushels
Postemergence	1 bushel	1 1/2 bushels
Grasshopper	1 bushel	1 1/2 bushels
For Stronger	1 bushel	1 1/2 bushels

[illegible][illegible][illegible]

General Weed Control A 2,4-D, 2,4,5-T, and 2,4,6-T trisodium salt formulation. Controls broadleaf weeds in corn, soybeans, sorghum, and cotton. **Bulky 4D** controls broadleaf weeds in corn, soybeans, sorghum, and cotton.

Storage and Disposal

Storage—Do not store the water in the container for more than 24 hours.

[illegible][illegible]

EPA Est No 5905 1A-1

75-70-65

The diagram shows a 4x4 grid of points. The points are arranged in a regular pattern, with some points highlighted in black and others in white. The grid is labeled with 'a' and 'b' at the bottom, indicating the spacing between points.