

ACCEPTED

2-172

NET CONTENTS

GALLONS

EPA REG. NO. 677-297-AA

READ ENTIRE LABEL  
BEFORE USING THIS PRODUCT

GENERAL INFORMATION

AMINE 2D/2T is an amine salt mixture of 2,4-D and 2,4,5-T which forms a solution in water. It will not mix with diesel oil, kerosene or gasoline.

The following is a partial list of some weedy plants controlled by AMINE 2D/2T:

Aleppo	Bam	Sorrelberry
Amaranth	Carex	Poison ivy
Amaranth	Cattail	Poison oak
Barnyard	Hemp nettle	Spurred
Buckwheat	Lettuce	W. sycamore
Burdock	Mulch	Wild grape
Burdock	Oats	Willow
Bindweed	Overorange	

When stored at temperatures above freezing, it may be necessary to warm contents to 50° F. and mix thoroughly before using.

DIRECTIONS FOR USE

PREPARATION OF THE SPRAY: Fill the spray tank with half the required amount of water. Add the recommended amount of AMINE 2D/2T with agitation and continue filling the spray tank with balance of water.

FOGGER TREATMENT: For use in the control of weeds growing in areas where there is no ground cover.

AMINE 2D/2T may be applied to brush, shrubs, trees, vines, ground covers, etc., by hand spraying, knapsack sprayer, or by airless pressure sprayer. After application, AMINE 2D/2T will follow the water leading to the underlying soil, and may take 3 to 4 weeks to penetrate plant roots. Coverage should be complete, wetting all parts of the plant, including leaves and stems to the point of run-off. Applications are more effective when applied to actively growing plants. DO NOT treat during periods of severe drought or in early fall when leaves have lost their healthy green color. AMINE 2D/2T acts more slowly than ester spray. Therefore, a retreatment of that may be necessary should be delayed until the following season. Certain woody plants are less resistant to treatment with this product so be controlled by using 10 VOC or 10 VOC 2D/2T as a basal or stump spray.



®  
**Diamond Shamrock  
Chemical Company**

**Brush Killer**

**AMINE 2D/2T**

Contains 20 Pounds Each of 2,4-D and 2,4,5-T Acid Equivalent Per Gallon

**ACTIVE INGREDIENTS:**

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid\* 24.8%  
Triethylamine Salt of 2,4,5-Trichlorophenoxyacetic Acid\*\* 28.8%

**INERT INGREDIENTS:**

TOTAL 100.0%

\*Equivalent to 2,4-Dichlorophenoxyacetic Acid 20.6%

\*\*Equivalent to 2,4,5-Trichlorophenoxyacetic Acid 20.6%

**CAUTION: KEEP OUT OF REACH OF CHILDREN.  
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.**

**AGRICULTURAL CHEMICALS DIVISION-CLEVELAND, OHIO 44115**

AU 246

DIAMOND SHAMROCK CORPORATION

3752 A 146 L 571

Aerial application may be of use in control of brush and woody plants where there would be no danger of drift to susceptible crops. Applications should only be made by applicators experienced in the use of 2,4-D and 2,4,5-T formulations. Regulations governing aerial application of herbicides are in effect in many states. Consult local regulatory agencies concerning requirements before making applications.

**CAUTION**

Harmful if swallowed or absorbed through the skin.

Causes irritation of eyes and skin.

DO NOT get in eyes. In case of contact, flush eyes with plenty of water for at least 15 minutes and get medical attention.

DO NOT allow contact with skin or clothing. Wash exposed skin gently with plenty of soap and water.

DO NOT store near fertilizers, seeds, insecticides or fungicides.

DO NOT contaminate irrigation ditches or water used for domestic purposes.

Use care to avoid spray drift to 2,4-D and 2,4,5-T susceptible plants such as cotton, tobacco, flowers, grapes, fruit trees and ornamentals. Trace amounts of spray drift may cause severe injury to both growing and dormant plants. Coarse sprays are less likely to drift. Spray only on calm days using low pressure and lowered booms.

Flush sprayer out on suitable non-crop area after use. DO NOT use the same spray equipment for applying other materials to plants as injury will result.

DO NOT reuse empty container. Return to drum records, incinerate or destroy by puncturing, crushing and burying or discarding in a safe place.

DO NOT graze dairy animals on treated areas within 6 weeks after application. DO NOT graze meat animals on treated areas within 2 weeks of slaughter.

**WARRANTY AND LIMITATION OF DAMAGES**

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and BUYER assumes the risk of any use contrary to such directions. SELLER makes no express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY, and no agent of SELLER is authorized to do so except in writing with a specific reference to this warranty. In no event shall SELLER's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.