NET CONTENTS [redacted]

[redacted]

BRAVO® 6F

Agricultural Fungicide

[redacted]

DANGER: [redacted]

AGRICULTURAL [redacted] CLEVELAND, OHIO 44114

DIAMOND SHAMROCK CORPORATION

A Broad

REA FNTI

NOTE: Slowly invert container several times to

N (SNAP)

CUCUMBER

BBAGE, CAULIFLOWER BROCCOLI, BRUS
S SPROUTS

CANTALOUPE,
MELON

ROT

LERY

WATERMELON

SWEET

BRAVO® 6F

Agricultural Fungicide

Contains 6.0 Pounds Chlorothalonil Per Gallon

ACTIVE INGREDIENT:
Chlorothalonil (tetrachloroisophthalonitrile)*.....54.00%
INERT INGREDIENTS:.....46.00%
TOTAL 100.00%

*Covered under U.S. Patent No. 3,290,353 and No. 3,381,785

A Broa

READ ENT

BRAVO 6F is an excellent fungicide when used a
be used effectively in dilute or concentrate spray
DO NOT combine BRAVO 6F in the spray tank
physically compatible and non-injurious under y
NOTE: Slowly invert container several times to
The required amount of BRAVO 6F should be a
amount of BRAVO 6F in a clean container and
during spray operations.
DO NOT use on greenhouse-grown crops.

Dosage rates on this label indicate
to obtain adequate coverage of foli
range from 20 to 150 gallons per
applications.

BEANS (SNAP): Rust—Use 3 pints per acre in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when disease first threatens and repeat at weekly intervals or as necessary to maintain control. DO NOT apply within 7 days of harvest. DO NOT graze treated areas or feed treated plant parts to livestock.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS: Alternaria leaf spot (Southeastern states only) and downy mildew—Use 1½ pints per acre in sufficient water to obtain adequate coverage. Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals or as necessary to maintain control. Ring spot on field-seeded brussels sprouts (California only)—Use 2 pints per acre. Begin applications at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals or as necessary to maintain control.

CELERY: Early blight and late blight—Use 1½ to 2 pints per acre in sufficient water to obtain adequate coverage. Start applications when disease threatens and repeat at 7 to 10 day intervals or as necessary to maintain control.

CORN: Early blight, late blight, basal stalk rot, Helminthosporium leaf spot—Use 1 to 1½ pints per acre on 5 to 6 day spray schedule, or use 2 to 3 pints per acre on a 7 day schedule. Supression of pint rot—Use 3 pints per acre on a 7 day schedule. Start applications when transplants are set in the field. Apply BRAVO 6F in sufficient water to obtain adequate coverage. DO NOT apply within 7 days of harvest.

For control of early blight and late blight in celery seedbeds, use 1½ to 2 pints BRAVO 6F per 100 gallons of water and apply 125 gallons per acre twice weekly or as needed to maintain control. Start applications shortly after crop emergence. Use the higher rate under severe disease conditions.

CORN (SWEET): Helminthosporium leaf blight—Use 1½ to 2 pints per acre in sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development and repeat at 4 to 7 day intervals or as required to maintain control.

DO NOT apply with
fresh market crop
corn to be proces
graze in treated fi
or use as livestock.
CUCUMBERS: Tar
nose and downy
acre. Powdery
states), gummy st
Use 2 to 3 pints p
rot (Rhizoctonia
Apply BRAVO 6F
adequate covera
plants are in first t
are favorable for
applications at
disease conditions
CANTALOUPE,
MELON: Anthracn
to 2 pints per a
Southwestern S
gummy steam blig
3 pints per acre
wat tain a
tions n plants
condit are fa
Repeat applicati
severe disease co
WATERMELON,
nose and downy
acre. Cercospora
blight and scab—
BRAVO 6F in su
coverage. Begin
first true leaf state
for disease devel
day intervals. U
shorten spray inte
PEANUT: Cercos
per acre. Rust—
BRAVO 6F in su
coverage. Start
appears and repe
necessary to m
disease condition
DO NOT apply w
allow livestock to
hay or threshinge

AGRICULT

LJA Rev. 077-TX-1

BRAVO® 6F

A Broad-Spectrum Agriculture Fungicide

READ ENTIRE LABEL CAREFULLY AND USE ONLY AS DIRECTED

GENERAL INFORMATION

BRAVO 6F is an excellent fungicide when used according to label directions for control of a broad spectrum of plant diseases. BRAVO 6F can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control.

DO NOT combine BRAVO 6F in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible and non-injurious under your conditions of use.

NOTE: Slowly invert container several times to assure uniform mixture.

The required amount of BRAVO 6F should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of BRAVO 6F in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

DO NOT use on greenhouse-grown crops.

DIRECTIONS FOR USE

Dosage rates on this label indicate pints of BRAVO 6F per acre, unless otherwise stated. Applications should be made in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth. Spray volume usually will range from 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground sprays and aircraft applications.

BEAN (SNAP): Rust—Use 3 pints per acre in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when disease first threatens and repeat at weekly intervals or as necessary to maintain control. DO NOT apply within 7 days of harvest. DO NOT graze treated areas or feed treated plant parts to livestock.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS: Alternaria leaf spot (Southeastern states only) and downy mildew—Use 1½ pints per acre in sufficient water to obtain adequate coverage. Begin applications after transplants are set in field or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals or as necessary to maintain control. Ring spot on field-seeded brussels sprouts (California only)—Use 2 pints per acre. Begin applications at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals or as necessary to maintain control.

CARROT: Early blight and late blight—Use 1½ to 2 pints per acre in sufficient water to obtain adequate coverage. Start applications when disease threatens and repeat at 7 to 10 day intervals or as necessary to maintain control.

CELERY: Early blight, late blight, basal stalk rot (Rhizoctonia solani)—Use 1 to 1½ pints per acre on a 5 day spray schedule, or use 2 to 3 pints per acre on a 7 day schedule. Supression of pint rot—Use 3 pints per acre on a 7 day schedule. Start applications when transplants are set in the field. Apply BRAVO 6F in sufficient water to obtain adequate coverage. DO NOT apply within 7 days of harvest.

For control of early blight and late blight in celery seedbeds, use 1½ to 2 pints BRAVO 6F per 100 gallons of water and apply 125 gallons per acre once weekly or as needed to maintain control. Start applications shortly after crop emergence. Use the higher rate under severe disease conditions.

CORN (SWEET): Helminthosporium leaf blight—Use 1½ to 2 pints per acre in sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development and repeat at 4 to 7 day intervals or as required to maintain control. DO NOT apply within 14 days of harvest. For use on fresh market crop only. DO NOT apply to sweet corn to be processed. DO NOT allow livestock to graze in treated fields. DO NOT ensile treated corn or use as livestock forage.

CUCUMBER: Target spot (Florida only), anthracnose and downy mildew—Use 1½ to 2 pints per acre. Powdery mildew (except Southwestern states), gummy stem blight, leaf blight and scab—Use 2 to 3 pints per acre. For suppression of fruit rot (Rhizoctonia solani)—Use 3 pints per acre. Apply BRAVO 6F in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7-day intervals. Under severe disease conditions, shorten spray interval.

CANTALOUPE, MUSKMELON, HONEYDEW MELON: Anthracnose and downy mildew—Use 1½ to 2 pints per acre. Powdery mildew (except in Southwestern States), Cercospora leaf spot, gummy steam blight, leaf blight and scab—Use 2 to 3 pints per acre. Apply BRAVO 6F in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals. Under severe disease conditions, shorten spray interval.

WATERMELON, SQUASH, PUMPKIN: Anthracnose and downy mildew—Use 1½ to 2 pints per acre. Cercospora leaf spot, gummy stem blight, leaf blight and scab—Use 2 to 3 pints per acre. Apply BRAVO 6F in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf state or when conditions are favorable for disease development. Repeat applications at 7 day intervals. Under severe disease conditions, shorten spray interval.

PEANUT: Cercospora leaf spots—Use 1 to 1½ pints per acre. Rust—Use 1½ pints per acre. Apply BRAVO 6F in sufficient water to obtain adequate coverage. Start applications when disease first appears and repeat at 10 to 14 day intervals or as necessary to maintain control. Under severe disease conditions, use the 1½ pints per acre rate. DO NOT apply within 14 days of harvest. DO NOT allow livestock to graze treated areas. DO NOT feed hay or threshings from treated fields to livestock.

POTATO: Early blight, late blight, Botrytis vine rot (Botrytis spp.)—Use 1 to 1½ pints per acre in sufficient water to obtain adequate coverage. Begin applications when plants are 6 inches high or when disease threatens and continue at 7 to 10 day intervals or as needed to maintain disease control. Under severe disease conditions, use the 1½ pints per acre rate and shorten spray interval.

TOMATO: Early blight, late blight, gray leaf spot and gray leaf mold—Use 1½ to 2 pints per acre. Anthracnose—Use 2 to 3 pints per acre. Rhizoctonia fruit rot and Botrytis gray mold—Use 3 pints per acre. Apply BRAVO 6F in sufficient water to obtain adequate coverage. Begin applications when disease threatens and repeat at 7- to 10-day intervals. Under severe disease conditions, shorten spray interval. Alternaria fruit rot or black-mold (California only)—Make three applications at the rate of 3 pints per acre on a 10- to 14-day schedule beginning 5 to 6 weeks before anticipated harvest.

CONIFERS: Applications of BRAVO 6F with ground equipment are recommended for control of the following needle diseases: Lophodermium needlecast—Apply at the rate of 2½ pints per 100 gallons of water in hydraulic spray equipment or 3 qts. per 100 gallons in high-pressure mist blower. Begin applications in early August in the North Central and Northeastern states before infection occurs. Three sprays applied at 3 to 4 week intervals are usually adequate to control this disease. Scirrhia brown spot—Apply at rate of 3 qts. per 100 gallons of water in high-pressure mist blower. Begin applications when new needles are approximately one-half their full grown length. If rainfall is heavy, apply two additional sprays at three week intervals. Scleroderris canker—Use 1½ qts. per 100 gallons in hydraulic spray equipment or 3 qts. per 100 gallons in high-pressure mist blower to nursery stock. Make first application in the spring as soon as new growth appears. Repeat at 2 to 3 week intervals until early July. Continue applications at 4 week intervals until early September. Sirococcus tip blight and Phoma spp. (West Coast only) Apply at rate of 1 qt. per 100 gallons in hydraulic spray equipment. Start applications in early November and repeat at 2 to 4 week intervals as needed to maintain disease control. Use the shorter interval when disease conditions are moderately severe to severe. Cool, damp conditions are favorable for Phoma spp.

DANGER

CAUSES EYE DAMAGE

DO NOT get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for at least 15 minutes. Call a physician.

Avoid contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using.

DO NOT take internally.

Avoid breathing spray mist.

NOTE: BRAVO 6F may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Affected persons respond to treatment with antihistamines, steroid creams and/or systemic steroids.

DO NOT store feed near feed or foodstuffs. Store in a cool place. Protect from excessive heat.

DO NOT reuse empty container. Destroy by perforating, crushing, and burying or discarding in a safe place.

Apply only to areas specified on label.

This product is toxic to fish. Keep out of lakes, streams or ponds. DO NOT apply where runoff is likely to occur. DO NOT apply when weather conditions favor drift from areas treated. DO NOT contaminate water by cleaning of equipment or disposal of wastes.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and BUYER assumes the risk of any use contrary to such directions. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY [illegible] SELLER [illegible] authorized [illegible] except in writing and with specific reference to this warranty [illegible] shall [illegible]

AGRICULTURAL CHEMICALS DIVISION-CLEVELAND, OHIO 44114

DIAMOND SHAMROCK CORPORATION

EPA Est. 677-TX-1

3300-M-5-174