



Agricultural Chemicals Division

# Bravo 500



**Diamond Shamrock**

## Supplemental Labeling

SUPPLEMENTAL DIRECTIONS FOR THE USE OF

BRAVO 500  
(EPA Reg. No. 677-313)

**ACCEPTED**  
SEP 20 1982  
Under the Federal Insecticide,  
Fungicide and Rodenticide Act,  
as amended for the pesticide  
registered under  
EPA Reg. No. 677-313

FOR CONTROL OF DISEASES ON PEACH, NECTARINE,  
APRICOT, CHERRY, PLUM AND PRUNE

### Directions for Use on Tree and Orchard Crops

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply BRAVO 500 in sufficient water and proper calibration to obtain uniform coverage of tree canopy. Application through ground equipment is recommended. When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate of BRAVO 500 listed may be used. The following spray volumes are recommended as gallons of spray per acre:

CROP	SPRAY VOLUME (Gallons per Acre)	
	Dilute	Concentrate
Peach, Nectarine, Apricot, Tart Cherry, Plum, Prune	300	50 to 150
Sweet Cherry	400	65 to 200

Warranty and Limitation of Remedies

Implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of seller is authorized to do so except in writing with a specific reference to this warranty.

Seller makes no other express or implied warranty.

CROP	DISEASES	BRAVO 500 ACRE	RATE PER 100 GAL*	APPLICATION DIRECTIONS
Peach, Nectarine, Apricot, Cherry, Plum, Prune	Leaf curl, Coryneum blight (shothole) on peach, nectarine, apricot	4½ to 6 pts	1½ to 2 pts	Use only full dilute spray volumes for dormant applications. Make one application in late autumn to early winter before hard freezing occurs. Make one or two additional applications in mid to late winter before buds begin to swell. Where Coryneum blight (shothole) occurs, apply once at petal fall or at shuck-split to prevent fruit infections
	Brown rot blossom blight	4½ to 8 pts	1½ to 2 pts	Use 6 to 8 pints per acre on trees taller than 20 ft. and 4½ to 6 pints per acre on smaller trees. Make one application at popcorn (pink, red or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
	Cherry leafspot; peach, nectarine, apricot scab	4½ to 6 pts	1½ to 2 pts	In addition to the bloom applications listed above, make one application at shuck-split. DO NOT apply BRAVO 500 after shuck-split and before harvest. If additional disease control is needed before harvest, use another registered fungicide.  For control of cherry leafspot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leafspot incidence, make a second application 10-14 days later.

\*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

NOTE: It is important to read and observe the precautionary statements and all other information appearing on the BRAVO 500 label. Use only as directed.

NOTE: This labeling must be in the possession of the user at the time of pesticide application.