

DACONATE® 6

GALLONS

MSMA plus Surfactant For Postemergence Weed Control
READ ENTIRE LABEL BEFORE USING THIS PRODUCT

EPA REG. NO. 677-317-AA

GENERAL INFORMATION

DACONATE 6 TURF HERBICIDE is a postemergence herbicide for use on turf and certain non-crop areas. DACONATE 6 contains a surfactant (wetting agent) in proper ratio for best results. It is not necessary to add additional surfactant to the spray tank. Local conditions and recommendations vary; consult local agricultural experiment station or extension service weed specialists for recommendations on use of this product in your area.

A partial list of weeds controlled with DACONATE 6 TURF HERBICIDE includes:

(P1) Bahiagrass	Crabgrass	Nutsedge
Barnyardgrass	Dallisgrass	Flower
Brachiaria spp.	Foxtail	Puncturevine
Bullmoose	Goosegrass	Ragweed
Chickweed	Johnsongrass	Sandbur
Cocklebur	Morningglory	Wood sorrel

MIXING INSTRUCTIONS: DACONATE 6 TURF HERBICIDE must be thoroughly dissolved. Fill the spray tank with half the required amount of water and, with pump or agitator operating, add the recommended amount of DACONATE 6 TURF HERBICIDE, then continue filling the tank with balance of water and mix thoroughly.

If preferred, DACONATE 6 TURF HERBICIDE can be pre-mixed by dissolving the required amount of DACONATE 6 in 3 to 5 gallons of water in a clean, open-head bucket. This pre-mix should then be poured in the partially filled spray tank and the tank filled with balance of water. Mix thoroughly before using.

DO NOT store spray solution in tank overnight. Clean application equipment thoroughly after use by flushing with water in a safe place.

APPLICATION METHODS: DACONATE 6 TURF HERBICIDE should be applied with a low-volume, properly calibrated sprayer having satisfactory pumping and bypass action. ADJUST NOZZLES IN A MANNER TO ALLOW MAXIMUM COVERAGE OF WEEDS. Adequate coverage is very important for effective weed control.

Apply during warm weather when weeds are in an active stage of growth.

DIRECTIONS FOR USE

(S2) **SOLE COURSE AND ORNAMENTAL TURFGRASS:** DACONATE 6 TURF HERBICIDE can be used for selective control of bahiagrass, barnyardgrass, chickweed, smooth and large crabgrass, dallisgrass, nutsedge, sandbur, and wood sorrel with little or no injury to well established actively growing turfgrasses. Mow turfgrass to height of 1 to 1½ inches before treatment. Mix 1 fluid ounce (2 tablespoons) of DACONATE 6 in 5 gallons of water for application to an area of 1,000 square feet. Application should be uniform and thorough to adequately wet all undesirable plants.

Two or more repeat treatments at 14-day intervals may be necessary. Make applications during warm weather when temperature is between 80 and 90°F. DO NOT water turf for at least 24 hours after application. Turfgrasses may be temporarily discolored. Bermudagrass, bluegrass, and zoysiagrass have shown tolerance to properly applied DACONATE 6. Injury may result if applied to bentgrasses and fescues. DO NOT apply to St. Augustinegrass, carpetgrass, centipede grass, or to Dichondra lawns. DO NOT reseed until 2 weeks after last application.

(S1) **NON-CROP:** DACONATE 6 TURF HERBICIDE is effective in control of the above listed weeds and many similar weeds on drainage ditch banks, fence rows, sand traps, storage yards and similar non-crop areas. Application should be made when weeds are small and conditions are favorable for good weed growth. Mix at a rate of 2½ to 2¾ pints of DACONATE 6 in 100 gallons of water for application to one acre. Spray undesirable vegetation thoroughly to point of run-off. Adequate coverage and complete wetting of foliage is important for effective control. Repeat applications may be necessary if regrowth occurs. Use only as spot treatment in Florida.

CAUTION

Harmful if swallowed.

Avoid contact with skin and eyes. In case of contact, flush eyes with plenty of water.

Avoid breathing spray mist.

Wash hands after handling or using.

Keep away from children and domestic animals and do not allow them on treated areas until material has been washed into the soil.

DO NOT store near food and feed products.

DO NOT feed treated clippings to livestock or graze treated areas.

DO NOT spray or allow drift onto vegetables, ornamentals or other desirable plants.

DO NOT contaminate waters used for domestic purposes or by animals, including wildlife and aquatic life, or water used for irrigation purposes.


DO NOT reuse empty container. Return to drum reconditioner, or destroy by perforating, crushing or burying or discarding in a safe place away from water supplies.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and BUYER assumes the risk of any use contrary to such directions. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of SELLER is authorized to do so except in writing with a specific reference to this warranty. In no event shall SELLER'S liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Diamond

Shamrock



Chemical

Company


DACONATE® 6

TURF HERBICIDE

MSMA plus Surfactant
For Postemergence Weed Control

ACTIVE INGREDIENT:	
Monosodium Acid Methanesulfonate.....	48.35%
INERT INGREDIENTS.....	51.65%
TOTAL.....	100.00%

Total Arsenic, all in water-soluble form,
expressed as elemental ... 22.38%
Product contains 6.0 pounds MSMA per gallon



ACCEPTED

April 18, 1974

FEDERAL INSECTICIDE
AND FUNGICIDE
REGISTRY
677-317

CAUTION:

KEEP OUT OF REACH
OF CHILDREN
SEE SIDE PANEL FOR
OTHER PRECAUTIONS.

AGRICULTURAL CHEMICALS DIVISION-CLEVELAND, OHIO 44114

DIAMOND SHAMROCK CORPORATION