

FREIGHT CLASSIFICATION #312  
TEXTILE SOFTENERS NOI (DRY)

Net Weight      lbs.

# TEX-FLUFF

WITH

A FABRIC CONDITIONING AGENT

USE DRY ● APPLY DIRECT TO WHEEL ● GIVES A FINE SOFT FINISH ● CONTAINS A POTENT BACTERIOSTATIC AGENT

## DANGER

KEEP OUT OF REACH OF CHILDREN

*FOR INDUSTRIAL USE ONLY*



**Diamond Shamrock  
Chemical Company**

TEXTILE CARE DEPARTMENT  
CLEVELAND, OHIO 44114

DIAMOND SHAMROCK CORPORATION

### FOR TECHNICAL USE ONLY

#### ACTIVE INGREDIENTS:

|   |      |
|---|------|
| n-Alkyl (60%C., 30%C., 5%C., 5%C.)      |      |
| Dimethyl Benzyl Ammonium Chloride       | 8.0% |
| n-Alkyl (68%C., 32%C.)                  |      |
| Dimethyl Ethylbenzyl Ammonium Chlorides | 8.0% |

INERT INGREDIENTS:\* 84.0%

\*includes fabric softeners

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

## DANGER

KEEP OUT OF REACH OF CHILDREN. CORROSIVE. CAUSES EYE DAMAGE AND SKIN IRRITATION. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. WEAR GOGGLES OR FACE SHIELD AND RUBBER GLOVES WHEN HANDLING. HARMFUL OR FATAL IF SWALLOWED. AVOID CONTAMINATION OF FOOD.

### FIRST AID:

In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink PROMPTLY a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

### Directions For Use:

1. Use dry to the wheel.
2. For loads up to 100 lb. use 3 oz. per 100 lb. dry weight clothes.
3. For loads of more than 100 lb. use 2 oz. per 100 lb. dry weight clothes.
4. Can be used in starch bath or sour bath.
5. Can be used in automatic equipment.

E.P.A. REGISTRATION NO. 677-323  
E.P.A. Est. 677-MD-01

### NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.