

Precautionary Statements

Hazards to Humans

Caution

Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash eyes with plenty of water. If irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Do not apply in dwellings.

Environmental Hazards

This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a dry place.

Pesticide Disposal: Pesticide that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into mixing equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and Local authorities.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing and with specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Net Weight 12 1/2 Pounds/5.65 kilograms



Diamond Shamrock

Rabon[®] Livestock Dust

A Ready-To-Use Insecticide Dust

Controls Horn Flies, Cattle Grubs and Lice and aids in control of Face Flies on Beef Cattle and Dairy Cattle and controls Lice on Swine

Active Ingredient:	By Weight
2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	3.00%
Inert Ingredients:	97.00%
Total:	100.00%

Keep Out of Reach of Children

CAUTION

See other precautions on side panel

*Rabon Insecticide
Rabon is a registered trademark of Shell Oil Company

Diamond Shamrock Corporation
Animal Health Business
1100 Superior Avenue
Cleveland, Ohio 44114

EPA Reg. No. 677-3
EPA Est. No.

DSL-899 4/82

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Rabon Livestock Dust is a ready to use insecticide dust which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

Animal	Insect	Remarks
Beef Cattle & Dry Cattle	Horn Flies Face Flies & Lice	<p>HAND DUSTING Apply approximately 2 1/2 oz of dust by shaker, can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in control of face flies. Rub thoroughly to carry the dust beneath the hair, if possible, as necessary.</p> <p>SELF-TREATING DUST BAG - FORCED USE Put Rabon Livestock Dust in cotton cloth or double burlap bags or use prepacked weather proof cattle dust bags and hang in door exits or alleyways leading from animal buildings, salt or mineral blocks or watering holes. Protect cloth or burlap bags from weather.</p> <p>FREE CHOICE USE Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice aids in the control of lice.</p>
	Cattle Grubs	After the grubs have encysted (formed warbles) apply 3-4 oz down the backline and rub in thoroughly, taking care to get the dust into the warble if possible.
Same	Lice	HAND DUSTING - Apply 3-4 oz of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% dust should be applied per 150 sq. ft. of bedding.