



PARATHION 2 DUST

OBSERVE THE FOLLOWING USE PRECAUTIONS

Wear natural rubber gloves, protective clothing and goggles.
Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection.
Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated.
Do not contaminate feed and foodstuffs.
Wash hands, arms, and face thoroughly with soap and water before eating or smoking.
Wash all contaminated clothing with soap and hot water before re-use.

Observe use limitations given on the label for specific crops.

DO NOT apply during bloom when honey bees or other pollinators are active. Avoid dust drift to adjoining food and forage crops about to be harvested, or crops in bloom.

Do not apply to fish and wildlife areas, including treated areas, rivers, lakes, ponds, or any body of water, or to any growing susceptible crops.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

Consult local agricultural authorities for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

DIRECTIONS FOR USE

This product is an insecticide dust formulation for use by the commercial grower or commercial applicator. Apply when air is calm. To obtain most effective results, apply with a ground duster or by airplane. Make thorough coverage of plants with a uniform deposit of dust. Apply at the first sign of infestation and repeat at 7 to 10 day intervals when needed to maintain control, but observe use limits given for specific crops.

BEANS—Aphids, armyworms, corn earworm, cucumber beetle, leafhopper, leaf miner, Mexican bean beetle, spider mites, white fly—Apply up to 25 lbs. per acre. Do not apply within 15 days of harvest.

BEETS, SWEET—Aphids, leaf miner, webworms—Apply up to 17 lbs. per acre. Do not apply within 15 days of harvest.

BROCCOLI, CAULIFLOWERS, BRUSSELS SPROUTS—Aphids, leaf miner, webworms, cabbage root maggot, black root, larvae, seedling, leaf miner, thrips—Use 25

lbs. per acre at first sign of infestation and repeat as needed. Do not apply more than 7 days before harvest.

ACTIVE INGREDIENT

Parathion (D, O diethyl O p nitrophenyl thiophosphate)

INERT INGREDIENTS

7%

98%

100%



WARNING



KEEP OUT OF REACH OF CHILDREN

PARATHION IS A DEADLY POISON. IT KILLS MAN AND ANIMALS.
KEEP IT OUT OF REACH OF CHILDREN.

PARATHION IS A DEADLY POISON. IT KILLS MAN AND ANIMALS.
KEEP IT OUT OF REACH OF CHILDREN.

POISONING SYMPTOMS: Irritation, blurred vision, weakness, nausea, vomiting, diarrhea, and convulsions.

SYMPTOMS OF POISONING: Obtain emergency medical attention immediately. Keep on hand.

SEE A PHYSICIAN IMMEDIATELY if persons handling or working with parathion become ill.

FIRST AID TREATMENT

IF SWALLOWED: Give plenty of water to drink. Wash mouth out with water. Do not induce vomiting.

IF IN EYES: Remove dust from contaminated atmosphere. Flush eyes with water for 15 minutes.

IF ON SKIN: Remove dust from contaminated atmosphere. Wash skin with soap and water.

IF INHALED: If contact, flush immediately with water for at least 15 minutes.

IF INFORMATION FOR PHYSICIAN: Give symptoms, preferably give product name and amount ingested. Give every hour or other as directed.

IF SWALLOWED: Give plenty of water to drink. Wash mouth out with water. Do not induce vomiting.

NET CONTENT 6-2-3

ST

King

EN

1990

CUCURBITS (Cucumbers, Melon, Squash)—Apply only when cabbage looper, squash beetles, flea beetle, leafhoppers, leafminer, melon worm, pickleworm, spider mite, thrips, white fly—Apply ~~to~~ to 40 lbs. per acre. Apply only when plants are dry. Do not apply before vining or within 15 days of harvest. *Apply only when plants are dry. Do not apply before vining or within 15 days of harvest.*

• **UGPLANT** -Aphids, serpentine leaf miner, spider mites -Apply up to 22 lbs per acre
 • **Do not apply within 15 days of harvest.**

FORAGE CROPS (Alfalfa, barley, clover, oats, vetch, wheat)—Aphids, armyworms, blister beetles, grasshoppers. Apply ~~up to~~ 2 lbs per acre. Do not apply when crops are in bloom. Do not apply within 15 days of harvest or forage use.

CAUTION—Aphids, armyworms, cabbage looper, vegetable weevil, serpentine leaf miner.
Apply ~~4.0 to 8.0~~ lbs. per acre. Do not apply within ~~7~~ days of harvest.

CAUTIONS—Injury—Apply ~~up to~~ ¹⁰ 10 lbs. per acre. Do not apply on seed crops after the flowers are open because of hazard to pollinators. Do not apply within 14 days of harvest.

PEAS - Aphids, pea weevil, serpentine leaf miner, thrips - Apply ~~20~~ 30 lbs. per acre. Do not apply when plants are in bloom. Do not apply within 10 days of harvest. 15 ~~20~~ 16

PEPPERS- Aphids, flea beetles, pepper maggot, serpentine leaf miner. Apply insecticide per acre. Do not apply within 15 days of harvest.

ROTATIONS - Aphids, armyworms, Colorado potato beetle, flea beetle, leafhopper, spring
line leaf miner, vegetable weevil, white fly. Apply 20-25 lbs. per acre. Do not apply
within 3 days of harvest.

FINACH: Aphids—Apply ~~up to~~ 20 per acre. Do not apply within 14 days of harvest.

STRAWBERRY: Aphids and spider mites usually up to 20% are more or less active in the field.

~~MAIO~~ Aphids, army worms, flea beetle, leather beetle, serpentine leaf miner, etc. Do not apply within 10 days of harvest.

42-2

13. IF THEY EMIT ANY FUEL BY BURNING, LIST - FUEL TYPE, USE, LOCATION

KEEP OUT OF RANGE OF FUMES

CP, INC. (1943-1944)