



ACCEPTED
5-23-68
682-53
 UNDER THE FEDERAL INSECT
 FERTILIZER ACT, 1948, AS AMENDED
 FOR FERTILIZERS, FERTILIZER REGISTRATION
 ED UNDER THE ACT, AND SUBJECT
 TO ALL APPLICABLE COMMENTS.

SEVIN® 5% DAMP DUST™
 *Trade Mark of Union Carbide and Carbon Company



OBSERVE THE FOLLOWING USE PRECAUTIONS
 This material is **TOXIC TO HONEY BEES**. DO NOT APPLY TO CROPS DURING BLOOM WHEN HONEY BEES OR OTHER POLLINATORS ARE ACTIVE. AVOID DRIFT TO ADJOINING FOOD AND FORAGE CROPS ABOUT TO BE HARVESTED OR CROPS IN BLOOM.

CROPS

BEANS: Mexican Bean Beetle — Use 20-25 pounds per acre. Leaf Hoppers — use 25-30 pounds per acre. DO NOT EXCEED 42 pounds per acre. No time limitations.

CORN: Corn earworm and European Corn Borer — use 40 pounds per acre at two to three day intervals, depending upon the growth of silk, and degree of infestation. No time limitation.

GRAPES: Grape leaf folder, Grape leafhopper, Grape Berry Moth, Japanese Beetle — use 50 pounds 5% dust per acre at first sign of infestation and repeat as necessary. Do not exceed 50 pounds per acre. No time limitation.

LETTUCE (Outdoor) Head: Cabbage looper (1st and 2nd instar), Imported Cabbag Moth, Corn earworm, Leafhoppers, grasshoppers — use 30-40 pounds per acre at first sign of infestation and repeat if necessary. DO NOT EXCEED 40 POUNDS PER ACRE. Last application no closer to harvest than 3 days.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methyl-carbamate)	5.0%
INERT INGREDIENTS	95.0%
	100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

- Harmful if swallowed, or inhaled.
- Avoid prolonged breathing of dust.
- Avoid prolonged or repeated contact with skin.
- Wash thoroughly after using.
- Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

AVOID STORAGE NEAR FEED OR FOODSTUFFS
SAMPLE
 CONTENTS U.S.D.A. Reg. 682-53

LABEL

LETTUCE (Outdoor) Leaf: For the control of the same insects as for head lettuce. Use 40 pounds per acre. Do not exceed this rate. Last application no closer than 14 days prior to harvest.

ORNAMENTALS: For control of various beetle larvae, dust foliage at weekly intervals. Cicada, periodical — on foliage when insects appear. Japanese beetle, leafhoppers, rose chafer, Rose curculio, rose slug, use 2 pounds dust per 1000 square feet of area. Dust lightly but thoroughly on foliage. Repeat when necessary. OBSERVE USE PRECAUTIONS.

POTATOES: Flea beetles and Leaf Hoppers, Colorado Potato beetle, Cucumber Beetle, Flea Beetles, Lygus bugs, use from 30 to 40 pounds per acre. Foliage application only. Use at first sign of infestation and repeat as necessary. DO NOT EXCEED 40 POUNDS PER ACRE PER APPLICATION.

RUTABAGA: For control of Leafhoppers, Flea-beetle, corn earworm, climbing cutworms, use 40 pounds per acre as needed for control of infestation. DO NOT EXCEED 40 POUNDS PER ACRE. Do not apply closer to harvest than 3 days.

SPINACH: For control of Cucumber beetles, Leaf

hopper, Armyworm, use 30-40 pounds per acre. Do not exceed 40 pounds per acre. Last application should be no closer than 14 days before harvest.

STRAWBERRIES: Meadow spittlebugs, leaf roller — use 30 to 40 pounds per acre as required for infestation. Repeat as necessary. DO NOT EXCEED 40 pounds per acre. Last application no closer to harvest than 1 day.

SWISS CHARD: Beet webworm, Flea Beetles, Army worm — use 30-40 pounds per acre as required by infestation and foliage density. Repeat as required. DO NOT EXCEED 40 pounds per acre per application. Do not use closer to harvest than 14 days.

SUGAR BEETS: For control of Alfalfa Loopers, Armyworm, Flea beetles, Flea beetles, use 30-40 pounds per acre. Last application no closer to harvest than 14 days.

TOMATOES: Army worms, Corn earworm, flea beetles, Cucumber Beetle, Hornworm, Colorado Potato Beetle, Stink bugs, Climbing Cutworms, use 40-75 pounds per acre as required by infestation and foliage density. Repeat as necessary for control. No time limitation. DO NOT EVER EXCEED 75 POUNDS PER ACRE.

IMPORTANT NOTICE TO PURCHASER AND USER OF THIS PRODUCT — This product is manufactured, labeled and sold in accordance with state and federal laws. Adequate directions for use, necessary warnings and cautions, and other essential information on the safe handling and use of this product are set forth on this label. However, the use and handling of this product is always attended with risk of personal injury or property damage if the user or handler departs to any degree from the instructions on this label. The manufacturer makes no warranties, either expressed or implied, and assumes no responsibility whatsoever for the effectiveness or safety of this product except to the extent that the product is used and handled in strict accordance with information and directions set forth on this label. The foregoing limitation of responsibility may not be altered in any way except by written agreement signed by officers of the manufacturer. Unless the intended purchaser agrees to be bound by the terms and limitations expressed herein and on this label, the manufacturer refuses to consummate the sale of this product to him and the product must be returned to the manufacturer.
 CROP KING CHEMICALS, Yakima, Washington



5-23-68
682-53

SEVIN* 5% DAMP DUST T.M.

*Trade Mark of Union Carbide and Carbon Company

OBSERVE THE FOLLOWING USE PRECAUTIONS. This material is TOXIC TO HONEY BEES. DO NOT APPLY TO CROPS DURING BLOOM WHEN HONEY BEES OR OTHER POLLINATORS ARE ACTIVE. AVOID DRIFT TO ADJOINING FOOD AND FORAGE CROPS ABOUT TO BE HARVESTED OR CROPS IN BLOOM.

CROPS

BEANS: Mexican Bean Beetle -- Use 20-25 pounds per acre. Leaf Hoppers -- use 25-30 pounds per acre. DO NOT EXCEED 42 pounds per acre. No time limitations.

CORN: Corn earworm and European Corn Borer -- Use 40 pounds per acre at two to three day intervals, depending upon the growth of silk, and degree of infestation. No time limitation.

GRAPES: Grape leaf folder, Grape leafhopper, Grape berry moth, Japanese Beetle -- use 50 pounds 5% dust per acre at first sign of infestation and repeat as necessary. Do not exceed 50 pounds per acre. No time limitation.

LETTUCE (Outdoor) Head: Cabbage looper (1st and 2nd instar), Imported Cabbage worm, corn earworm, Leafhoppers, grasshoppers -- use 30-40 pounds per acre at first sign of infestation and repeat if necessary. DO NOT EXCEED 40 POUNDS PER ACRE. Last application no closer to harvest than 3 days.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methyl-carbamate) 5.0%

INERT INGREDIENTS

95.0%

100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, or inhaled.

Avoid prolonged breathing of dust.

Avoid prolonged or repeated contact with skin.

Wash thoroughly after using.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

AVOID STORAGE NEAR FEED OR FOODSTUFFS

SAMPLE

CONTENTS

U.S.D.A. Reg. 682-53

LETTUCE (Outdoor) Leaf: For the control of same insects as for head lettuce. Use 40 per acre. Do not exceed this rate. Last application no closer than 14 days prior to harvest.

ORNAMENTALS: For control of various beetle dust foliage at weekly intervals. Cicada, Japanese leafhoppers, rose chafer, Rose curculio, etc. use 2 pounds dust per 1000 square feet of area lightly but thoroughly on foliage. Repeat as necessary. OBSERVE USE PRECAUTIONS.

POTATOES: Flea beetles and Leaf Hoppers, Potato beetle, Cucumber Beetle, Flea Beetle bugs, use from 30 to 40 pounds per acre. application only. Use at first sign of infestation and repeat as necessary. DO NOT EXCEED 40 POUNDS PER ACRE PER APPLICATION.

RUTABAGA: For control of Leafhoppers, Flea beetle, corn earworm, climbing cutworms, use 40 pounds per acre as needed for control of infestation. DO NOT EXCEED 40 POUNDS PER ACRE. Do not apply closer to harvest than 3 days.

SPINACH: For control of Cucumber beetle

LABEL

IMPORTANT NOTICE TO PURCHASER AND USER OF THIS PRODUCT: This product is manufactured, labeled and sold in accordance with state and federal laws. Adequate directions for the safe handling and use of this product are set forth on this label. However, the use and handling of this product should be attended with a sense of personal injury or property damage. The manufacturer makes no warranty, either expressed or implied, as to the effectiveness of this product, or the effect upon the health of the user of this product, except as stated on the label. The foregoing limitation of responsibility may not be altered in any way except by written agreement signed by officials of the manufacturer. The manufacturer reserves the right to discontinue the sale of this product at any time and without notice. U.S.D.A. Reg. 682-53. CROP KING CHEMICALS

SEVIN* 5% DAMP DUST T.M.

*Trade Mark of Union Carbide and Carbon Company



INGREDIENTS:

aryl (1-naphthyl N-methyl- carbamate)	5.0%
INGREDIENTS	95.0%
	<hr/> 100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

- if swallowed, or inhaled.
- prolonged breathing of dust.
- prolonged or repeated contact with skin.
- Wash thoroughly after using.
- Use empty container. Destroy it by burying or burning. Stay away from smoke and

STORAGE NEAR FEED OR FOODSTUFFS

SAMPLE

U.S.D.A. Reg. 682-53

LABEL

This product is manufactured, labeled and sold in accordance with state and federal laws. Adequate directions for use, necessary warnings and other important information are provided on the label and packaging of this product. Always attend to these directions, warnings and other information. The degree of control may vary with the degree of infestation, the type of pest, the type of crop, the weather, and the user's responsibility. The user assumes responsibility for the effective use of this product. The manufacturer's liability for the effectiveness of this product is limited to the extent that the product is used in accordance with the directions on the label. The manufacturer's liability for the effectiveness of this product is limited to the extent that the product is used in accordance with the directions on the label. The manufacturer's liability for the effectiveness of this product is limited to the extent that the product is used in accordance with the directions on the label.

LETTUCE (Outdoor) Leaf: For the control of the same insects as for head lettuce. Use 40 pounds per acre. Do not exceed this rate. Last application no closer than 14 days prior to harvest.

ORNAMENTALS: For control of various beetle larvae, dust foliage at weekly intervals. Cicada, periodical — on foliage when insects appear. Japanese beetle, leafhoppers, rose chafer, Rose curculio, rose slug, use 2 pounds dust per 1000 square feet of area. Dust lightly but thoroughly on foliage. Repeat when necessary. OBSERVE USE PRECAUTIONS.

POTATOES: Flea beetles and Leaf Hoppers, Colorado Potato beetle, Cucumber Beetle, Flea Beetles, Lygus bugs, use from 30 to 40 pounds per acre. Foliage application only. Use at first sign of infestation and repeat as necessary. DO NOT EXCEED 40 POUNDS PER ACRE PER APPLICATION.

RUTABAGA: For control of Leafhoppers, Flea-beetle, corn earworm, climbing cutworms, use 40 pounds per acre as needed for control of infestation. DO NOT EXCEED 40 POUNDS PER ACRE. Do not apply closer to harvest than 3 days.

SPINACH: For control of Cucumber beetles, Leaf

hopper, Armyworm, use 30-40 pounds per acre. Do not exceed 40 pounds per acre. Last application should be no closer than 14 days before harvest.

STRAWBERRIES: Meadow spittlebugs, leaf roller — use 30 to 40 pounds per acre as required for infestation. Repeat as necessary. DO NOT EXCEED 40 pounds per acre. Last application no closer to harvest than 1 day.

SWISS CHARD: Beet webworm, Flea Beetles, Army worm — use 30-40 pounds per acre as required by infestation and foliage density. Repeat as required. DO NOT EXCEED 40 pounds per acre per application. Do not use closer to harvest than 14 days.

SUGAR BEETS: For control of Alfalfa Loopers, Armyworm, blister beetles, Flea beetles, use 30-40 pounds per acre. Last application no closer to harvest than 14 days.

TOMATOES: Army worms, Corn earworm, flea beetles, Cucumber Beetle, Hornworm, Colorado Potato Beetle, Stink bugs, Climbing Cutworms, use 40-75 pounds per acre as required by infestation and foliage density. Repeat as necessary for control. No time limitation. DO NOT EVER EXCEED 75 POUNDS PER ACRE.