



SEVIN® 5% DUST

*Trade Mark of Union Carbide and Carbon Company



OBSERVE THE FOLLOWING USE PRECAUTIONS
This material is **TOXIC TO HONEY BEES**. DO NOT APPLY TO CROPS DURING BLOOM WHEN HONEY BEES OR OTHER POLLINATORS ARE ACTIVE. AVOID DRIFT TO ADJOINING FOOD AND FORAGE CROPS ABOUT TO BE HARVESTED OR CROPS IN BLOOM.

CROPS

BEANS: Mexican Bean Beetle — Use 20-25 pounds per acre. Leaf Hoppers — use 25-30 pounds per acre. DO NOT EXCEED 42 pounds per acre. No time limitations.

CORN: Corn earworm and European Corn Borer — use 40 pounds per acre at two to three day intervals, depending upon the growth of silk, and degree of infestation. No time limitation.

GRAPES: Grape leaf folder, Grape leafhopper, Grape berry moth, Japanese Beetle — use 50 pounds 5% dust per acre at first sign of infestation and repeat as necessary. Do not exceed 50 pounds per acre. No time limitation.

LETTUCE (Outdoor) Head: Cabbage looper (1st and 2nd instar), Imported cabbage worm, corn earworm, Leafhoppers, grasshoppers — use 30-40 pounds per acre at first sign of infestation and repeat if necessary. DO NOT EXCEED 40 POUNDS PER ACRE. Last application no closer to harvest than 3 days.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methyl-carbamate) -----

5.0%

INERT INGREDIENTS -----

95.0%

100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed.

Avoid prolonged breathing of dust.

Avoid prolonged or repeated contact with skin.

Wash thoroughly after using.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

AVOID STORAGE NEAR FEED OR FOODSTUFFS

CONTENTS

U.S.D.A. Reg. 682-57

LETTUCE (Outdoor) Leaf: For the control of the same insects as for head lettuce. Use 40 pounds per acre. Do not exceed this rate. Last application no closer than 14 days prior to harvest.

ORNAMENTALS: For control of various beetle larvae, dust foliage at weekly intervals. Cicada, periodical — on foliage when insects appear. Japanese Beetle, leafhoppers, pickleworms, rose chafer, Rose curculio, rose slug, use 2 pounds dust per 1000 square feet of area. Dust lightly but thoroughly on foliage. Repeat when necessary. OBSERVE USE PRECAUTIONS.

POTATOES: Flea beetles and Leaf Hoppers, Colorado Potato beetle, Cucumber Beetle, Flea Beetles, Lygus bugs, use from 30 to 40 pounds per acre. Foliage application only. Use at first sign of infestation and repeat as necessary. DO NOT EXCEED 40 POUNDS PER ACRE PER APPLICATION.

RUTABAGA: For control of Leafhoppers, Flea-beetle, corn earworm, cutworms, use 40 pounds per acre as needed for control of infestation. DO NOT EXCEED 40 POUNDS PER ACRE. Do not apply closer to harvest than 3 days.

SPINACH: For control of Cucumber beetles, Leaf

hopper, Armyworm, use 30-40 pounds per acre. Do not exceed 40 pounds per acre. Last application should be no closer than 14 days before harvest.

STRAWBERRIES: Meadow spittlebugs, leaf roller — use 30 to 40 pounds per acre as required for infestation. Repeat as necessary. DO NOT EXCEED 40 pounds per acre. Last application no closer to harvest than 1 day.

SWISS CHARD: Beat webworm, Flea Beetles, Army worm — use 30-40 pounds per acre as required by infestation and foliage density. Repeat as required. DO NOT EXCEED 40 pounds per acre per application. Do not use closer to harvest than 14 days.

SUGAR BEETS: For control of Alfalfa Loopers, Armyworm, Blister beetles, Flea beetles, use 30-40 pounds per acre. Last application no closer to harvest than 14 days.


TOMATOES: Army worms, Corn earworm, flea beetles, Cucumber Beetle, Hornworm, Colorado Potato Beetle, Stink bugs, Climbing Cutworms, use 40-75 pounds per acre as required by infestation and foliage density. Repeat as necessary for control. No time limitation. DO NOT EVER EXCEED 75 POUNDS PER ACRE.

IMPORTANT NOTICE TO PURCHASER AND USER OF THIS PRODUCT — This product is manufactured, labeled and sold in accordance with state and federal laws. Adequate directions for use, necessary warnings and cautions, and other essential information on the safe handling and use of this product are set forth on this label. However, the use and handling of this product is always attended with risk of personal injury or property damage if the user or handler departs to any degree from the instructions on this label. The manufacturer makes no warranties, either expressed or implied, and assumes no responsibility whatsoever for the effectiveness or safety of this product except to the extent that the product is used and handled in strict accordance with information and directions set forth on this label. The foregoing limitation of responsibility may not be altered in any way except by written agreement signed by officers of the manufacturer. Unless the intended purchaser agrees to be bound by the terms and limitations expressed herein and on this label, the manufacturer refuses to consummate the sale of this product to him and the product must be returned to the manufacturer.

CROP KING CHEMICALS, Yakima, Washington

DAMP-DUST

30 LBS. NET WT.



SEVIN® 5% DUST

Trade Mark of Union Carbide and Chemical Company



ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methyl-
carbamate) ----- 5.0%

INERT INGREDIENTS ----- 95.0%
100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Dangerous if swallowed.

Avoid prolonged breathing of dust.

Avoid prolonged or repeated contact with skin.

Wash thoroughly after using.

Do not reuse empty container. Destroy it by burying
in waste or burning. Stay away from smoke and
flames.

DO NOT STORE NEAR FEED OR FOODSTUFFS

CONTENTS

U.S.D.A. Reg. 682-57

LETTUCE (Outdoor): Leaf For the control of the
same insects as for leafy lettuce. Use 40 pounds
per acre. Do not exceed this rate. Last application
no closer than 14 days prior to harvest.

ORNAMENTALS: For control of various beetle larvae,
dust foliage at weekly intervals. Check, periodically
— on foliage when insects appear. Japanese beetle,
leafhoppers, pickleworms, rose chafer, Rose cur-
culio, rose slug, use 2 pounds dust per 1000 square
feet of area. Dust lightly but thoroughly on foliage.
Repeat when necessary. **OBSERVE USE PRECAU-
TIONS.**

POTATOES: Flea beetles and Leaf Hoppers, Colorado
Potato beetle, Cucumber Beetle, Flea Beetles, Lygus
bugs, use from 30 to 40 pounds per acre. Foliage
application only. Use at first sign of infestation
and repeat as necessary. **DO NOT EXCEED 40
POUNDS PER ACRE PER APPLICATION.**

RUTABAGA: For control of Leafhoppers, Flea-beetle,
corn earworm, cutworms, use 40 pounds per acre as
needed for control of infestation. **DO NOT EXCEED
40 POUNDS PER ACRE.** Do not apply closer to har-
vest than 3 days.

SPINACH: For control of Cucumber beetles, Leaf

hoppers, Armyworms, use 30-40 pounds per acre. Do
not exceed 40 pounds per acre. Last application
should be no closer than 14 days before harvest.

STRAWBERRIES: Flea beetles, leaf miner —
use 30 to 40 pounds per acre as required for infesta-
tion. Repeat as necessary. **DO NOT EXCEED 40
pounds per acre.** Last application no closer to har-
vest than 1 day.

SWISS CHARD: Beet webworm, Flea Beetles, Army
worm — use 30-40 pounds per acre as required by
infestation and foliage density. Repeat as required.
DO NOT EXCEED 40 pounds per acre per application.
Do not use closer to harvest than 14 days.

SUGAR BEETS: For control of Alfalfa Loopers, Army-
worm, Blister beetles, Flea beetles, use 30-40 pounds
per acre. Last application no closer to harvest than
14 days.

TOMATOES: Army worms, Corn earworm, flea
beetles, Cucumber Beetle, Hornworm, Colorado Po-
tato Beetle, Stink bugs, Climbing Cutworms, use
40-75 pounds per acre as required by infestation
and foliage density. Repeat as necessary for con-
trol. No time limitation. **DO NOT EVER EXCEED
75 POUNDS PER ACRE.**

This product is manufactured, labeled and sold in accordance with state and federal laws. Adequate directions for use, necessary warnings and cautions, and other essential information
are provided on the label. However, the use and handling of this product is always attended with risk of personal injury or property damage if the user or handler departs to any degree from the instructions
on the label, or implied, and assumes no responsibility whatsoever for the effectiveness or safety of this product except to the extent that the product is used and handled in strict accordance with
the label. No responsibility may not be altered in any way except by written agreement signed by officers of the manufacturer. Unless the intended purchaser agrees to be bound by the
label, the manufacturer refuses to consummate the sale of this product to him and the product must be returned to the manufacturer.

CROP KING CHEMICALS, Yakima, Washington