



SEVIN® 5% DUST

Trade Mark of Union Carbide and Carbon Company

OBSERVE THE FOLLOWING USE PRECAUTIONS
 This material is **TOXIC TO HONEY BEES**. **DO NOT APPLY TO CROPS DURING BLOOM WHEN HONEY BEES OR OTHER POLLINATORS ARE ACTIVE**. **AVOID DRIFT TO ADJOINING FOOD AND FORAGE CROPS ABOUT TO BE HARVESTED OR CROPS IN BLOOM.**

CROPS

BEANS: Mexican Bean Beetle — Use 20-25 pounds per acre. Leaf Hoppers — use 25-30 pounds per acre. **DO NOT EXCEED 42 pounds per acre. No time limitations.**

CORN: Corn earworm and European Corn Borer — use 40 pounds per acre at two to three day intervals, depending upon the growth of silk, and degree of infestation. No time limitation.

GRAPES: Grape leaf folder, Grape leafhopper, Grape berry moth, Japanese Beetle — use 50 pounds 5% dust per acre at first sign of infestation and repeat as necessary. Do not exceed 50 pounds per acre. No time limitation.

LETTUCE (Outdoor) Head: Cabbage looper (1st and 2nd instar), Imported Cabbage worm, corn earworm, Leafhoppers, grasshoppers — use 30-40 pounds per acre at first sign of infestation and repeat if necessary. **DO NOT EXCEED 40 POUNDS PER ACRE.** Last application no closer to harvest than 3 days.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methyl-carbamate) ----- 5.0%

INERT INGREDIENTS ----- 95.0%
 100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed.

Avoid prolonged breathing of dust.

Avoid prolonged or repeated contact with skin.

Wash thoroughly after using.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

AVOID STORAGE NEAR FEED OR FOODSTUFFS

CONTENTS

U.S.D.A. Reg. 682-57

LETTUCE (Outdoor) Leaf: For the control of same insects as for head lettuce. Use 40 pounds per acre. Do not exceed this rate. Last application no closer than 14 days prior to harvest.

ORNAMENTALS: For control of various beetle dust foliage at weekly intervals. Cicada, perl — on foliage when insects appear. Japanese leafhoppers, pickleworms, rose chafer, Rosculio, rose slug, use 2 pounds dust per 1000 feet of area. Dust lightly but thoroughly on foliage. Repeat when necessary. **OBSERVE USE PRECAUTIONS.**

POTATOES: Flea beetles and Leaf Hoppers, Colorado Potato beetle, Cucumber Beetle, Flea Beetles, bugs, use from 30 to 40 pounds per acre. [application only. Use at first sign of infestation and repeat as necessary. **DO NOT EXCEED 40 POUNDS PER ACRE PER APPLICATION.**

RUTABAGA: For control of Leafhoppers, Flea-corn earworm, cutworms, use 40 pounds per acre needed for control of infestation. **DO NOT EXCEED 40 POUNDS PER ACRE.** Do not apply closer than 3 days.

SPINACH: For control of Cucumber beetles

IMPORTANT NOTICE TO PURCHASER AND USER OF THIS PRODUCT — This product is manufactured, labeled and sold in accordance with state and federal laws. Adequate directions on the safe handling and use of this product are set forth on this label. However, the use and handling of this product is always attended with risk of personal injury or property damage on this label. The manufacturer makes no warranties, either expressed or implied, and assumes no responsibility whatsoever for the effectiveness or safety of this product except to the information and directions set forth on this label. The foregoing limitation of responsibility may not be altered in any way except by written agreement signed by officers of the manufacturer. In accordance with the terms and limitations expressed herein and on this label, the manufacturer refuses to consummate the sale of this product to him and the product must be returned to the manufacturer.

CROP KING CHEMICALS