

ACCEPTED
 JUN 16 1972
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. **68258**



TEDION 400

FOR VOLUME SPRAYS on fruit trees use from 200-400 gallons of water per acre. Volume to be determined by age/size of trees and density of foliage, time of application and degree of insect infestation.

ROW CROPS: Use from 50-100 gallons of water per acre depending upon crop, foliage density, and insect infestation.

Airplane Application: Use 10 gallons of water per acre; where equipment and economy permit use 20 gallons per acre.

Consult State Extension Service Specialists for further information concerning local problems and conditions.

HOPS

For Two-spotted mite, apply 1 quart. DO NOT make more than 2 applications per season during fruiting period. DO NOT apply within 14 days of harvest. DO NOT feed straw to livestock.

PEPPERMINT AND SPEARMINT

For Two-spotted mite. Apply 2 1/2 quart. DO NOT make more than 2 applications per season. DO NOT apply within 10 days of cutting. DO NOT feed mint hay or spent mint hay to livestock.

FRUIT CROPS

APPLICATIONS MAY BE MADE UP TO DAY OF HARVEST

APPLE, CRABAPPLE, PEAR AND QUINCE: For European red, Two-spotted, McDaniel, Canadianis, Schoenii, and Carpini mites, use 1 1/4 to 2 1/2 quarts. DO NOT make more than 3 applications after petal fall. DO NOT feed pomace from treated apples to livestock.

CITRUS: (ORANGES, GRAPEFRUIT, TANGERINES, LEMONS, LIMES, TANGELOS, and CITRUS CITRON).

FOR CITRUS RED MITE, SIX-SPOTTED MITE and TEXAS CITRUS MITE: Use 2 to 2 1/2 quarts. Apply when mites first appear. Application may be made right up to harvest; however, DO NOT make more than one application per season while fruit is present.

APRICOT, CHERRY, NECTARINE AND PEACH: For European red, Two-spotted, and McDaniel mites, use 1 1/4 to 2 quarts. DO NOT make more than one application after shuck-rot.

MITICIDE

ACTIVE INGREDIENTS:
 *Tetraflon 40%

INERT INGREDIENTS
 *p-chlorophenyl 2, 4, 5-trichlorophenyl sulphone 60%

PROTECT FROM TEMPERATURES BELOW 32° F.
 (Product freezes but thaws unimpaired)

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified. DO NOT allow to drift to adjoining food or forage crops. DO NOT graze or feed contaminated cover crop to livestock.

Keep container closed. Leaking packages should be removed to a safe place.

Use with adequate ventilation.

Avoid breathing vapor.

Avoid contact with skin.

Shrimp and crab may be killed at application rates recommended on this label. Keep out of Tidal Marshes and Estuaries.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies.

Containers should be disposed of by punching holes in them and burying with wastes.

(See right hand panel for further cautions)

PLUMS AND PRUNES: Use 1 1/4 to 2 quarts per acre during fruiting period.

GRAPE: For Phlox mite DO NOT make more than 3 applications.

BUSHBERRIES (Blackberries, Raspberries): 1 mite first appear. Ret

STRAWBERRY: For Two less than 30 days during

fruiting period.

DO NOT apply within 14 days of harvest.

DO NOT feed straw to livestock.

DO NOT apply within 14 days of harvest.

DO NOT feed straw to livestock.

DO NOT apply within 14 days of harvest.

DO NOT feed straw to livestock.

DO NOT apply within 14 days of harvest.

DO NOT feed straw to livestock.

DO NOT apply within 14 days of harvest.

DO NOT feed straw to livestock.

DO NOT apply within 14 days of harvest.

DO NOT feed straw to livestock.

DO NOT apply within 14 days of harvest.

DO NOT feed straw to livestock.

DO NOT apply within 14 days of harvest.

DO NOT feed straw to livestock.

DO NOT apply within 14 days of harvest.

DO NOT feed straw to livestock.

DO NOT apply within 14 days of harvest.

IMPORTANT NOTICE TO PURCHASER AND USER OF THIS PRODUCT — This product is manufactured, labeled and sold in accordance with state and federal laws. Adequate directions for use, care on the safe handling and use of this product are set forth on this label. However, the use and handling of this product is always attended with risk of personal injury or property damage if the information and directions set forth on this label. The foregoing limitation of responsibility may not be altered in any way except by written agreement signed by officers of the manufacturer. Terms and limitations expressed herein and on this label, the manufacturer refuses to consummate the sale of this product to him and the product must be returned to the manufacturer.

CROP KING CHEMICALS, Yakima, WA

EXPERIMENTAL PASTOR []



ACCEPTED
 JUN 16 1978
 Federal Insecticide,
 Fungicide and Rodenticide Act.
 under the pesticide
 No. 68258

682-58



TEDION 400

MITICIDE

ACTIVE INGREDIENTS: Tetrathion 60%

INERT INGREDIENTS: p-chlorophenyl 2, 4, 5-trichlorophenyl sulphone 40%

PROTECT FROM TEMPERATURES BELOW 32° F.
 (Product freezes but thaws unimpaired)

CAUTION
KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified. DO NOT allow to drift to adjoining food or forage crops. DO NOT graze or feed contaminated cover crop to livestock.

- Keep container closed. Leaking packages should be removed to a safe place.
- Use with adequate ventilation.
- Avoid breathing vapor.
- Avoid contact with skin.
- Shrimp and crab may be killed at application rates recommended on this label. Keep out of Tidal Marshes and Estuaries.
- Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies.
- Containers should be disposed of by punching holes in them and burying with wastes.

(See right hand panel for further cautions)

PLUMS AND PRUNES: For European red, two-spotted, and McDaniel mites, use 1 1/2 to 2 quarts. DO NOT make more than 3 applications during the fruiting period.

GRAPE: For Pacific mite, two-spotted spider mite, and Wilsonella mite use 1 quart. DO NOT make more than 3 applications during the fruiting period.

BUSHBERRIES (Blackberries, Boysenberries, Blueberries, Gooseberries, Loganberries, Raspberries): For Two-spotted mites, apply 1 quart after harvest when mites first appear. Repeat as required.

STRAWBERRY: For Two-spotted mite use 2 pints. DO NOT apply at intervals of less than 35 days during fruiting, or within 3 days of picking.

CUCUMBERS: For Two-spotted mites, apply 1/2 pint per 100 gallons of water for greenhouse cucumbers and 1 pint per acre for field cucumbers. Apply when mites first appear and repeat as required. DO NOT make more than 3 applications during the growing season.

TOMATO: For Tropical and Two-spotted mites use 1 pint. May be used up to day of harvest, but DO NOT make more than 3 applications during fruiting period.

MELONS, PUMPKIN, WINTER SQUASH: Two-spotted mite, apply 1 pint. May be applied up to the day of harvest.

ORNAMENTALS

(Greenhouse and out-of-doors): For Two-spotted spider mite and citrus red mite use 1/2 pint per 100 gallons of water. Spray thoroughly when mites first appear and repeat as required. Application may cause leaf drop in White Butterfly and Cinderella Roses.

EVERGREENS: For Spruce mite, apply 1/2 pint per 100 gallons of water when mites first appear and repeat as required.

NOTE: Kill of adult mites is not fast and it may take several days before full effect is seen. However, this product has long residual action and should be effective for several weeks.

THOROUGHLY MIX BEFORE USE, OR USE ENTIRE CONTAINER AT ONE TIME.

NET CONTENTS

EPA Reg. No. 682-58

This product is manufactured, labeled and sold in accordance with state and federal laws. Adequate directions for use, necessary warnings and cautions, and other essential information are on the label. However, the use and handling of this product is always attended with risk of personal injury or property damage if the user or handler departs in any degree from the instructions on the label, and assumes no responsibility whatsoever for the effectiveness or safety of this product except to the extent that the product is used and handled in strict accordance with the instructions on the label. No liability or responsibility may not be altered in any way except by written agreement signed by officers of the manufacturer. Unless the intended purchaser agrees to be bound by the manufacturer's terms of sale, the manufacturer reserves the right to refuse to consummate the sale of this product to him and the product must be returned to the manufacturer.

CROP KING CHEMICALS, Yakima, Washington

