



MALATHION 8-E

For control of aphids, including the Yellow Clover aphid, alfalfa leafhopper, red spider mite, grasshoppers and alfalfa weevil larvae — use one and one-quarter pints per acre in sufficient water volume for thorough coverage. Begin application at first sign of infestation and repeat as needed. If used for control of alfalfa weevil larvae, application may be made not later than 7 days prior to harvest. CAUTION: May cause injury to stems and portions of leaves with wet spray.

ALFALFA AND CLOVER: For control of aphids, including the Yellow Clover aphid, alfalfa leafhopper, red spider mite, grasshoppers and alfalfa weevil larvae — use one and one-quarter pints per acre in sufficient water volume for thorough coverage. Begin application at first sign of infestation and repeat as needed. If used for control of alfalfa weevil larvae, application may be made not later than 7 days prior to harvest.

PEACHES: Mexican Leaf beetle, leaf hoppers, mites, use one pint per acre in sufficient water for thorough coverage. Last application 3 days before harvest.

CORN: Corn earworm, Dusky Sapbeetles, use one pint per acre applied as a dilute spray. Last application 5 days prior to harvest.

GRAPES: For control of Leafhoppers, Mealybugs, Spider mites, use one and one-half pints per acre in sufficient water for thorough coverage. Last application 3 days prior to harvest. CAUTION: May cause injury to foliage of Ribier, Almeria, Cardinal and Italian varieties.

PODS: Aphids, Army worms, corn earworms, spider mites, use eight-tenths pint per acre in sufficient water for thorough coverage. Repeat as necessary for control. Last application not later than 7 days prior to harvest.

PEACHES: Codling Moth, Plum Curculio, Redbanded leaf miner, use same spray schedule as for peaches. Mealy bug, pear psylla, spider mites, use 1 1/2 pints per acre in sufficient water in a volume spray. CAUTION: May cause injury to some peach varieties. Last application 7 days prior to harvest.

For control of aphids, including the Yellow Clover aphid, alfalfa leafhopper, red spider mite, grasshoppers and alfalfa weevil larvae — use one and one-quarter pints per acre in sufficient water volume for thorough coverage. Begin application at first sign of infestation and repeat as needed. If used for control of alfalfa weevil larvae, application may be made not later than 7 days prior to harvest. CAUTION: May cause injury to stems and portions of leaves with wet spray.

CAUTION

KEEP OUT OF REACH OF CHILDREN

FOR ADDITIONAL CAUTION INFORMATION SEE OTHER PANEL

Harmful if inhaled, swallowed, or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after using. Keep out of reach of children and domestic animals. Avoid contamination of feed and foodstuffs.

NOTE TO PHYSICIANS: Malathion, upon repeated, prolonged careless use, may cause cholinesterase inhibition. Atropine is antidotal.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

Thoroughly mix before use. Use entire container by spraying. Rinse spray equipment and containers and dispose of rinse water in a ditch or non-crop lands located away from water supplies. Do not use in conditions of rain or high humidity.

Malathion

8-E

MALATHION 8-E



Active Ingredients:

Malathion 8-E (80% Malathion)
Inert Ingredients (20%)

Kylene Range Aromatic Hydrocarbon Solvent

INERT INGREDIENTS

5.7
4.6
15.2%
100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

FOR ADDITIONAL CAUTION INFORMATION SEE OTHER PANEL

harmful if inhaled, swallowed, or absorbed through the skin. Avoid breathing
by mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap
l water after using. Keep out of reach of children and domestic animals. Avoid
contamination of feed and foodstuffs.

NOTE TO PHYSICIANS: Malathion, upon repeated, prolonged careless use, may
cause cholinesterase inhibition. Atropine is antidotal.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

Shake well before use or use entire container at one time.

Use spray equipment and containers and dispose of liquid wastes in a pit in
open lands located away from water supplies or return empty containers to
original conditioner or manufacturer.

NET CONTENTS

U.S. PAT. No. 682,550

APPLES: For control of fruit rots, use one and one-quarter pints per acre in sufficient water for thorough coverage. Repeat as needed for control. Last application 3 days prior to harvest on Green Onion. No time limitation on dry onions.

ONIONS: Onion thrip use one and one-quarter pints per acre in sufficient water for thorough coverage. Repeat as needed for control. Last application 3 days prior to harvest on Green Onion. No time limitation on dry onions.

PEAS: Pea aphid, use two (2) pints per acre in sufficient water for thorough coverage. Last application no closer than 3 days prior to harvest.

POTATOES: Aphids — use 2 pints per acre in sufficient water for good coverage. No time limitation.

PLUMS AND PRUNES: Prune louse, aphids, leafhoppers, spider mites — use one pint per 100 gallons of water in full coverage spray (600-800 gallons per acre). Last application 3 days prior to harvest.

TOMATOES: Aphids, spider mites, use one to two pints per acre in sufficient water for good coverage. Last application one day before harvest.

PEPPERMINT AND SPEARMINT: For control of aphids, arm worm, alfalfa loopers and to assist in the control of mites. Use 3 1/2 pint in sufficient water volume for thorough coverage. Repeat application as necessary. Last application 7 days before harvest.

Do not apply to any crop in bloom when bees or other pollinators are active. Avoid drift to adjoining crops about to be harvested or crops in bloom. Consult local agricultural extension service personnel for additional information. For airplane application use 10 gallons of water per acre; where equipment and economy permit use 20 gallons of water per acre. When using boom sprayers on low crops, use a minimum of 50 gallons of water per acre.

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply to any area not specified on this label.