ACCEPTED

April 1, 1975

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 682-67

Crop King

COLLOIDAL COPPER SPRAY 150

CAUTION

Avoid contact with skin. In case of skin contact, wash with soap and water. Do not breathe spray mist. Harmful if swallowed.

Avoid contamination of feed and foodstuffs. Do not store with food. Do not store with feed or forage to be fed to dairy animals or to animals being finished for slaughter.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not allow drift onto adjoining areas. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Mix thoroughly before use or use entire container at one time. Do not combine with colloidal suspensions, wettable powder or emulsifiable concentrates unless previous experience assures compatibility. Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

ACTIVE INGREDIENT:	
Metallic Copper Equivalent*	4.15%
INERT INGREDIENTS	95.85%
	100.00%

*From basic copper sulfate

CAUTION

KEEP OUT OF REACH OF CHILDREN

DIRECTIONS FOR USE

General Classification — Available to the public.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not reuse container. Return to drum reconditioner, or destroy by perforating or crushing and bury in a safe place.

APRICOTS and NECTARINES: For C
8-16 gals. per acre in a minimum o
to swell bud stage.

CHERRIES: For Coryneum Blight a
— use 10-16 gals. per acre in a mini
prior to swell bud stage.

PEACHES: For Coryneum Blight sup
in a minimum of 400 gallons of wat

BARTLETT PEARS: For Fire Blight
gallons of water starting at full bloo
until all bloom is gone. Do not sp
russeting. Do not use until polle
suppress pollenation

NET CONTENTS