

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE BACK OF PANEL FOR ADDITIONAL CAUTIONS.

U.S.D.A. Reg. No. 690-33

FOR RATS TO DRINK

NET WEIGHT 1/3 OZ.

PERK'S NEW

D - B

DOUBLE BARRELED ATTACK

RAT KILLER

MOUSE KILLER

Mix Contents of This Package With
1 QT. OF WATER

D-B Water Soluble Warfarin

A concentrate of Sodium Salt of
Warfarin for use against rats and mice

Active Ingredients:

Sodium Salt of Warfarin (Sodium Salt of 3 (4-acetonylbenzyl)-

4-hydroxycoumarin) 0.54%

Inert Ingredients: (Sugar) 99.46%

Total 100.00%

Manufactured under license from Wisconsin Alumni Research
Foundation — Patent applied for.
Equivalent 0.5% Warfarin

DIRECTIONS

DIRECTIONS FOR USE OF WATER SOLUBLE WARFARIN:

One-third ounce of Water Soluble Warfarin will make one quart of solution. Drinking water in a Mason Jar can be used. No mixing or stirring required. Do not attempt to dissolve small silica particles which remain on the bottom of the jar. Place a chicken water fount over the mouth of the jar and turn over quickly. Set fount on a clean level surface where rats are known to feed. Discard at once water in which mold or filth is noted. Keep fount clean and well supplied at all times. Do not place fount where it can be reached by children or pets and domestic animals.

ACCERTED

Jun 10, 1968

UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
AND UNDER NO. 2417-22

with comments

DIRECTIONS

DIRECTIONS FOR USE OF WATER SOLUBLE WARFARIN:

One-third ounce of Water Soluble Warfarin will make one quart of solution. Drinking water in a Mason Jar can be used. No mixing or stirring is required. Do not attempt to dissolve small silica particles which remain on the bottom of the jar. Place a chicken water fount over the mouth of the jar and turn over quickly. Set fount on a clean level surface where rats are accustomed to feed. Discard at once water in which mold or filth is noted. Keep fount clean and well supplied at all times. Do not place fount where freezing may occur. Protect founts from children or pets and domestic animals.

Where a continuous source of infestation prevails from nearby dumps or fields, permanent bait stations should be used and baits replenished as needed.

MICE:

For controlling mice follow the same general directions as for rats except bait placements may be smaller and more placements should be made. Mice are more difficult to control than rats and complete control may take a longer period of baiting.

IMPORTANT:

Baiting should continue until complete lack of feeding is noted. This should be from 5 to 14 days.

We recommend that dry warfarin bait be used in conjunction with this product, especially where there is an ample supply of fresh water.

CAUTION:

D-B Water Soluble Warfarin contains as its active ingredient an anti-coagulant chemical which if taken accidentally by humans, domestic animals, or pets may reduce the clotting ability of the blood and serious hemorrhage may result. In case baits are accidentally swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS:

When a human has been known to have accidentally ingested Water Soluble Warfarin, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol. (R)

PERK PRODUCTS CO.

1412 PICKERSON RD.

NASHVILLE, TENN.