

Perkerson's

# BLAM

# Warfarin

## RODENTICIDE

## KILLS RATS and MICE

### Anti-coagulant for Rodent Control

## GUARANTEED

### INGREDIENTS:

Active ingredient	Warfarin 3-(3-acetylbenzyl)-4-hydroxycoumarin)	0.5%
Inert ingredient	Starch	99.5%
	<b>Perk Products Company</b>	<b>100.0%</b>

**Perk Products Company**

1338 LEWIS STREET

MEMPHIS, TENNESSEE - 37210

### DIRECTIONS

- (a) Thoroughly mix one part (by weight) of rodenticide with nineteen parts of dry cereal bait
- (b) Place bait in locations where rats and mice feed, water or travel.
- (c) Bait placements along walls, in corners, and in out-cast locations where rats and mice feed are desirable.
- (d) Place baits in locations not accessible to children, pets, and domestic animals, or in tamper proof bait boxes.

#### For Rat Control:

- (a) Place at least four ounces of bait in each bait box.
- (b) Maintain an uninterrupted supply of bait in bait boxes for approximately 10 days.

#### For Mouse Control:

- (a) Place tablespoon amount of bait in each bait box.
- (b) Maintain an uninterrupted supply of bait in bait boxes for 10 days.

#### Note:

If bait molds, decomposes, or becomes stale, replace with fresh materials.

Keep away from humans, domestic animals, and birds. If ingested or inhaled by humans, domestic animals, or pets, the drug may reduce the clotting ability of the blood and cause hemorrhage. In such case, the intravenous and oral administration of Vitamin K combined with blood transfusion are indicated in the treatment of hemorrhage caused by overdoses or fishhook poisoning.

U.S.D.A. Reg. No.

ACCEPTED WITH COMMENTS

**RODENTICIDE**  
**KILLS RATS and MICE**  
**Anti-coagulant for Rodent Control**  
**GUARANTEED**

**INGREDIENTS:**

Active ingredient	Warfarin (3 -- (a-acetylbenzyl) —4 -- hydroxycoumarin)	0.5%
Inert ingredient	Starch	99.5%
	<b>Perk Products Company</b>	100.0%

1338 LEWIS STREET

MEMPHIS TENNESSEE - 37210

(d) Place baits in locations not accessible to domestic animals, or in tam...

**For Rat Control:**

- (a) Place at least four ounces of
- (b) Maintain an uninterrupted... for approximately 10 days

**For Mouse Control:**

- (a) Place tablespoon amounts of
- (b) Maintain an uninterrupted... days.

**Note:**

If bait molds, decomposes, use fresh materials.

Keep away from humans. Administered by humans, domestic animals, reduce the clotting ability of the blood. In such case, the intravenous and oral administration of K combined with blood transfusion may be necessary to control hemorrhage caused by over...

U.S.D.A. Reg

**A**  
**rin**

**E**  
**MICE**  
**nt Control**

### DIRECTIONS

- (a) Thoroughly mix one part (by weight) of rodenticide with nineteen parts of dry cereal bait
- (b) Place bait in locations where rats and mice feed, water or travel.
- (c) Bait placements along walls, in corners, and in concealed locations where rats and mice feed are desirable
- (d) Place baits in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes

#### For Rat Control:

- (a) Place at least four ounces of bait at each location
- (b) Maintain an uninterrupted supply of bait at all locations for approximately 10 days

#### For Mouse Control:

- (a) Place tablespoons amount of bait at each location
- (b) Maintain an uninterrupted supply of bait at all locations for 10 days.

#### Note:

If bait molds, decomposes, or becomes stale, replace with fresh materials.

Keep away from humans, domestic animals, and pets. Avoid ingestion followed by humans, domestic animals, or pets. The active ingredient reduces the clotting ability of the blood and causes internal hemorrhage. In such case, the intravenous and oral administration of Vitamin K combined with blood transfusion are indicated as a means of control of hemorrhage caused by overdoses of bishydroxycoumarin.

enzyl	0.5%
oumarin)	
	99.5%
	100.0%