

ACCEPTED

OCT 1 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 690-51

WARNING: Chlordane is toxic and can be absorbed through skin. Harmful or fatal if swallowed. Avoid inhaling spray mist. In case of spillage on skin, wash with soap and water. If swallowed drink emetic of mustard and call physician. Toxic to fish & wildlife. Do not contaminate any body of water nor apply to any area not specified on the label. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated.

WARNING: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse container. Destroy when empty.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS

LAWN AND GOLF GREEN INSECTS: Mix the indicated amount of Chlordane EC-8 with 5 gallons of water and spray or sprinkle on 1000 square feet to control the insects listed. **Armyworm, Fall Armyworm, Crickets, Chiggers, Ticks, Ants and Grasshoppers** - use 1 1/2 tablespoons. Do not sprinkle for 48 hours. **Mole Crickets** - sprinkle thoroughly before application. Apply only 1 1/2 tablespoons. **Earwings** - use 1 1/2 tablespoons. Spray around **foundations, along fences, around refuse, trash and other possible hiding places.** **Cutworms** - use 3 tablespoons and sprinkle thoroughly after application. **Wireworms** - use 2 tablespoons on sandy soil, 4 on clay and loams and 6 on peat or mucks. Water thoroughly after application. **Sod Webworm and lawn moths** -

PERKERSON'S (BLAM)

CHLORDANE EC-8 EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENTS:

*Technical Chlordane 72.8%
Deodorized Kerosene 22.0%

INERT INGREDIENTS: 5.2%
E. P. A. REG. NO. 690-51 TOTAL 100.0%

*Equivalent to 43.7% Octachlora-4, 7-methanotetrahydroindane and 29.1% related compounds.

WARNING: KEEP OUT OF REACH OF CHILDREN.

Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. See side panel for additional precautions.

Net contents _____

sprinkle before application. Apply 2 to 4 tablespoons, then wait 48 hours before watering again.

CRABGRASS: To control crabgrass in lawns apply this material at the rate of 8 gallons per acre (8 oz. to 350 sq. ft.). Make application in early spring before crabgrass seeds germinate. Do not use on newly seeded lawns. Keep children and pets off treated areas until this material has been thoroughly watered into the soil and the grass has been thoroughly dried. Birds feeding on treated areas may be harmed.

CUTWORMS, MOLE CRICKETS: On **beets, celery, collards, peppers, potatoes** and in flower gardens apply 1 to 2 pmt per acre in sufficient water for coverage. Apply to soil around the base of plants. Do not apply after edible parts of plants begin to form.

WHITE GRUBS, JAPANESE LARVAE, WHITE-FRINGED BEETLE LARVAE: On **grasslands, golf greens, lawns, flower gardens.** Apply at a rate of 5 to 10 pints per acre (1/4 to 1/2 cup in 5 gallons of water applied to 1000 square feet) in sufficient water for coverage. On lawns and golf greens water thoroughly after application. Keep children and pets off treated area until grass is dry.

NOTICE

Perk Products Company makes no warranty, expressed or implied, concerning this material, except that it conforms to the chemical description on the label. Neither Perk Products Company nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.