PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.

CLAIRE

MULTI-USE

INSECTICIDE

SPRAY

Contains the Synthetic Pyrethroid Permanone

Controls Residential, Industrial and Garden Insects

For Use On Dogs and on Premises to Control Fleas and Ticks

Aqueous, Stable

Controls Pests for Up to Four Weeks

ACTIVE INGREDIENT:

#Permethrin INERT INGREDIENTS:

4(3-phenoxyphenyl) methyl (+) cis/trans 3-(2,2dichlorgethenyl; 2,2-dimethylcyclopropanecarboxylate. Cis/Trans ratio: Min. 35% (+) cis and max. / 65% (+) trans.

BEST AVAILABLE COPY

Keep Out of Reach of Children

CAUTION

See back panel for additional precautionary statements.

Recommended for Use by Commercial Applicators

NET CONTENTS __ GZ. With COMMENTS AUG 23 1990 -

PERMANONE - Registered Trademark of Fairfield Unite the for an investigation of the president of the pr American Corporation

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply directly to food. Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms, but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Non-food areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mon closets and storage (after canning or bottling).

Not for use in USDA Meat and Poultry Plants. Do not use this product in or on velectrical equipment due to the possibility of shock hazard.

SHAKE WELL BEFORE USING

Houseflies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Blow Flies and Small Flying Moths: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Repeat treatment as necessary.

Cockroaches, Materbugs, Palmetto Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sombugs, and Pillbugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moths, Angoumois Grain Moths: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application as necessary.

Exposed states of <u>Confused Flour Beetles</u>, <u>Saw-Toothed Grain Beetles</u>, <u>Red Flour Beetles</u>, <u>Rice Weevils</u>, <u>Granary Weevils</u>, <u>Grain Mites</u>: Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these inserts tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as pacessary. I fested stored food products in the home should be disposed of.

<u>Carpet Beetles:</u> Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

<u>Bedbugs:</u> Spray mattresses, particularly around tufts and seass. Take beds apart and spray into all joints. Allow mattresses and other articles to dry thoroughly before use. Treat baseboards, moldings and floors. Repeat treatment as necessary.

BEST AVAILABLE COPY

To control <u>Clothes Hoths:</u> Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

IN ANIMAL QUARTERS AND MILK ROOMS: For rapid control of <u>Houseflies</u>, <u>Mosquitoes</u>, <u>Gnats</u>, <u>Wasps</u>, <u>Hornets</u>, <u>Bees</u> and <u>Small Flying Moths</u>, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate before reoccupying.

TO TREAT LIVESTOCK: To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of <u>Stable Flies</u>. Horse <u>Flies</u>, <u>Deer Flies</u>, <u>Face Flies</u>, <u>House Flies</u>, <u>Horn Flies</u>, <u>Mosquitoes</u>, and <u>Gnats</u>: Spray about 3 seconds on each side being careful to spray back, withers and forelegs thorough; y. To protect from face flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome. For Swine: Do not ship animals for slaughter within 5 days of last treatment.

To control <u>Blood Sucking Lice:</u> Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control <u>Poultry Lice</u>, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of <u>Bedbugs</u> and <u>Mites</u> in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

Fleas and Ticks: (Adults and Larvae): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before general use.

To control <u>Fleas</u>, <u>Ticks</u> and <u>Lice</u> on <u>Dogs</u>: Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Repeat application weekly if necessary to control infestations. NOTE: Not recommended for use on pupples (Sess than four weeks old.

For the Control of Garden Insects: For use on roses, dahlias, astems and other ornamentals to kill <u>Japanese Beetles</u>, <u>Whiteflies</u>, <u>Aphids</u>, <u>exposed Thrips</u>, <u>Spuders</u>, <u>Red Mites</u> and <u>Leafminers</u>, hit by spray.

May be used on plants, as well as a multi-purpose spray in small gardens for spot treatments of incipient insect infestations to prevent spreading in large garden as was:

May also be used on other plants such as African violets, asters, azaleas,...'begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds. rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.



Effective when used as directed against Whiteflies, Aphids, Spider and Red exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Morms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat as infestations occur.

STORAGE: Store in a cool, dry place (preferably locked) inaccessible to children and pets and away from direct sunlight, windows, flames and other sources of heat or ignition. Do not transport or store below 32°F. DISPOSAL: Replace cap, wrap container in several layers of newspaper and discard in trash. Do not incinerate or puncture.

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid spraying in eyes. Avoid breathing spray mist. Avoid contact with skin. Wash hands thoroughly after use. Vacate room after treatment and ventilate before reoccupying. Avoid contamination of feed and foodstuffs. Cover or remove fishbowls. No not allow children or pets to contact treated areas until surfaces are dry. Wear full-faced gas mask with canister type recommended for general insecticide protection when applying indoors as a space spray. THIS PRODUCT IS NOT FOR USE ON HUMANS. If lice infestations should occur on humans, consult either a physician or pharmacist for a product for use on humans. Animals: Do not spray directly in/on eyes, mouth or genitalia. Do not cause exposure to puppies less than four weeks old.

Statement of Practical Treatment

If Inhaled: Remove affected person to fresh air. Apply artificial respiration if indicated. If In Eyes: Flush with plenty of water. Contact a physician if irritation persists. If On Skin: Immediately remove contaminated clothing and wash affected areas immediately with soap and water. Contact a physician if irritation persists. Wash clothing prior to reuse. If Swallowed: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

24 HOUR EMERGENCY NUMBER --- 1-800-228-5635

Physical or Chemical Hazards

Contents under pressure. Do not use or store near fire, sparks or heated surfaces. Do not store in direct sunlight. Do not smoke while using this product. Do not puncture or incinerate container. Exposure to temperatures above 120°F may cause bursting. Incineration of container may cause explosion.

EPA Reg. No. 706-DL

EPA Est. 706-IL-1

CLAIRE MANUFACTURING COMPANY
500 Vista Avenue, Addison, Illinois 60101

Buver assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

BEST AVAILABLE GURY