

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. May cause skin irritation. Avoid contact with skin and eyes.

FRUITIVE CLOTHING consisting of long pants, long sleeve shirt, gloves, hat & boots must be worn during mixing and loading.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with large amounts of water for at least 15 minutes. Call a physician if irritation persists.

IF INHALED: Move subject to fresh air.

IF ON SKIN: Wash affected skin areas with soap and water. Call a physician if irritation persists.

IF SWALLOWED: Institute giving 2 glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person.

Before using this product read the entire Precautionary Statement, Conditions of Sale and Warranty, Directions for Use, Use Precautions, and Storage and Disposal Instructions. If the conditions of sale and Warranty are not acceptable, return the product in perfect condition thirty days of purchase to the place of purchase.

REENTRY AND REENTRY PROTECTION STATEMENTS

Do not reenter treated areas until protective clothing and sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in a area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

CAUTION

Closest area to field description treated with DITHIOM 7073. Do not enter treated areas until protective clothing and sprays have dried. In case of accidental exposure call a doctor, physician, hospital or hospital immediately. Explain that the victim has been exposed to DITHIOM 7073 and describe his condition. For further information see the Precautionary Statement of Practical Treatment.

Oral warnings must include the same information as is in the written warnings.

STORAGE AND DISPOSAL

For safe disposal, do not contaminate water, soil, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environment Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

NEVER CONTAMINATE FEEDSTUFFS, FODDER OR ANY OTHER FEEDS. Do not use or store this product in a way that will contaminate feedstuffs, or other products approved by state and local authorities.

DO NOT BE TAKEN IN CASE MATERIAL IS HELD IN CONTAINERS like and can spray with inert material (sand, earth, filler's earth, etc.) or packed liquid and solid diving material to separate containers for re-use or disposal. Remove contaminated clothing and wash affected skin areas with soap and water. Wash clothing before reuse. Keep sprays out of streams and other bodies of water. Refer to Precautionary Statements.

ACCEPTED

JUL 17 1985

Under this contract, the buyer
For your information, the
On 7/17/85, the following
Registered under
EPA Reg. No. 7073

ACTIVE INGREDIENT

Nabam (Disodium ethylene
bisdithiocarbamate) 22%
INERT INGREDIENTS 78%
TOTAL 100%


2 lbs. active per gallon

EPA Reg No 7073

EPA Est No 9386 GA 1

NET CONTENTS

BEST AVAILABLE COPY GALLONS



**ROHM
AND
HAAS**

PHILADELPHIA, PA. 19105

Further details on mixing, compatibility, and for additional crops see the recommendations on this label and in special circulars. Then consult the current State Agricultural Experiment Station spray schedules or State Extension Service Specialists for details on dosage, timing and frequency of sprays.

Thorough, uniform and timely coverage is absolutely essential for disease control. Either standard high volume, concentrate or aerial sprayers may be used. Use 2 qts. of Dithane D-14 plus metallic sulfate per 100 gallons of water in standard sprayers, or the equivalent in concentrate machines or aerial sprayers. Apply 2 to 3 qts. of Dithane D-14 per acre to vegetable and field crops.

■ MIXING DIRECTIONS

For each 2 quarts of DITHANE D-14, use one of the following:
Zinc sulfate (36% zinc) $\frac{3}{4}$ pound
Zinc sulfate (22.8% zinc) $1\frac{1}{4}$ pound
Steps to follow—

1. Fill tank half to almost full with water.
2. Add required zinc and dissolve thoroughly.
3. Add required DITHANE D-14. Agitate thoroughly.
4. Add insecticide, spreader, or nutritional sprays last if used.

VEGETABLES, FIELD CROPS, ORNAMENTALS. Apply Dithane D-14 plus zinc sulfate sprays on a regular 7 to 10 day schedule, or oftener under severe disease conditions. Begin applications when disease is first present and continue as long as disease conditions prevail. To avoid excessive residues observe the following last application dates:

5 days for: Cantaloupes, Cucumbers, Tomatoes, and Watermelons.

7 days for: Beets, Broccoli, Cabbage, Carrots, Cauliflower and Turnips.

10 days for: Kale, Lettuce, Mustard Greens, Swiss Chard.

14 days for: Celery (remove residue remaining at harvest by clipping, washing or trimming) and hops.

USE DITHANE D-14 PLUS ZINC SULFATE TO CONTROL THE FOLLOWING DISEASES:

ASPARAGUS—Rust. Apply to new fern growth after harvest or to new plantings at first sign of disease. Repeat at 10-day intervals until fern growth is completed.

BEANS (lima and snap)—Downy mildew, Rust, Anthracnose (Begin after first blossoms)

BEETS, KALE, MUSTARD GREENS, SWISS CHARD, TURNIPS—Cercospora leaf spot, Downy mildew

CABBAGE, BROCCOLI, CAULIFLOWER—Downy mildew, Alternaria. Use in plant bed and field

CANTALOUPE, CUCUMBERS, WATERMELONS—Downy mildew, Alternaria (Macrosporium) leaf spot, Anthracnose, Gummy stem blight. Direct sprays especially to underside of leaves.

CARROTS—Alternaria and Cercospora leaf spot.

LETTUCE—Downy mildew

ONIONS—Downy mildew, Botrytis leaf blight, Purple blotch. Apply as directed above. Onion Smut, apply one gallon Dithane D-14 per acre without zinc sulfate as a furrow drench at planting. Dilute Dithane D-14 as necessary in sufficient water for adequate application in available equipment

SPINACH—Downy mildew, White rust

TOMATOES—Early and late blight, Anthracnose, Stemphylium and Septoria leaf spots, Cladosporium gray mold

AZALEAS and CAMELIAS—Petal blight. Use only 1 qt. of DITHANE D-14 plus 6 ozs. of zinc sulfate (36% zinc) per 100 gallons. Spray 2 or 3 times a week while flowers are opening. Apply also to soil and litter around plants while they are in bloom

CHRYSANTHEMUMS, CARNATIONS, SNAPDRAGONS—Rust, Leaf spot, Anthracnose.

GLADIOLUS—Curvularia, Botrytis blight, Stemphylium.

FRUITS

Apply sprays thoroughly in accordance with timings recommended by State Agricultural Experiment Station spray schedule for the specific crops. Application may be made either by suitable ground or aerial equipment. To avoid residues in excess of tolerances do not apply within 7 days of harvest, except as otherwise noted

USE DITHANE D-14 PLUS ZINC SULFATE FOR THE FOLLOWING DISEASES:

GRAPES—Black rot, Downy mildew, Ripe rot.

STRAWBERRIES—Leaf spot, Leaf scorch

SOUR CHERRIES—Leaf spot. Use only 1 qt. DITHANE D-14 plus 6 ozs. of zinc sulfate (36% zinc)

DITHANE D-14 contains ingredients which provide adequate wetting on many crops. On certain hard-to-wet crops, the addition of 2 to 6 ounces of Triton® B-1956® per 100 gallons of dilute spray may be desirable for optimum coverage of the plant surface.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Crop injury, ineffectiveness or other unintended consequences resulting from such factors as failure to follow label directions, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property, handling, storage and manner of use are beyond the control of Rohm and Haas and Seller. All such risks will be assumed by the Buyer or User. In no case will Rohm and Haas or Seller be held liable for consequential, special or indirect damages resulting from the use or handling of this product.

BEST AVAILABLE COPY