

DIRECTIONS FOR USE

General: KARATHANE is a fungicide and miticide. It is effective against a wide range of fungi and mites on many crops.

powdery Mildew Control: KARATHANE Liquid Concentrate is a systemic fungicide. It is effective against powdery mildew on many crops. Apply KARATHANE Liquid Concentrate to the foliage of the crop at the first appearance of the disease. Repeat applications at 14-day intervals until the disease is under control.

Mite Control: KARATHANE Liquid Concentrate is a systemic miticide. It is effective against a wide range of mites on many crops. Apply KARATHANE Liquid Concentrate to the foliage of the crop at the first appearance of the mites. Repeat applications at 14-day intervals until the mites are under control. KARATHANE Liquid Concentrate may also be used in concentrate ground or aerial sprayers. Be sure to adjust dosage accordingly.


Directions are given for single strength sprays. KARATHANE Liquid Concentrate may also be used in concentrate ground or aerial sprayers. Be sure to adjust dosage accordingly.

APPLES, PEARS: To control powdery mildew use 4 to 6 ounces KARATHANE Liquid Concentrate to 100 gallons water, applying through full coverage spray at 500-800 gallons per acre for dilute sprays. Adjust for spray for concentrate sprayers. Begin applications at development of pink stage and continue in every regular spray at 14-day intervals until the cover sprays until terminal growth or mildew activity is completed. At no time should applications of KARATHANE be more than 14 days apart. Thorough coverage is necessary for mildew control. When not visible mildew growth is evident, add a spreader/shaker to the spray to insure thorough wetting of the leaves and fruit. Certain stick spreaders will reduce the deposit of KARATHANE on apples and pears. If a stick spreader is to be employed with KARATHANE Liquid Concentrate, use TRITON® B-1956 at 1 to 2 ounces per 100 gallons of dilute spray.

KARATHANE is a systemic fungicide and miticide. It is effective against a wide range of fungi and mites on many crops. Apply KARATHANE Liquid Concentrate to the foliage of the crop at the first appearance of the disease. Repeat applications at 14-day intervals until the disease is under control.

KARATHANE is a systemic fungicide and miticide. It is effective against a wide range of fungi and mites on many crops. Apply KARATHANE Liquid Concentrate to the foliage of the crop at the first appearance of the disease. Repeat applications at 14-day intervals until the disease is under control.

KARATHANE® LIQUID CONCENTRATE



MMS

PHILADELPHIA, PA 19105

AGRICULTURAL FUNGICIDE AND MITICIDE

CONTAINS 4 LBS. TECHNICAL KARATHANE® PER GALLON

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin or eyes. May cause skin irritation or sensitization. Wash thoroughly after use. Avoid inhalation of mist. Harmful if swallowed. Avoid contamination of food stuffs and feeds. Inflammable—Keep away from flames.

Do not apply when weather conditions favor drift. Spray from ground. Treated foliage, equipment and containers should be disposed of waste by burying in non-crop lands away from water supplies.

This product is for use only for the purposes for which it is registered. Keep out of any body of water.

USDA Registered under 240.241
EPA Reg. No. 240.241

ACTIVE INGREDIENTS


Technical Karathane 4.00
Fungicide 1.00
Miticide 1.00
Inert Ingredients 1.00

INERT INGREDIENTS

Water 1.00
Fungicide 1.00
Miticide 1.00
Inert Ingredients 1.00

NET CONTENTS

1.00 GALLONS



DIRECTIONS FOR USE

General: KARATHANE Liquid Concentrate is a highly effective fungicide and miticide for use on a wide variety of crops.

Powdery Mildew Control: KARATHANE Liquid Concentrate is a systemic fungicide that controls powdery mildew on a wide variety of crops. It is recommended for use on crops with various degrees of resistance to powdery mildew. The following directions are applicable to your area.

Mite Control: KARATHANE Liquid Concentrate is an effective miticide for control of mites on a wide variety of crops. It is recommended for use on crops with various degrees of resistance to mites. The following directions are applicable to your area.

Directions are given for single strength sprays. KARATHANE Liquid Concentrate may also be used in concentrate ground or aerial sprayers. Be sure to adjust dosage accordingly.

APPLES, PEARS: To control powdery mildew use 4 to 6 ounces KARATHANE Liquid Concentrate to 100 gallons water applying thorough full coverage spray at 500-800 gallons per acre for dilute sprays. Adjust accordingly for concentrate sprayers. Begin applications at delayed dormant to pink stage and continue in every regular spray at 7-14 day intervals until the cover sprays until terminal growth or mildew activity is completed. At no time should applications of KARATHANE be more than 14 days apart. Thorough coverage is necessary for mildew control. When powdery mildew growth is evident add a spreader sticker to the spray to insure thorough wetting of mildew spores and mycelium. Certain sticker spreaders will reduce the deposit of KARATHANE on apples and pears. If a sticker spreader is to be employed with KARATHANE Liquid Concentrate use TRITON® B-1956 at 1 to 2 ounces per 100 gallons of dilute spray.

KARATHANE Liquid Concentrate is also effective against various other diseases of apples and pears. For complete directions on the use of KARATHANE Liquid Concentrate on apples and pears see the full directions on the label.



KARATHANE Liquid Concentrate is also effective against various other diseases of apples and pears. For complete directions on the use of KARATHANE Liquid Concentrate on apples and pears see the full directions on the label.

KARATHANE Liquid Concentrate is also effective against various other diseases of apples and pears. For complete directions on the use of KARATHANE Liquid Concentrate on apples and pears see the full directions on the label.

TRUS

PEACHES AND APRICOTS

For complete directions on the use of KARATHANE Liquid Concentrate on peaches and apricots see the full directions on the label.



**ROHM
AND
HAAS**
PHILADELPHIA, PA. 19105

**AGRICULTURAL FUNGICIDE
AND MITICIDE**
CONTAINS 4 LBS. TECHNICAL KARATHANE® PER GALLON

**CAUTION: KEEP
OUT OF REACH OF CHILDREN.** Avoid contact with skin or eyes. May cause skin irritation or sensitization. Wash thoroughly after use. Avoid inhalation of mist. Harmful if swallowed. Avoid contamination of food, feed, and feedstuffs. Inflammable—Keep away from flames.

Do not apply when weather conditions favor drift of spray from areas treated. Rinse equipment and containers and dispose of waste by burying in non-crop lands away from water supply.

This product is for use on crops. Keep out of any body of water. EPA Reg. No. 100-100000-01

ACTIVE INGREDIENTS

Technical Karathane 4.00 lbs. per gallon

INERT INGREDIENTS

Water 96.00 lbs. per gallon

NET CONTENTS

1.00 GALLONS

DIRECTIONS

RASPBERRIES: KARATHANE Liquid Concentrate is a systemic fungicide that controls powdery mildew on a wide variety of crops. It is recommended for use on crops with various degrees of resistance to powdery mildew. The following directions are applicable to your area.

STRAWBERRIES: KARATHANE Liquid Concentrate is a systemic fungicide that controls powdery mildew on a wide variety of crops. It is recommended for use on crops with various degrees of resistance to powdery mildew. The following directions are applicable to your area.

GOOSEBERRIES: KARATHANE Liquid Concentrate is a systemic fungicide that controls powdery mildew on a wide variety of crops. It is recommended for use on crops with various degrees of resistance to powdery mildew. The following directions are applicable to your area.

GRAPES: KARATHANE Liquid Concentrate is a systemic fungicide that controls powdery mildew on a wide variety of crops. It is recommended for use on crops with various degrees of resistance to powdery mildew. The following directions are applicable to your area.


CANTALOUPE, WATERMELONS: KARATHANE Liquid Concentrate is a systemic fungicide that controls powdery mildew on a wide variety of crops. It is recommended for use on crops with various degrees of resistance to powdery mildew. The following directions are applicable to your area.

ORNAMENTAL PEACHES, CHERRIES: KARATHANE Liquid Concentrate is a systemic fungicide that controls powdery mildew on a wide variety of crops. It is recommended for use on crops with various degrees of resistance to powdery mildew. The following directions are applicable to your area.

NOTICE: KARATHANE Liquid Concentrate is a systemic fungicide that controls powdery mildew on a wide variety of crops. It is recommended for use on crops with various degrees of resistance to powdery mildew. The following directions are applicable to your area.

body with a few drops of water. Concentrate on the head and thorax. Powder can be used on the legs with care to avoid application to the area of the joint. As an effective molluscicide, KARATHANE liquid is a very short residual. It may provide effective control of KARATHANE for a few days after emerging. If adult applications are infrequent, alternating with KARATHANE may be used to give the insect a constant contact of resistance.

Use 4 to 6 ounces
dilute with water applying
one quart per acre for dilute
sprays. Begin applica-
tions in early spring and
continue in every regular
spray until terminal
fruit time should applica-
tions be continued.
For thorough coverage
of the plant, use a
fine mist or spray. It is
advisable to insure thorough
coverage of the plant with
certain types of spreaders
and sprayers. If a
RATHANE Liquid Con-
centrate is used, use 1
quart per 100 gallons of



W
K-A-
a-
M-

CONTAINS 4 LBS. TECHNICAL KARATHANE® PER GALLON

Do not apply when weather conditions favor drift of spray from treated areas. Treat fish, equipment and containers, and dispose of waste by burying in non-ropic soils away from water supplies.

Don't let the opportunity
pass you by.

INERT INGREDIENTS

10. 1. 2010

For the purpose of this study, the following hypotheses were formulated:

GALLONS

A

NET CONTENTS

709-31

REMARKS:

709-31

[illegible][illegible][illegible]

GRAPES Do not use on grapes in California

CANTALOUPE, MONEYDEW MELONS, MUSKMELONS, BUMPKINS, WATERMELONS, WINTER MELONS, CHAMP OF SUMMER SQUASH
To control powdery mildew use 8-4-4 or 16-16-16 ATHANE 1-10-10 Concentrate in 100 gallons water and apply 1-10-10 1/2 oz. per acre. Begin applications when the first signs of infection are noticed. If necessary, repeat applications at 7-10 day intervals. In severe cases, if necessary, repeat and increase application rate. The amount of water used will vary according to water content of the crop and soil conditions. For best application, use the spray nozzle which gives the finest spray possible. Apply in the early morning or late afternoon. Apply to the underside of leaves.

ORNAMENTAL TREES, SHRUBS, NURSERY STOCK, APPLES,
PEACHES, CHERRIES

NOTICE