


DITHANE M-22 SPECIAL WITH ZINC



AGRICULTURAL FUNGICIDE

CAUTION KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT

INERT INGREDIENTS

EPA Reg. No. 707-85-AA

NET CONTENTS

3 LBS.



ROHM AND HAAS
PHILADELPHIA, PA. 19103

FRUITS AND NUTS

ALMONDS—Fruit rot. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications when fruit is 1/2 grown and repeat every 14 days through petal fall and up to final timing recommendations and disease severity. For fruit brown rot use again at 4 to 6 weeks before harvest, repeating applications every 7 to 14 days. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Apply only in good applications. Begin when leaves first appear and repeat every two or three weeks or as required.

APPLES—Fruit rot. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications when fruit is 1/2 grown and repeat every 14 days through petal fall and up to final timing recommendations and disease severity. For fruit brown rot use again at 4 to 6 weeks before harvest, repeating applications every 7 to 14 days. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Apply only in good applications. Begin when leaves first appear and repeat every two or three weeks or as required.

APRICOTS, NECTARINES, PEACHES—Western States. Brown rot, leaf curl. Begin at formation to petal stage, repeat every 7 to 14 days through petal fall and up to final timing recommendations and disease severity. For fruit brown rot use again at 4 to 6 weeks before harvest, repeating applications every 7 to 14 days. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Apply only in good applications. Begin when leaves first appear and repeat every two or three weeks or as required.

BANANAS—Sigatoka. Use 2 to 4 lbs. DITHANE M-22 SPECIAL per acre in sufficient water to provide adequate coverage. Apply only in good applications. Begin when leaves first appear and repeat every two or three weeks or as required.

CRANBERRIES—Fruit rot. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

ES—Fruit rot. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

FIELD CROPS

CEREAL GRAINS—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Crop	Rate	Application
Barley	1.0-2.0 lb.	tamping off seed in its seed bed
Oats	1.0-2.0 lb.	tamping off seed in its seed bed
Rye	1.0-2.0 lb.	tamping off seed in its seed bed
Wheat	1.0-2.0 lb.	tamping off seed in its seed bed

GRASSES FOR SEED—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

FIELD CROPS

SUGAR BEETS—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

TOBACCO—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

ORNAMENTALS AND TURF—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

VEGETABLES—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

FRUITING PLANTS—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

FLOWERING DOGWOOD—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

PEONY—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

CARNATION—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

DAHLIA—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

GLADIOLUS—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

LILY—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

PANSY—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

ROSE—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

SNAPDRAGON—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

ZINNIA—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

TURF GRASSES—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

CELERY—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

COLLARDS, KALE, MUSTARD GREENS, TURNIPS—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

CORN—Fungal diseases. Use 1 to 2 lbs. DITHANE M-22 SPECIAL per acre in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

DIRECTIONS FOR USE GENERAL INSTRUCTIONS

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

Apply DITHANE M-22 SPECIAL in sufficient water for good coverage. Begin applications at mid bloom and repeat at 14 day intervals up to final application within 14 days of harvest.

acre in
Beet
repeat
not free
EGGPL
Apply
Dithane
LETTU
leaf
rot
DITHANE
M-22
SPECIAL
per
acre
in
sufficient
water
for
good
coverage
Begin
applications
at
mid
bloom
and
repeat
at
14
day
intervals
up
to
final
application
within
14
days
of
harvest
ONION
purple
in
the
leaves
during
POTA
7 to 10
days
before
harvest
PUMPK
in
the
leaves
Start
and re
Do not
SPINA
rot
A
at 7 to
quate
harvest
TOMATO
blight,
leaf
sg
tomato
Begin
potato
wilt
or
ions
of
severe
Do not
NOTE
1
sticker
the
spr
to
smo
TRITICO
able
p
used
is
recol
spread
DITHANE
M-22
SPECIAL
per
acre
in
sufficient
water
for
good
coverage
Begin
applications
at
mid
bloom
and
repeat
at
14
day
intervals
up
to
final
application
within
14
days
of
harvest
NOTICE
forms
f
ably
fit
used
in
condi
any
off
for
a
l
end
produc
abnorm
reason
sumes
include
crops
l
pled
fi
drift

THANEM-22 SPECIAL WITH ZINC

AGRICULTURAL FUNGICIDE

CAUTION KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT

Methyl N-propylmaleate 80%

INERT INGREDIENTS

20%

EPA Reg. No. 707-83 AA

NET CONTENTS

3 LBS.

ACCEPTED

JAN 23 1973

FIELD CROPS

SPRING PLANTS—Apply to soil or foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

SUGAR BEETS

Apply to soil or foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

For soil application use 1 lb. of THANEM-22 per 100 lbs. seed.

Do not feed treated seed to any farm animals.

TOBACCO

Apply to soil or foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

ORNAMENTALS AND TURF

CHRYSANTHEMUM—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

FLOWERING DOGWOOD—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

PEONY—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

CARNATION—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

DAHLIA—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

GLADIOLUS—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

LILY—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

PANSY—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

ROSE—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

SNAPDRAGON—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

ZINNIA—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

TURF GRASSES—Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

DIRECTIONS FOR USE GENERAL INSTRUCTIONS

THANEM-22 is a fungicide designed for the control of various plant diseases. Apply to soil or foliage at 1 to 2 lbs. per acre. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

Apply to soil or foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

Apply to soil or foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

VEGETABLES

BEANS—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

CANTALOUPE, CUCUMBERS, MELONS, SQUASH, WATERMELONS—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

CARROTS—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

CELERY—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

COLLARDS, KALE, MUSTARDGREENS, TURNIPS—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

CORN—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

Apply to soil or foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

EGGPLANT, PEPPERS—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

LETTUCE, ENDIVE—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

ONIONS—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

POTATOES—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

PUMPKINS—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

SPINACH—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

TOMATOES—Apply to foliage every 2 to 3 weeks. Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

NOTE—THANEM-22 SPE may be used without any sticker spreader. When necessary to improve the spray coverage on foliage difficult to wet and to smooth out spray deposits on maturing fruit TRITON® B 1946® is recommended. Other suitable sticker spreaders may also be used, except in situations where TRITON B 1946 is recommended as the only acceptable sticker spreader.

THANEM-22 SPE FOR DUSTING—Apply to soil at 1 to 2 lbs. per acre. Apply to foliage at 1 to 2 lbs. per acre. Do not apply to soil or foliage of plants in bloom.

NOTICE—The user warrants that the product conforms to the label description and is reasonably fit for the purpose stated on the label when used in accordance with the directions. The user assumes the risk of injury to himself or any other person, property, or animal, or damage to crops, if the product is used in any manner not reasonably foreseeable to the seller, and buyer assumes the risk of injury to himself or any other person, property, or animal, or damage to crops, if the product is used in any manner not reasonably foreseeable to the buyer. These risks include, but are not limited to, damage to plants, crops, and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

PRINTED IN U.S.A.