

GENERAL

FORE Turf and Ornamental Fungicide is a unique and different fungicide which when used according to directions, provides very high fungicidal activity and is safe for use on both turf grasses and certain ornamentals. This new fungicide provides proven control of most common turf grass diseases and it is also effective in controlling major fungus diseases of certain ornamentals.

Diseases of turf grass and ornamentals can attack suddenly and unexpectedly causing severe damage and may even result in total loss of large areas of valuable turf grass and ornamental plants. The use of a regular protective spray program will minimize the risk of disease damage and can generally be accomplished with lower rates and less frequent fungicide applications. Once diseases have become established higher rates of fungicide and more frequent applications are required to bring them under control. Follow a regular protective program for maximum FORE Turf and Ornamental Fungicide performance.

To mix, slowly sift FORE Turf and Ornamental Fungicide into spray tank while it is being filled and the agitator is running. Add insecticide and nutritional sprays last. FORE Turf and Ornamental Fungicide is believed compatible with most commonly used pesticides.

DIRECTIONS FOR USE - TURF DISEASES

Fusarium Blight (F. roseum), Red Thread, Copper Spot, Slime Molds In a protective spray program use FORE at 4 to 6 oz. per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens-up in the spring and repeat every 7 to 14 days.

When conditions are especially favorable for disease development increase FORE Turf and Ornamental Fungicide rate to 6 to 8 oz. per 1000 sq. ft. and repeat applications every 7 days until disease threat is past.

Helminthosporium Melting Out, Rhizoctonia Brown Patch, Rust Use FORE at 4 oz. per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens-up in the spring and repeat every 7 to 14 days for Helminthosporium and Rust and every 7 days for Rhizoctonia control.

Pythium blight Use FORE at 8 oz. per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications at first sign of disease and repeat at 5 day intervals or oftener if conditions are especially favorable for disease development. Continue until disease threat is past.


Algae control in turf grass Use FORE at 6 oz. per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when algae begin to appear and repeat every 7 days until threat is past.

Follow local turf disease control recommendations for further details on timing of applications.

ACCEPTED

APR 17 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 207-8 SUBJECT
NO ATTACHED COMMENTS.



**ROHM
AND
HAAS**

PHILADELPHIA, PA. 19105

FORE[®]

A FUNGICIDE FOR TURF AND ORNAMENTALS

CAUTION: May cause irritation of nose throat, eyes and skin. Do not breathe dust or spray mist. In case of skin contact wash with water; for eyes wash copiously with water and get medical attention. Contamination of fish-bearing waters with this product may harm fish. **KEEP OUT OF REACH OF CHILDREN.**

IMPORTANT: Keep away from fire and sparks. Store in a cool, dry place. Do not allow to become wet or overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of FORE and may also generate flammable vapors. Keep container closed when not in use.

U.S. Patents 2,504,404; 2,974,156 and Patent Pending.

USDA Registration No. 707-87

ACTIVE INGREDIENT

A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate	80%
In which the ingredients are	
Manganese + +	16%
Zinc + +	2%
Ethylene bisdithiocarbamate ion (C ₄ H ₈ N ₂ S ₄) =	62%
INERT INGREDIENT	20%
	100%

**NET CONTENTS
50 LBS.**

707-87

04/17/1967

1/2

Dollar Spot (Sclerotinia) In a protective spray program use FORE at 6 to 8 oz. per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens-up in the spring and repeat every 7 to 14 days.

When conditions are especially favorable for dollar spot development use FORE at 8 oz. per 1000 sq. ft. and repeat applications every 7 days until disease threat is past. A cadmium fungicide may be added to supplement dollar spot control under severe disease conditions.

Fusarium Snow Mold Use FORE at 6 to 8 oz. per 1000 sq. ft. in sufficient water to provide adequate coverage. Make applications during the winter at a 2 to 6 week interval.

ORNAMENTALS

Use FORE at $1\frac{1}{2}$ lbs. per 100 gallons ($1\frac{1}{2}$ level tablespoons per gallon) in full coverage spray. Begin spraying when plants are well leafed-out and apply at 7 to 10 day intervals throughout the season or follow local commercial florists or State Extension Service recommendations for control of:

ROSES	Black spot
IRIS	Leaf spot
DAHLIA	Botrytis blight
PANSY	Anthrachnose
ZINNIA	Leaf blight
CARNATION	Leaf spot
LILY	Botrytis blight
SNAPDRAGON	Rust
TULIP	Botrytis blight (fire)
HOLYHOCK	Leaf spot, anthracnose, rust
HOLY	Purple spot
HONEYSUCKLE	Herpobasidium blight
ORNAMENTAL CRABAPPLE	Scab, cedar-apple rust
PEONY	Phytophthora and Botrytis blight. Apply in early spring and early fall, drenching soil around plants as well as the foliage. Destroy all infected parts promptly.
PACHYSANDRA	Volutella blight. Use a drenching spray of 2 lbs. FORE in 50 gallons of water per 5000 sq.ft. of bed area. At least 5 applications about 10 to 14 days apart are required. Start at first sign of disease.
FLOWERING DOGWOOD	Anthrachnose. Apply when buds begin to open, when bracts have fallen, 4 weeks later and again in late summer after flower buds for next season are formed.
AZALEAS, CAMELLIAS, RHODODENDRON	Petal blight. Spray 2 to 3 times a week while flowers are opening. Direct spray into flowers and spray ground under bushes thoroughly.