

Container Label

SEE FOLDER IN BOTTOM FLAP FOR USE DIRECTIONS

FORE

FUNGICIDE AND ORNAMENTAL FUNGICIDE




6 1984

KEEP OUT OF REACH OF CHILDREN

CAUTION

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal instructions on back panel and attached use directions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.



**ROHM
AND
HAAS**
PHILADELPHIA, PA 19105

ACTIVE INGREDIENTS

Mancuzeb	80%
A coordination product of zinc ion and manganese ethylene bisdithiocarbamate in which the ingredients are	
Manganese**	16%
Zinc**	2%
Ethylene bisdithiocarbamate ion (C ₂ H ₄ N ₂ S ₂)	62%

INERT INGREDIENTS	20%
TOTAL	100%

EPA Reg. No. 707-87
EPA Est. No. 8590 PA 1
U.S. Patent 3,179,131
© Copyright 1984 Rohm and Haas Company

NET CONTENTS 4 LBS. (1.814 KG)

BEST DOCUMENT AVAILABLE

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

May cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. PROTECTIVE CLOTHING consisting of long pants, long sleeve shirt, gloves, hat and boots must be worn during mixing and loading.

STATEMENTS OF PRACTICAL TREATMENT

- IF SWALLOWED:** Dilute by giving 2 glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person.
- IF INHALED:** Move subject to fresh air.
- IF IN EYES:** Flush eyes with large amounts of water for at least 15 minutes. Consult a physician if irritation persists.
- IF ON SKIN:** Wash affected skin areas with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Crop injury, ineffectiveness or other unintended consequences resulting from such factors as failure to follow label directions, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property, handling, storage and manner of use are beyond the control of Rohm and Haas and Seller. All such risks will be assumed by the Buyer or User. In no case will Rohm and Haas or Seller be held liable for consequential, special or indirect damages resulting from the use or handling of this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **SEE FOLDER IN BOTTOM FLAP FOR USE DIRECTIONS**

STORAGE AND DISPOSAL

STORAGE: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area. Do not allow to become wet or overheated in storage. decomposition, impaired activity or fire may result. Keep container closed when not in use.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or empty container may result in contamination. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED: Sweep or shovel into containers for disposal or reworking. Keep dusting to a minimum. Flush contaminated area with a large amount of water to a chemical or sanitary sewer containing a settling pit.

FORE

TURF AND ORNAMENTAL FUNGICIDE

EPA Reg. No. 707-87
©Copyright 1983
Rohm and Haas Company
U.S. Patent 3,375,616



NOTICE Before using this product read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

May cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. **PROTECTIVE CLOTHING** consisting of long pants, long sleeve shirt, gloves, hat and boots must be worn during mixing and loading.

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Dilute by giving 2 glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person.

IF INHALED: Move subject to fresh air.

IF IN EYES: Flush eyes with large amounts of water for at least 15 minutes. Consult a physician if irritation persists.

IF ON SKIN: Wash affected skin areas with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Crop injury, ineffectiveness or other unintended consequences resulting from such factors as failure to follow label directions, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property, handling, storage and manner of use are beyond the control of Rohm and Haas and Seller. All such risks will be assumed by the Buyer or User. In no case will Rohm and Haas or Seller be held liable for consequential, special or indirect damages resulting from the use or handling of this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL USE INFORMATION

FORE Turf and Ornamental Fungicide is a unique and different broad spectrum protectant fungicide which when used according to label directions provides high levels of disease control. This fungicide is safe for use on turf grasses and most ornamentals.

Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program. The addition of a spreader/sticker will improve fungicide performance by providing a uniform spray deposit, increased foliar redistribution, and fungicide retention during periods of wet weather.

USE RATE DETERMINATION

Carefully read, understand, and follow label rates and use limitations.

Under low disease conditions, minimum label rates per application can be used while maximum label rates and shortened spray schedules are recommended for severe or threatening disease conditions.

For proper application to turf, determine the square footage to be treated, divide footage by 1000 and multiply by the recommended fungicide use rate per 1000 sq. ft., and then determine the amount of water required to provide adequate coverage. When treating ornamentals, determine the recommended fungicide use rate and the spray gallonage required to provide a thorough coverage. Prepare only the amount of spray solution required to treat the desired area.

When FORE fungicide is applied by a hand sprayer, 1 pound per 100 gallons of acre is equivalent to 1 level tablespoon per gallon spray solution.

MIXING

Slowly place into spray tank as it is being filled or thoroughly pre-mix in a nurse tank for concentrate sprayers. Add other co-applied fungicides, insecticides, growth regulators, micronutrients, and spray adjuvants after FORE fungicide has been placed into suspension.

When preparing spray solutions for use in a hand sprayer, pre-mix as a slurry in a small container and then add to sprayer containing 1/2 to 1/3 the desired final water volume.

COMPATIBILITY

FORE fungicide is compatible with most commonly used agricultural fungicides, insecticides, and growth regulators. When preparing tank mixes, use should consult spray compatibility charts or State Extension Service Specialists prior to actual use.

The physical spray compatibility of specific tank mixes can be checked by preparing a sample suspension in a glass container and observing the suspension for any adverse effects which might interfere with proper spray application.

BEST DOCUMENT AVAILABLE

MAR 6 1984

Under the Federal Insecticide, Fungicide, and Rodenticide Act, it is recommended, for the pesticide registered under EPA Reg. No.

SPRAY ADJUVANTS

The addition of agricultural surfactants to FORE fungicide sprays will improve initial spray deposits, fungicide redistribution, and weatherability. The following spray adjuvants have been especially formulated to optimize the performance of other applied agricultural chemicals:

- TRITON® CS 7 A spreader binder designed specifically for use in concentrate and low volume sprays applied by aircraft or ground equipment.
- TRITON® B 1956® A water dispersible resin based nonionic surfactant which resists re-wetting and removal by rain. Effective with dilute sprays applied by ground equipment.
- TRITON® AG 98 A low foam nonionic general purpose spray adjuvant. May be used in dilute or concentrate sprays applied by aircraft or ground equipment.

Place FORE fungicide into suspension prior to adding an adjuvant to the spray mixture. Read and carefully observe the precautionary statements and all other information appearing on both product labels prior to spray preparation.

APPLICATION

FORE is a protectant fungicide. Optimum disease control is achieved with thorough coverage sprays applied on a regular basis.

DISEASE SCOUTING

FORE fungicide is a broad spectrum protectant fungicide. If not applied on a routine protectant spray schedule, turf and ornamental plants should be frequently observed for disease signs or symptoms. Fungicide application should be made at the recommended label use rate and spray schedule when disease is first observed, reported in local area or during environmental conditions favorable for disease development.

RESTRICTIONS

Users should carefully read, understand and follow all use restrictions prior to using FORE fungicide.

USE DIRECTIONS

Crop	Diseases Controlled	Rate of FORE Fungicide Per Application	REMARKS (Also Refer to Directions For Use)
TURF	Helminthosporium melting-out Rust (leaf stem stripe)	oz/1000 ft. 4	GENERAL USE DIRECTIONS Start application when grass greens-up in spring or when disease first appears, and repeat at 7 to 14 day intervals or until disease threat is past. When conditions are especially favorable for disease development, apply maximum fungicide use rate on a 7-day spray schedule. Apply in sufficient water to provide adequate coverage. RESTRICTIONS Do not graze treated areas. Do not feed clippings to livestock. Apply on a 7-day spray schedule. Apply at 5 day intervals or more frequently if conditions are especially favorable for disease development. Apply at 2 to 6 week intervals during winter.
	Copper spot fusarium blight Red thread Slime mold	4 to 8	
	Algae	6	
	Dollar spot	6 to 8	
	Rhizoctonia brown patch	4	
	Pythium blight	8	
	Fusarium snow mold	6 to 8	

101 DOCUMENT AVAILABLE

USE DIRECTIONS

Crop	Diseases Controlled	Rate of FORE Fungicide Per Application	REMARKS (Also Refer to Directions For Use)
ORNAMENTALS		1 lb 100 gal	GENERAL USE DIRECTIONS Begin spraying when plants are well leaved or at first sign of disease in a full coverage spray at 7 to 10 day intervals throughout season or follow State Extension Service recommendations for disease control on the ornamental plants listed.
Azalea	Petal blight		Apply in a full coverage spray 2 to 3 times a week while flowers are opening. Direct spray into flowers and thoroughly spray ground under bushes.
Camellias	Petal blight		
Carnation	Leaf spot		
Chrysanthemum	Botrytis petal spot		Apply twice weekly during blooming period
Crabapple (Ornamental)	Cedar apple rust Scab		
Dahlia	Botrytis blight		
Dogwood flowering	Anthracoze		Apply when buds begin to open when bracts have fallen 4 weeks later and again in late summer after flower buds for the next season have formed.
Gladiolus	Curvularia leaf spot Botrytis blossom blight		Make regular weekly applications starting before diseases appear and increase to 2 or 3 applications per week during periods of heavy disease and during rainy weather. On flower spikes reduce spray concentration to 1/2 lb per 100 gals.
Holly	Purple spot		
Hollyhock	Anthracoze Leaf spot Rust		
Honeysuckle	Herpobasidium blight		
Iris	Leaf spot (fire)		
Lily	Botrytis blight		
Pachysandra	Volutella blight		Use a drenching spray of 2 lbs per 50 gallons of water per 5,000 sq ft of bed. Start application at first sign of disease and apply at least 5 applications at 10 to 14 day intervals.
Pansy	Anthracoze		
Peony	Botrytis blossom blight Phytophthora blight		Apply in early spring and early fall; drenching soil around plants as well as the foliage. Promptly destroy all infected plant parts.
Rhododendron	Petal blight		Refer to Azalea
Rose	Black spot		
Snapdragon	Rust		
Tulip	Botrytis blight (fire)		
Zinnia	Leaf blight		

BEST DOCUMENT AVAILABLE

STORAGE AND DISPOSAL

STORAGE Keep away from fire and sparks. Store in a cool, dry, well-ventilated area. Do not allow to become wet or overheated in storage. Decomposition, impaired activity, or fire may result. Keep container closed when not in use.

Pallets of containers should not be stacked more than three high. Provide access aisle for each two rows. Decomposition produces a foul odor. If observed, check for hot containers and immediately remove to open areas for disposal.

PESTICIDE DISPOSAL Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL

Paper Bags — Completely empty bag into application equipment. Then dispose of it in a sanitary landfill or by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Fiber Drums with liners — Do not reuse empty containers. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose drum and liner in a sanitary landfill or by incineration if allowed by State and local authorities.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED
Sweep or shovel into containers for disposal or reworking. Keep dusting to a minimum. Flush contaminated area with a large amount of water to a chemical or sanitary sewer containing a settling pit.

FORE® TRITON® CS? TRITON® B 1956 and TRITON® AG 96
are registered trademarks of Rohm and Haas Company.

8107 B0

MADE IN U.S.A.

BEST DOCUMENT