

that may have germinated during this time. Maximum herbicidal activity from TOK, while dependent upon prevailing temperatures, will be somewhere between 7 and 14 days after flood water has first started across the field.

NOTE: DRAINING OR SPILLING HEAVY WATER FLOW FROM THE FIELD ANYTIME WITHIN 3 WEEKS AFTER THE TOK APPLICATION WILL PROBABLY RESULT IN A SIGNIFICANT REDUCTION OR COMPLETE LOSS OF WEED CONTROL.

DOSAGE AND EQUIPMENT

Apply TOK at 2 pounds active (1 gallon E-25 formulation) to the established flood waters in 10 to 12 gallons of water per acre by aircraft. Nozzling appropriate for phenoxy herbicides will be satisfactory and must be used for the application of TOK.

PLANTING OF RICE

Before seeding rice make certain that a positive water flow has been employed during the previous 24 hour period. Seeding with presoaked or sprouted rice seed should not be done until 2 days after TOK treatment into flood. Grower may now start his usual program on water flow management with consideration of draining or heavy water flow as detailed in NOTE above.

DIRECTIONS FOR USE

PREFLOOD APPLICATION

FIELD MANAGEMENT BEFORE TREATMENT

Field should be level, well worked and essentially free of trash and clods larger than fist size. It is important to break up all clods thus permitting the more uniform distribution of TOK E-25 herbicide and assuring earlier weed seed germination during the period TOK herbicide is most effective.

DOSAGE AND EQUIPMENT

Apply TOK at 3 pounds active (1½ gallons of E-25 formulation) to the bare soil in 10 to 12 gallons of water per acre by aircraft or in 40 to 60 gallons of water per acre by ground equipment. Nozzling appropriate for phenoxy herbicides will be satisfactory and must be used for the application of TOK. DO NOT INCORPORATE TOK INTO THE SOIL.

WATER MANAGEMENT AFTER TREATMENT

As soon as soil is treated with TOK E-25 flooding may begin. The initial flood should not be established so rapidly as to cause soil erosion or excess soil movement within the field. Flooding should be completed to a depth of 2 to 6 inches. As soon as flooding is completed the water flow should be stabilized so that not more than 1 inch of water flows over the last outflow box. Grower preference for the amount of water flow can be resumed 3 days after the reduced flow period.

DRAINING OF THE FIELD ANYTIME WITHIN 3 WEEKS AFTER THE TOK APPLICATION WILL PROBABLY RESULT IN A SIGNIFICANT REDUCTION OR COMPLETE LOSS OF WEED CONTROL.

PLANTING OF RICE

Seeding of presoaked or sprouted rice seed can follow any time after the flood waters have been established.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Those risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops and personal injury.

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be irritating to the skin. Avoid breathing spray mist. In case of contact, flush skin and eyes with plenty of water; for eyes, get medical attention. Harmful if swallowed. If taken internally, induce vomiting. Drink copious quantities of water and obtain medical attention for gastric lavage. Do not store near feed and food products.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on the label.

Rinse equipment and containers and dispose of waste by burying in non-crop lands away from water supply. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

IMPORTANT: DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Protect from freezing. If stored at 20°F or below, warm to 45°F or above and thoroughly mix before using.

ACTIVE INGREDIENT

2, 4-dichlorophenyl p-nitrophenyl ether 25%*

INERT INGREDIENT 75%
100%


*Equivalent to 2 lbs. active ingredient per gallon.

EPA Registration No. 707-88-AA

EPA Est. No. 707-PA-1

Patented in U.S. and Foreign Countries
U.S. Pat. No. 3,000,225

NET CONTENTS



ROHM HAAS

PHILADELPHIA, PA. 19105

ACCEPTED

TOK E-25



DHMAAS

PHILADELPHIA, PA. 19105

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be irritating to the skin. Avoid breathing spray mist. In case of contact, flush skin and eyes with plenty of water; for eyes, get medical attention. Harmful if swallowed. If taken internally, induce vomiting. Drink copious quantities of water and obtain medical attention for gastric lavage. Do not store near feed and food products.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on the label.

Rinse equipment and containers and dispose of waste by burying in non-crop lands away from water supply. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

IMPORTANT: DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Protect from freezing. If stored at 20°F or below, warm to 45°F or above and thoroughly mix before using.

ACTIVE INGREDIENT	
2, 4-dichlorophenyl p-nitrophenyl ether	25%*
INERT INGREDIENT	75%
	<u>100%</u>

*Equivalent to 2 lbs. active ingredient per gallon.
EPA Registration No. 707-88-AA
EPA Est. No. 707-PA-1
Patented in U.S. and Foreign Countries
U.S. Pat. No. 3,080,225

NET CONTENTS

water in 10 to 12 gallons of water per acre by aircraft. Nozzling appropriate for phenoxy herbicides will be satisfactory and must be used for the application of TOK.

PLANTING OF RICE

Before seeding rice make certain that a positive water flow has been employed during the previous 24 hour period. Seeding with presoaked or sprouted rice seed should not be done until 2 days after TOK treatment into flood. Grower may now start his usual program on water flow management with consideration of draining or heavy water flow as detailed in NOTE above.

DIRECTIONS FOR USE

PREFLOOD APPLICATION

FIELD MANAGEMENT BEFORE TREATMENT

Field should be level, well worked and essentially free of trash and clods larger than fist size. It is important to break up all clods thus permitting the more uniform distribution of TOK E-25 herbicide and assuring earlier weed seed germination during the period TOK herbicide is most effective.

DOSAGE AND EQUIPMENT

Apply TOK at 3 pounds active (1½ gallons of E-25 formulation) to the bare soil in 10 to 12 gallons of water per acre by aircraft or in 40 to 60 gallons of water per acre by ground equipment. Nozzling appropriate for phenoxy herbicides will be satisfactory and must be used for the application of TOK. DO NOT INCORPORATE TOK INTO THE SOIL.

WATER MANAGEMENT AFTER TREATMENT

As soon as soil is treated with TOK E-25 flooding may begin. The initial flood should not be established so rapidly as to cause soil erosion or excess soil movement within the field. Flooding should be completed to a depth of 2 to 6 inches. As soon as flooding is completed the water flow should be stabilized so that not more than 1 inch of water flows over the last outflow box. Grower preference for the amount of water flow can be resumed 3 days after the reduced flow period.

DRAINING OF THE FIELD ANYTIME WITHIN 3 WEEKS AFTER THE TOK APPLICATION WILL PROBABLY RESULT IN A SIGNIFICANT REDUCTION OR COMPLETE LOSS OF WEED CONTROL.

PLANTING OF RICE

Seeding of presoaked or sprouted rice seed can follow any time after the flood waters have been established.

NOTICE: Seller warrants that the product conforms to its chemical description, and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Those risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops and personal injury.

ACCEPTED
April 1, 1975
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTRATION

707-88-AA

707-88-AA
11/15/75

TOK WP-50

WETTABLE POWDER
A SELECTIVE
AGRICULTURAL HERBICIDE

CAUTION
KEEP OUT OF REACH OF CHILDREN

May be irritating to the skin. Avoid breathing vapors. In case of contact, flush with plenty of water. For more information, contact your local poison control center or hospital nearest you for treatment.

This product is for use on lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Reseal promptly and store in original container. Keep out of reach of children. Wash hands thoroughly after use. Do not use if you are pregnant or nursing.


IMPORTANT
 Do not apply to lawns or golf courses.

ACTIVE INGREDIENT
 2,4-Dichlorophenoxyacetic acid, 2,4,5-trichlorophenoxyacetic acid

INERT INGREDIENT
 Patent Pending in U.S. and other countries. U.S. Pat. No. 3,088,000

PACKED FOR ROHM AND HAAS COMPANY
 EPA Reg. No. 707-92 AA
 EPA Est. No. 477-MD 1

NET CONTENTS 5 LBS.



ROHM AND HAAS
 PHILADELPHIA, PA 19105

General Information
ORNAMENTALS AND NON CROPS

GRASSES

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

FLOWERS
CARNATIONS

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

CHRYSANTHEMUMS

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

ROSES

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

NON CROP

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

SECTION 2

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

ONIONS

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

MULTIPLE ONIONS

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

CULTURAL CONSIDERATIONS

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

APPLICATION

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

NOTE

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

BROCCOLI, CABBAGE, CAULIFLOWER

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

NOTICE

Apply to lawns, golf courses, and parks. Do not use on ornamentals, shrubs, or trees. Apply when weeds are young. Do not apply to treated areas until the weeds have died.

Handwritten note: 10/10/74