

### General Information ORNAMENTALS AND NON-CROP

#### GROUND COVERS

Apply TOK WP 50 at 1.0 to 1.5 lb. per 1,000 sq. ft. of ground cover. Apply in early spring or late fall. For best results, apply to well-established ground covers.

<i>Asplenium</i>	<i>Polka Dot Plant</i>
<i>Epiphyllum</i>	<i>Spotted Begonia</i>
<i>Fittonia</i>	<i>Wandering Jew</i>
<i>Hemipeltis</i>	<i>Wax Plant</i>
<i>Peperomia</i>	<i>Zebra Plant</i>
<i>Polka Dot Plant</i>	
<i>Spotted Begonia</i>	
<i>Wandering Jew</i>	
<i>Wax Plant</i>	
<i>Zebra Plant</i>	

The water requirement of TOK WP 50 is 4 to 6 pounds of water per acre. Apply to the soil surface. The product is not to be used on the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

#### FLOWERS

**CARNATIONS** Apply TOK WP 50 at 1.0 to 1.5 lb. per 1,000 sq. ft. of ground cover. Apply in early spring or late fall. For best results, apply to well-established ground covers.

**CHRYSANTHEMUMS** Apply TOK WP 50 at 1.0 to 1.5 lb. per 1,000 sq. ft. of ground cover. Apply in early spring or late fall. For best results, apply to well-established ground covers.

**ROSES** Apply TOK WP 50 at 1.0 to 1.5 lb. per 1,000 sq. ft. of ground cover. Apply in early spring or late fall. For best results, apply to well-established ground covers.

#### NON CROP

Apply TOK WP 50 at 1.0 to 1.5 lb. per 1,000 sq. ft. of ground cover. Apply in early spring or late fall. For best results, apply to well-established ground covers.

### DIRECTIONS FOR USE General Information—VEGETABLES

**TOK WP 50** is a contact herbicide. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**DO NOT** apply TOK WP 50 to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**WATER MANAGEMENT** Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**Excessive moisture should be avoided, for if the soil is very wet from rain or irrigation at the time of crop seedling emergence, TOK may injure the crop.**

Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**CULTURAL CONSIDERATIONS** Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**APPLICATION** Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**NOTE** Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**BROCCOLI, CABBAGE, CAULIFLOWER** Apply TOK WP 50 at 1.0 to 1.5 lb. per 1,000 sq. ft. of ground cover. Apply in early spring or late fall. For best results, apply to well-established ground covers.

Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**BRUSSELS SPROUTS** Apply TOK WP 50 at 1.0 to 1.5 lb. per 1,000 sq. ft. of ground cover. Apply in early spring or late fall. For best results, apply to well-established ground covers.

**CAUTION** Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**HORSERADISH** Apply TOK WP 50 at 1.0 to 1.5 lb. per 1,000 sq. ft. of ground cover. Apply in early spring or late fall. For best results, apply to well-established ground covers.

**PREEMERGENCE** Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**POSTEMERGENCE** Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**ONIONS (dry bulb only)** Apply TOK WP 50 at 1.0 to 1.5 lb. per 1,000 sq. ft. of ground cover. Apply in early spring or late fall. For best results, apply to well-established ground covers.

**MULTIPLE APPLICATIONS** Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

**NOTICE** Apply TOK WP 50 to the soil surface. Do not apply to the soil surface. It is a contact herbicide and will kill the weeds by contact. It is not a systemic herbicide and will not be absorbed by the weeds. It is not a soil sterilant and will not prevent the growth of weeds. It is not a soil fumigant and will not prevent the growth of weeds.

Oct 2, 1970

999 99