


**AGRICULTURAL USE ONLY OR ONLY
USE AND STORAGE BY SERVICE PERSONS**



REACH OF CHILDREN

Product see side panel for additional precautionary state-
el text for use directions, use restrictions, storage and
s and limit of warranty and liability statements.

ACTIVE INGREDIENTS

A coordination product of zinc ion
and manganese ethylene bisdithio-
carbamate 72.0%

In which the ingredients are:

Manganese ++	14.4%
Zinc++	1.8%
Ethylene bisdithio- carbamate	55.8%
2,4-Dinitro-6-octyl* phenyl crotonate }	4.4%
2,6-Dinitro-4-octyl* phenyl crotonate }	
Nitrooctyl* phenols (principally dinitro)	0.3%
INERT INGREDIENTS	23.3%
	100.0%

*A mixture of 1-methylheptyl, 1-ethylhexyl, and
1-propylpentyl isomers.

EPA Reg. No. 707-93
EPA Est. No. 707-PA-1
U.S. Patent 3,379,610

©Copyright 1982 Rohm and Haas Company

NET CONTENTS 5 LBS (2.268 KG)

DIKAR has performed satisfactorily in high concentrate sprays in
both ground sprayers and in aircraft.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a man-
DIKAR is a fungicide-miticide preparation for use on apple
is effective for the control of the following diseases and s

Diseases: Apple Scab	Brown Rot
Powdery Mildew	Cedar Apple Rust
Black Rot	Fly Speck
Bitter Rot	Sooty Blotch

Use DIKAR throughout the season in a regular apple spr-
state spray schedule. Supplement with insecticides or o-
DIKAR is believed to be compatible with the various ins-
apples. If emulsifiable formulations of any pesticide are u-
to check compatibility. For further details, see recom-
consult the current State Agricultural Experiment Station
Service Specialists for information on compatibility, dos-

APPLES — Use DIKAR in a regular spray program beginn-
dormant and repeat as sprays are required in accord-
schedule. In natural rainfall areas of high scab incidence,
to 10 days through petal fall and every 10 to 14 days in t-
Use 1 1/2 to 2 pounds of DIKAR per 100 gallons of water in
sprays. Use the higher dosage early in the season espec-
mildew development. The lower dosage may be used in
For concentrate application adjust the mixing rate acco-
pounds of DIKAR per acre for mature trees of aver-
satisfactorily at concentrations of 2X to 3X pound of

NOTE

Do not apply to apples within 21 days of harvest in Ar-
Kansas, Kentucky, Maryland, Missouri, New Jersey, North
Carolina, Tennessee, Virginia and West Virginia, or with-
not specifically listed above.
Do not graze livestock in the treated areas.
Do not use DIKAR within 7 days before or after an oil app-
To avoid uneven, blotchy residues from high volume dilu-
B-1956® per 100 gallons of spray in the last 1 or 2 appl-
applications usually leave a uniform, even deposit.
NOTE: DIKAR has about 24 to 30 hours "backup" or erac-
weather conditions or other delays interfere with the
schedule, DIKAR may be augmented with dodine in or
sprays.
If the normal spray interval is extended to 2 weeks or mor-
especially severe, 1 or 2 sprays of KARATHA, LC o-
between the regular DIKAR applications for added mild-

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or dis-
PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse
to label instructions must be disposed of according to
procedures.
CONTAINER DISPOSAL: Completely empty bag by shak-
loosen clinging particles. Empty residue into application
dispose of them in a sanitary landfill or by incineration if

NOTICE: Seller warrants that the product conforms to its chemi-
purpose stated on the label when used in accordance with direc-
neither this warranty nor any other warranty of merchantability or
implied, extends to the use, storage or handling of this product
instructions, or under abnormal conditions, or under conditions
seller assumes the risk of any such use. These risks include, but
and animals to which the material is applied, failure to control pe-
or crops, and personal injury.

Protective clothing consisting of long pants, long sleeve shirt and boots must be worn during mixing and loading.

INSTRUCTIONS FOR USE

Federal law prohibits the use of this product in a manner inconsistent with its labeling. This is a fungicide-miticide preparation for use on apples and when used as recommended controls the following diseases and suppression of the following mites:

Apple Scab	Brown Rot	Mites: European Red
Powdery Mildew	Cedar Apple Rust	Schoene
Black Rot	Fly Speck	Two-spotted
Fire Blight	Sooty Blotch	Clover

Use throughout the season in a regular apple spray program in accordance with local recommendations. Supplement with insecticides or other fungicides as needed.

When used in conjunction with other pesticides, care should be taken to be compatible with the various insecticides recommended for use on apples. Formulations of any pesticide are used with DIKAR, care should be taken to avoid phytotoxicity. For further details, see recommendations on this label and then consult State Agricultural Experiment Station spray schedules or State Extension Service for information on compatibility, dosage, timing and frequency of spray.

Use DIKAR in a regular spray program beginning early with the green tip or delayed bloom as sprays are required in accordance with the local or state spray schedule. In areas of high scab incidence, sprays are generally required every 7 to 10 days from petal fall and every 10 to 14 days in the cover sprays.

Use 1.5 to 2.0 lbs of DIKAR per 100 gallons of water in dilute, single strength, full coverage sprays. Higher dosage early in the season especially during primary scab and rapid bloom. The lower dosage may be used in cover sprays.

Application rate adjust the mixing rate accordingly. Use approximately 8 to 10 lbs per acre for mature trees of average density. DIKAR has performed well at concentrations of 2X to 3X in ground sprayers and up to 70X in aircraft.

Use on apples within 21 days of harvest in Arkansas, Delaware, Illinois, Indiana, Maryland, Missouri, New Jersey, North Carolina, Ohio, Pennsylvania, South Carolina, Virginia and West Virginia, or within 30 days of harvest in other states listed above.

Use in the treated areas.

Do not apply within 7 days before or after an oil application.

Sticky residues from high volume dilute sprays add 2 to 3 ounces TRITON® per gallon of spray in the last 1 or 2 applications before harvest. Concentrate sprays should leave a uniform, even deposit.

For about 24 to 30 hours "backup" or eradication action against apple scab. If rains or other delays interfere with the normal timing of a protective scab spray, it may be augmented with dodine in one or more combination or separate applications.

If the spray interval is extended to 2 weeks or more and powdery mildew and mites are present, 1 or 2 sprays of KARATHANE® LC or KARATHANE® WD should be used in conjunction with DIKAR applications for added mildew or mite control.

DISPOSAL

Do not use for water, food or feed by storage or disposal. Open dumping is prohibited.

WATER: Pesticide, spray mixture or rinse water that cannot be used according to label must be disposed of according to applicable Federal, State or local regulations.

LANDFILL: Completely empty bag by shaking and tapping sides and bottom to dislodge contents. Empty residue into application equipment. Perforate bags and then dispose in a sanitary landfill or by incineration if allowed by State or local authorities.

This product conforms to its chemical description and is reasonably fit for the label when used in accordance with directions under normal conditions of use, but no warranty is made for any other warranty of merchantability or fitness for a particular purpose, express or implied, or for use, storage or handling of this product in a manner other than as directed by label directions, or under conditions not reasonably foreseeable to seller, and for any such use. These risks include, but are not limited to, damage to plants, crops or property if material is applied, failure to control pests, damage caused by drift to other plants or property.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through the skin or inhaled. Causes moderate eye, skin, nose and throat irritation. Avoid contact with skin, eyes and clothing. Avoid breathing dust or spray mist.

Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Call a physician. Do not induce vomiting or give anything by mouth to an unconscious person.


IF INHALED: Remove victim to fresh air. Treat the person symptomatically. If irritation persists, call a physician.

IF ON SKIN: Wash exposed areas with plenty of soap and water. If irritation persists, call a physician.

IF IN EYES: Flush eyes with plenty of water for at least 15 minutes. If irritation persists, call a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Apply this product only as specified on this label.



MADE IN U.S.A.