

STAM[®] M-3

3 LB. E.C.

FOR POST-EMERGENCE CONTROL OF
BARNYARDGRASS AND OTHER WEEDS IN SOUTHERN RICE

WARNING KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED or absorbed through skin. Causes severe eye irritation. May cause skin irritation. Avoid contact with eyes, skin or clothing. Avoid prolonged breathing of vapor or mist. In case of contact, flush skin or eyes with plenty of water for at least 15 minutes; for eyes, get medical attention.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not reuse container. Destroy when empty.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

ACTIVE INGREDIENT
3-(4-dimethylamino)propylamine 55%
INERT INGREDIENTS 65%
Total 100
Equivalent to 3.33 lb. active ingredient per gallon.
PATENT PENDING
EPA Reg. No. 707-110-AA

NET CONTENTS
30 GAL.

DIRECTIONS FOR USE—RICE ONLY

Apply this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

NOTE: When this product is applied, especially after the fourth leaf stage of growth of the rice, under some conditions, a severe injury to the rice may result. However, the rice plants usually outgrow such injury when caused by the later application.

USE IN CALIFORNIA: None. When used in California, apply as directed.


NOTE: Before use, read entire label.

To control watergrass, apply this product 10 to 40 days after planting before the rice is 10 days old. When watergrass is present in rice fields, apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

When watergrass is present in rice fields, apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

If watergrass is present after the rice has emerged, apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged. Apply 1 to 4 quarts per acre of this product to rice fields after the rice has emerged.

NOTICE: Seller warrants that the product conforms to its label and is reasonably fit for the purpose stated on the label when used in accordance with directions and normal application of use. Seller does not warrant, and other warranties, if any, are hereby disclaimed, that the product will perform as advertised. The use of this product may result in injury to the rice plants, and the use of this product may result in injury to the rice plants. The use of this product may result in injury to the rice plants. The use of this product may result in injury to the rice plants.



**ROHM
AND
HAAS**

PHILADELPHIA, PA. 19105

STAM[®] M-4

4 LB. E.C.


FOR POST-EMERGENCE CONTROL OF
BARNYUM GRASSES AND OTHER WEEDS IN RICE

WARNING
KEEP OUT OF REACH OF CHILDREN

STAM M-4 is a selective herbicide for the control of barnyard grasses and other weeds in rice. It is applied as a foliar spray to the weeds in the rice field. The active ingredient is bentazone. It is a systemic herbicide which kills the weeds by interfering with their photosynthesis. It is safe for rice and other crops when used as directed. It is also safe for humans and animals when used as directed. It is a non-flammable, non-toxic, and non-corrosive herbicide. It is a selective herbicide which kills the weeds in the rice field. It is safe for rice and other crops when used as directed. It is also safe for humans and animals when used as directed. It is a non-flammable, non-toxic, and non-corrosive herbicide.

NET WEIGHT
4 LB. E.C.

30



**ROHM
HAAS**

